



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:14 PM GMT

PDB ID : 1SQN  
Title : Progesterone Receptor Ligand Binding Domain with bound Norethindrone  
Authors : Williams, S.P.; Madauss, K.P.; Deng, J.-S.; Austin, R.J.H.; Lambert, M.H.;  
McLay, I.; Pritchard, J.; Short, S.A.; Stewart, E.L.; Uings, I.J.  
Deposited on : 2004-03-19  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1278 (1.48-1.44)
Clashscore	102246	1336 (1.48-1.44)
Ramachandran outliers	100387	1320 (1.48-1.44)
Sidechain outliers	100360	1320 (1.48-1.44)
RSRZ outliers	91569	1279 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	 6% 79% 15% • 5%
1	B	261	 3% 80% 15% • •

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called progesterone receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1976	1284	323	355	14			
1	B	250	Total	C	N	O	S	0	0	0
			1991	1290	330	357	14			

- Molecule 2 is (14BETA,17ALPHA)-17-ETHYNYL-17-HYDROXYESTR-4-EN-3-ONE (three-letter code: NDR) (formula:  $C_{20}H_{26}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			22	20	2		
2	B	1	Total	C	O	0	0
			22	20	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	126	Total 126	O 126	0	0
3	B	150	Total 150	O 150	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: progesterone receptor



- Molecule 1: progesterone receptor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.75Å 64.14Å 70.14Å 90.00° 95.84° 90.00°	Depositor
Resolution (Å)	19.80 – 1.45 37.78 – 1.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.80-1.45) 76.3 (37.78-1.45)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.17 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, $R_{free}$	0.186 , 0.216 0.189 , 0.186	Depositor DCC
$R_{free}$ test set	4847 reflections (7.59%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtriage
Anisotropy	0.556	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 68787 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4287	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NDR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.27	8/2017 (0.4%)	1.12	8/2729 (0.3%)
1	B	1.28	6/2033 (0.3%)	1.15	8/2753 (0.3%)
All	All	1.28	14/4050 (0.3%)	1.13	16/5482 (0.3%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	808	PHE	CD2-CE2	7.57	1.54	1.39
1	B	720	GLN	CG-CD	7.07	1.67	1.51
1	B	777	TYR	CG-CD2	6.86	1.48	1.39
1	A	772	SER	CA-CB	6.19	1.62	1.52
1	A	692	MET	SD-CE	-6.10	1.43	1.77
1	A	765	TRP	CE3-CZ3	6.08	1.48	1.38
1	A	698	VAL	CB-CG1	-6.00	1.40	1.52
1	B	848	TYR	CD2-CE2	5.79	1.48	1.39
1	A	721	LEU	CG-CD2	5.45	1.72	1.51
1	B	898	SER	CB-OG	-5.40	1.35	1.42
1	B	845	ARG	NE-CZ	-5.30	1.26	1.33
1	A	745	ASP	CB-CG	-5.26	1.40	1.51
1	B	728	SER	CB-OG	-5.09	1.35	1.42
1	A	775	MET	CG-SD	-5.02	1.68	1.81

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	869	ARG	NE-CZ-NH2	-11.55	114.52	120.30
1	B	869	ARG	NE-CZ-NH1	10.02	125.31	120.30
1	A	869	ARG	NE-CZ-NH1	7.85	124.22	120.30
1	B	878	ASP	CB-CG-OD2	7.69	125.22	118.30
1	A	825	LEU	CB-CG-CD2	7.34	123.48	111.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	880	LEU	CB-CG-CD1	-6.85	99.36	111.00
1	B	869	ARG	NE-CZ-NH2	-6.83	116.89	120.30
1	B	850	ARG	NE-CZ-NH1	6.57	123.58	120.30
1	A	740	ARG	NE-CZ-NH2	-6.56	117.02	120.30
1	A	697	ASP	CB-CG-OD2	6.32	123.98	118.30
1	A	753	TYR	CB-CG-CD2	-6.14	117.31	121.00
1	B	869	ARG	CD-NE-CZ	6.02	132.03	123.60
1	B	753	TYR	CB-CG-CD1	5.76	124.46	121.00
1	B	768	TYR	CB-CG-CD2	-5.42	117.75	121.00
1	A	745	ASP	CB-CA-C	-5.08	100.25	110.40
1	B	753	TYR	CB-CG-CD2	-5.03	117.98	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1976	0	2003	30	2
1	B	1991	0	2009	27	4
2	A	22	0	23	0	0
2	B	22	0	23	0	0
3	A	126	0	0	12	6
3	B	150	0	0	5	4
All	All	4287	0	4058	57	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:692:MET:HE3	3:A:265:HOH:O	1.54	1.07
1:B:775:MET:HE3	1:B:785:ASN:HB3	1.08	1.03
1:B:775:MET:CE	1:B:785:ASN:HB3	1.87	1.03

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:864:VAL:HG12	1:A:868:GLN:HE21	1.28	0.95
1:A:909:MET:SD	3:A:212:HOH:O	2.27	0.91
1:A:909:MET:CG	3:A:212:HOH:O	2.26	0.82
1:B:707:LYS:CB	3:B:70:HOH:O	2.28	0.80
1:A:698:VAL:HG23	3:A:95:HOH:O	1.81	0.79
1:A:692:MET:CE	3:A:265:HOH:O	2.20	0.77
1:B:723:GLU:O	1:B:727:LEU:HD13	1.86	0.75
1:A:899:ARG:HG3	1:A:899:ARG:HH11	1.51	0.74
1:B:775:MET:HE3	1:B:785:ASN:CB	2.03	0.74
1:B:703:HIS:HD2	1:B:717:SER:OG	1.74	0.71
1:A:698:VAL:CG2	3:A:95:HOH:O	2.36	0.71
1:A:899:ARG:HG3	1:A:899:ARG:NH1	2.08	0.65
1:B:845:ARG:NH1	3:B:94:HOH:O	2.34	0.60
1:B:743:HIS:HE2	1:B:836:ARG:HH11	1.48	0.59
1:B:888:HIS:CD2	1:B:917:LEU:HD21	2.38	0.59
1:A:864:VAL:HG12	1:A:868:GLN:NE2	2.09	0.57
1:A:703:HIS:CE1	1:A:705:ASN:HB3	2.42	0.55
1:B:817:GLU:OE1	1:B:869:ARG:HD3	2.07	0.55
1:A:888:HIS:CD2	1:A:917:LEU:HD21	2.42	0.54
1:A:909:MET:HG3	3:A:212:HOH:O	1.99	0.53
1:B:775:MET:CE	1:B:785:ASN:CB	2.76	0.52
1:A:919:LYS:NZ	3:A:193:HOH:O	2.43	0.52
1:A:696:PRO:HB3	1:A:728:SER:OG	2.10	0.51
1:A:881:HIS:HE1	1:A:928:LEU:O	1.94	0.51
1:A:729:VAL:HG12	1:A:751:ILE:HD12	1.94	0.50
1:A:723:GLU:CD	1:A:906:PRO:HB2	2.33	0.49
1:B:703:HIS:CD2	1:B:717:SER:OG	2.62	0.48
1:B:743:HIS:HE2	1:B:836:ARG:NH1	2.11	0.48
1:B:703:HIS:HE1	3:B:100:HOH:O	1.97	0.47
1:A:909:MET:CB	3:A:212:HOH:O	2.61	0.47
1:B:840:GLN:HG3	3:B:20:HOH:O	2.14	0.47
1:B:784:LEU:HD22	1:B:788:ARG:HD3	1.97	0.47
1:A:787:GLN:HG3	3:A:202:HOH:O	2.14	0.46
1:B:723:GLU:CD	1:B:906:PRO:HB2	2.37	0.45
1:B:692:MET:O	1:B:692:MET:HE3	2.16	0.45
1:A:899:ARG:HG2	3:A:164:HOH:O	2.17	0.45
1:A:692:MET:HE1	1:A:819:LEU:HD11	1.99	0.45
1:A:864:VAL:O	1:A:868:GLN:HG3	2.16	0.45
1:A:802:TRP:O	1:A:806:GLN:HG3	2.17	0.44
1:B:743:HIS:NE2	1:B:836:ARG:NH1	2.61	0.44
1:B:811:LEU:HB3	1:B:869:ARG:NH2	2.33	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:863:VAL:HG23	1:B:864:VAL:N	2.33	0.43
1:A:909:MET:HB2	3:A:212:HOH:O	2.19	0.43
1:B:760:VAL:HG22	1:B:804:ILE:HD12	2.01	0.43
1:A:704:ASP:O	1:A:705:ASN:C	2.57	0.42
1:A:704:ASP:C	1:A:705:ASN:OD1	2.58	0.42
1:A:685:PRO:HA	1:A:686:PRO:HD3	1.80	0.42
1:B:714:LEU:HD12	1:B:714:LEU:HA	1.87	0.42
1:B:811:LEU:O	1:B:869:ARG:NH2	2.45	0.42
1:B:917:LEU:HA	1:B:917:LEU:HD23	1.86	0.42
1:A:862:GLY:O	1:A:866:SER:HB3	2.20	0.41
1:B:863:VAL:HG22	3:B:157:HOH:O	2.21	0.41
1:A:917:LEU:HD23	1:A:917:LEU:HA	1.89	0.40
1:B:856:ILE:HG12	1:B:869:ARG:HG2	2.02	0.40

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:85:HOH:O	3:A:256:HOH:O[2_556]	0.03	2.17
3:A:86:HOH:O	3:A:259:HOH:O[2_556]	0.04	2.16
3:A:283:HOH:O	3:B:123:HOH:O[1_556]	0.72	1.48
1:B:847:SER:CB	3:A:257:HOH:O[1_554]	0.90	1.30
1:A:787:GLN:NE2	3:B:277:HOH:O[2_545]	1.37	0.83
1:B:847:SER:OG	3:A:257:HOH:O[1_554]	1.60	0.60
1:B:847:SER:CA	3:A:257:HOH:O[1_554]	1.94	0.26
1:A:854:LYS:CE	3:B:262:HOH:O[1_556]	1.95	0.25
1:B:904:GLU:OE1	3:B:272:HOH:O[2_555]	2.19	0.01

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/261 (93%)	236 (98%)	6 (2%)	0	100	100
1	B	246/261 (94%)	239 (97%)	7 (3%)	0	100	100
All	All	488/522 (94%)	475 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/243 (92%)	219 (98%)	4 (2%)	66	30
1	B	223/243 (92%)	219 (98%)	4 (2%)	66	30
All	All	446/486 (92%)	438 (98%)	8 (2%)	66	30

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	698	VAL
1	A	791	GLU
1	A	802	TRP
1	A	867	SER
1	B	805	PRO
1	B	815	GLN
1	B	882	ASP
1	B	909	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	689	ASN
1	A	725	GLN
1	A	741	ASN
1	A	743	HIS
1	A	812	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	860	GLN
1	A	868	GLN
1	A	881	HIS
1	B	689	ASN
1	B	703	HIS
1	B	705	ASN
1	B	720	GLN
1	B	725	GLN
1	B	741	ASN
1	B	747	GLN
1	B	803	GLN
1	B	815	GLN
1	B	881	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NDR	A	1001	-	25,25,25	3.05	9 (36%)	40,40,40	3.00	20 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NDR	B	2001	-	25,25,25	3.21	8 (32%)	40,40,40	3.22	18 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDR	A	1001	-	-	0/0/60/60	0/4/4/4
2	NDR	B	2001	-	-	0/0/60/60	0/4/4/4

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	NDR	C3-C8	-7.63	1.41	1.54
2	B	2001	NDR	C3-C8	-6.80	1.42	1.54
2	B	2001	NDR	C14-C17	-6.47	1.41	1.52
2	A	1001	NDR	C14-C17	-5.72	1.42	1.52
2	B	2001	NDR	C14-C8	-5.69	1.44	1.54
2	B	2001	NDR	C3-C1	-5.00	1.43	1.53
2	A	1001	NDR	C3-C1	-4.80	1.44	1.53
2	A	1001	NDR	C14-C8	-4.63	1.46	1.54
2	B	2001	NDR	C2-C5	-2.88	1.52	1.56
2	B	2001	NDR	O22-C21	-2.70	1.18	1.23
2	A	1001	NDR	C6-C2	-2.57	1.49	1.54
2	A	1001	NDR	C20-C21	2.04	1.54	1.49
2	A	1001	NDR	C6-C13	2.71	1.59	1.53
2	B	2001	NDR	C19-C17	5.51	1.44	1.34
2	A	1001	NDR	C19-C17	5.59	1.44	1.34
2	B	2001	NDR	C11-C16	6.21	1.28	1.18
2	A	1001	NDR	C11-C16	6.32	1.28	1.18

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	NDR	C17-C19-C21	-5.17	114.93	123.06
2	A	1001	NDR	C7-C2-C6	-4.02	103.90	109.80
2	B	2001	NDR	C15-C17-C19	-3.80	113.98	121.53
2	A	1001	NDR	C15-C17-C19	-3.53	114.53	121.53
2	B	2001	NDR	O22-C21-C19	-3.51	116.17	121.62
2	A	1001	NDR	C18-C20-C21	-3.45	106.02	111.66
2	B	2001	NDR	C10-C5-C11	-3.16	106.37	110.23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	NDR	C17-C19-C21	-2.89	118.52	123.06
2	B	2001	NDR	C5-C11-C16	-2.67	173.49	177.12
2	A	1001	NDR	O12-C5-C11	-2.42	105.59	108.59
2	A	1001	NDR	O22-C21-C19	-2.42	117.86	121.62
2	B	2001	NDR	C6-C2-C1	-2.29	105.80	108.32
2	A	1001	NDR	C15-C9-C3	2.04	115.08	111.67
2	B	2001	NDR	C10-C5-C2	2.06	104.30	102.91
2	A	1001	NDR	C13-C8-C14	2.07	114.50	111.91
2	A	1001	NDR	C20-C18-C14	2.21	116.14	111.51
2	A	1001	NDR	C13-C8-C3	2.31	115.85	111.19
2	B	2001	NDR	C4-C1-C2	2.31	106.74	103.75
2	A	1001	NDR	C18-C14-C8	2.55	118.05	111.87
2	A	1001	NDR	C7-C2-C5	2.65	110.38	108.09
2	B	2001	NDR	C13-C8-C14	2.92	115.56	111.91
2	A	1001	NDR	C6-C2-C5	2.95	119.35	116.93
2	B	2001	NDR	C20-C21-C19	2.95	121.24	116.70
2	A	1001	NDR	C20-C21-C19	2.99	121.31	116.70
2	B	2001	NDR	C8-C14-C17	3.44	117.57	110.40
2	A	1001	NDR	C15-C17-C14	3.66	121.28	115.72
2	A	1001	NDR	C2-C1-C3	3.79	117.54	113.57
2	B	2001	NDR	C4-C1-C3	4.10	125.50	119.03
2	B	2001	NDR	C6-C2-C5	4.50	120.61	116.93
2	A	1001	NDR	C5-C2-C1	4.52	102.95	100.00
2	B	2001	NDR	C5-C2-C1	4.80	103.14	100.00
2	B	2001	NDR	C14-C17-C19	5.12	128.93	122.47
2	A	1001	NDR	C4-C1-C3	6.18	128.78	119.03
2	B	2001	NDR	C8-C3-C1	6.75	121.38	109.65
2	B	2001	NDR	C2-C1-C3	6.78	120.67	113.57
2	A	1001	NDR	C8-C3-C1	7.67	122.98	109.65
2	A	1001	NDR	C14-C8-C3	9.01	122.66	111.88
2	B	2001	NDR	C14-C8-C3	9.44	123.16	111.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	248/261 (95%)	0.09	16 (6%) 22 22	17, 23, 42, 57	0
1	B	250/261 (95%)	-0.11	7 (2%) 56 57	16, 23, 36, 51	0
All	All	498/522 (95%)	-0.01	23 (4%) 36 36	16, 23, 37, 57	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	683	LEU	7.5
1	A	864	VAL	5.9
1	A	705	ASN	5.1
1	B	864	VAL	4.8
1	A	682	GLN	4.5
1	B	863	VAL	4.4
1	A	708	PRO	4.1
1	A	860	GLN	4.0
1	B	860	GLN	3.7
1	A	862	GLY	3.7
1	B	706	THR	3.6
1	A	932	LYS	3.4
1	B	933	LYS	3.4
1	A	704	ASP	3.3
1	A	865	SER	2.9
1	A	863	VAL	2.8
1	A	899	ARG	2.8
1	A	871	TYR	2.5
1	B	862	GLY	2.3
1	A	859	ARG	2.3
1	A	900	ALA	2.3
1	B	705	ASN	2.1
1	A	709	ASP	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NDR	B	2001	22/22	0.95	0.07	-0.13	16,19,21,26	0
2	NDR	A	1001	22/22	0.95	0.06	-0.85	15,18,20,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.