



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:10 PM GMT

PDB ID : 3T22  
Title : Crystal structure of OxyR mutant from Porphyromonas gingivalis  
Authors : Svintradze, D.V.; Wright, H.T.; Lewis, J.P.  
Deposited on : 2011-07-22  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	
1	B	223	
1	C	223	
1	D	223	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Redox-sensitive transcriptional activator OxyR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1683	1070	296	307	10			
1	B	207	Total	C	N	O	S	0	0	0
			1661	1057	292	303	9			
1	C	214	Total	C	N	O	S	0	0	0
			1711	1088	301	312	10			
1	D	208	Total	C	N	O	S	0	0	0
			1670	1063	291	306	10			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	SER	CYS	ENGINEERED MUTATION	UNP Q20K61
A	309	HIS	-	EXPRESSION TAG	UNP Q20K61
A	310	HIS	-	EXPRESSION TAG	UNP Q20K61
A	311	HIS	-	EXPRESSION TAG	UNP Q20K61
A	312	HIS	-	EXPRESSION TAG	UNP Q20K61
B	199	SER	CYS	ENGINEERED MUTATION	UNP Q20K61
B	309	HIS	-	EXPRESSION TAG	UNP Q20K61
B	310	HIS	-	EXPRESSION TAG	UNP Q20K61
B	311	HIS	-	EXPRESSION TAG	UNP Q20K61
B	312	HIS	-	EXPRESSION TAG	UNP Q20K61
C	199	SER	CYS	ENGINEERED MUTATION	UNP Q20K61
C	309	HIS	-	EXPRESSION TAG	UNP Q20K61
C	310	HIS	-	EXPRESSION TAG	UNP Q20K61
C	311	HIS	-	EXPRESSION TAG	UNP Q20K61
C	312	HIS	-	EXPRESSION TAG	UNP Q20K61
D	199	SER	CYS	ENGINEERED MUTATION	UNP Q20K61
D	309	HIS	-	EXPRESSION TAG	UNP Q20K61
D	310	HIS	-	EXPRESSION TAG	UNP Q20K61
D	311	HIS	-	EXPRESSION TAG	UNP Q20K61
D	312	HIS	-	EXPRESSION TAG	UNP Q20K61


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	106	Total 106	O 106	0	0
2	B	78	Total 78	O 78	0	0
2	C	102	Total 102	O 102	0	0
2	D	92	Total 92	O 92	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Redox-sensitive transcriptional activator OxyR



- Molecule 1: Redox-sensitive transcriptional activator OxyR




- Molecule 1: Redox-sensitive transcriptional activator OxyR



- Molecule 1: Redox-sensitive transcriptional activator OxyR





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.21Å 64.36Å 81.31Å 111.75° 90.56° 111.24°	Depositor
Resolution (Å)	18.36 – 2.20 23.24 – 1.90	Depositor EDS
% Data completeness (in resolution range)	90.6 (18.36-2.20) 76.3 (23.24-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.15 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, $R_{free}$	0.234 , 0.274 0.242 , 0.272	Depositor DCC
$R_{free}$ test set	2296 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	0 of 69444 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7103	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/1716	0.56	0/2318
1	B	0.41	0/1694	0.57	0/2289
1	C	0.44	0/1744	0.57	0/2355
1	D	0.45	0/1703	0.59	0/2302
All	All	0.43	0/6857	0.57	0/9264

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1683	0	1703	45	0
1	B	1661	0	1681	65	0
1	C	1711	0	1737	30	0
1	D	1670	0	1689	68	0
2	A	106	0	0	3	0
2	B	78	0	0	10	0
2	C	102	0	0	5	0
2	D	92	0	0	6	0
All	All	7103	0	6810	205	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.



The worst 5 of 205 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:250:SER:HB2	1:D:251:PRO:HD2	1.38	1.05
1:A:204:LEU:HG	1:A:210:MET:CE	1.96	0.95
1:C:300:LYS:HE2	2:C:338:HOH:O	1.72	0.90
1:D:242:PRO:O	1:D:245:THR:HG22	1.75	0.86
1:D:268:GLU:HG2	1:D:307:LEU:HD21	1.59	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/223 (92%)	199 (97%)	7 (3%)	0	100	100
1	B	203/223 (91%)	192 (95%)	11 (5%)	0	100	100
1	C	210/223 (94%)	204 (97%)	6 (3%)	0	100	100
1	D	204/223 (92%)	186 (91%)	17 (8%)	1 (0%)	34	35
All	All	823/892 (92%)	781 (95%)	41 (5%)	1 (0%)	56	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	280	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/197 (94%)	177 (96%)	8 (4%)	35	43
1	B	183/197 (93%)	170 (93%)	13 (7%)	18	19
1	C	188/197 (95%)	184 (98%)	4 (2%)	61	74
1	D	185/197 (94%)	175 (95%)	10 (5%)	27	31
All	All	741/788 (94%)	706 (95%)	35 (5%)	32	39

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	221	SER
1	B	286	LEU
1	D	220	TYR
1	B	245	THR
1	B	276	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	94	ASN
1	B	275	GLN
1	B	302	GLN
1	D	209	GLN
1	D	306	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	210/223 (94%)	0.10	13 (6%) 24 23	24, 36, 53, 62	0
1	B	207/223 (92%)	0.27	14 (6%) 20 20	25, 38, 54, 63	0
1	C	214/223 (95%)	0.05	10 (4%) 35 34	21, 37, 54, 63	0
1	D	208/223 (93%)	0.30	19 (9%) 11 11	22, 40, 56, 68	0
All	All	839/892 (94%)	0.18	56 (6%) 21 20	21, 38, 55, 68	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	145	ILE	4.1
1	A	209	GLN	3.8
1	B	145	ILE	3.7
1	D	123	HIS	3.7
1	D	306	HIS	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.