



## wwPDB/EMDataBank EM Map/Model Validation Summary Report ⓘ

Nov 21, 2016 – 06:47 PM EST

PDB ID : 5T4D  
EMDB ID: : EMD-8354  
Title : Cryo-EM structure of Polycystic Kidney Disease protein 2 (PKD2), residues 198-703  
Authors : Shen, P.S.; Yang, X.; DeCaen, P.G.; Liu, X.; Bulkley, D.; Clapham, D.E.; Cao, E.  
Deposited on : 2016-08-29  
Resolution : 3.00 Å(reported)

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report for a publicly released PDB/EMDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
EM map analysis : **NOT EXECUTED**  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028320


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924
Ramachandran outliers	111179	726
Sidechain outliers	111093	686

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	510	66% 25% 9%
1	B	510	66% 24% 9%
1	C	510	66% 25% 9%
1	D	510	66% 24% 9%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 14876 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called hPKD:198-703, Polycystin-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	463	Total	C	N	O	S	0	0
			3677	2441	587	631	18		
1	B	463	Total	C	N	O	S	0	0
			3677	2441	587	631	18		
1	C	463	Total	C	N	O	S	0	0
			3677	2441	587	631	18		
1	D	463	Total	C	N	O	S	0	0
			3677	2441	587	631	18		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	193	GLY	-	expression tag	UNP Q13563
A	194	ALA	-	expression tag	UNP Q13563
A	195	MET	-	expression tag	UNP Q13563
A	196	GLY	-	expression tag	UNP Q13563
A	197	SER	-	expression tag	UNP Q13563
B	193	GLY	-	expression tag	UNP Q13563
B	194	ALA	-	expression tag	UNP Q13563
B	195	MET	-	expression tag	UNP Q13563
B	196	GLY	-	expression tag	UNP Q13563
B	197	SER	-	expression tag	UNP Q13563
C	193	GLY	-	expression tag	UNP Q13563
C	194	ALA	-	expression tag	UNP Q13563
C	195	MET	-	expression tag	UNP Q13563
C	196	GLY	-	expression tag	UNP Q13563
C	197	SER	-	expression tag	UNP Q13563
D	193	GLY	-	expression tag	UNP Q13563
D	194	ALA	-	expression tag	UNP Q13563
D	195	MET	-	expression tag	UNP Q13563
D	196	GLY	-	expression tag	UNP Q13563
D	197	SER	-	expression tag	UNP Q13563

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	N	O	0
			42	24	3	15	
2	A	1	Total	C	N	O	0
			42	24	3	15	
2	A	1	Total	C	N	O	0
			42	24	3	15	
2	B	1	Total	C	N	O	0
			42	24	3	15	
2	B	1	Total	C	N	O	0
			42	24	3	15	
2	B	1	Total	C	N	O	0
			42	24	3	15	
2	C	1	Total	C	N	O	0
			42	24	3	15	
2	C	1	Total	C	N	O	0
			42	24	3	15	
2	C	1	Total	C	N	O	0
			42	24	3	15	
2	D	1	Total	C	N	O	0
			42	24	3	15	
2	D	1	Total	C	N	O	0
			42	24	3	15	
2	D	1	Total	C	N	O	0
			42	24	3	15	


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: hPKD:198-703, Polycystin-2




- Molecule 1: hPKD:198-703, Polycystin-2




- Molecule 1: hPKD:198-703, Polycystin-2

Chain C: 



• Molecule 1: hPKD:198-703, Polycystin-2

Chain D: 



## 4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of particles used	93805	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Not provided	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	Not provided	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	41132	Depositor
Image detector	Not provided	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >2	RMSZ	# Z  >2
1	A	0.55	0/3774	0.51	0/5139
1	B	0.55	0/3774	0.51	0/5139
1	C	0.55	0/3774	0.51	0/5139
1	D	0.55	0/3774	0.51	0/5139
All	All	0.55	0/15096	0.51	0/20556

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	432	ASN	Peptide
1	B	432	ASN	Peptide
1	C	432	ASN	Peptide
1	D	432	ASN	Peptide



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3677	0	3556	104	0
1	B	3677	0	3556	104	0
1	C	3677	0	3556	103	0
1	D	3677	0	3556	97	0
2	A	42	0	39	1	0
2	B	42	0	39	1	0
2	C	42	0	39	1	0
2	D	42	0	39	1	0
All	All	14876	0	14380	360	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 360 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:396:SER:H	1:D:402:THR:HG22	1.42	0.85
1:A:396:SER:H	1:A:402:THR:HG22	1.42	0.83
1:C:396:SER:H	1:C:402:THR:HG22	1.42	0.83
1:B:396:SER:H	1:B:402:THR:HG22	1.42	0.81
1:A:677:LEU:HD23	1:D:674:ASN:HB3	1.64	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/510 (90%)	437 (96%)	20 (4%)	0	100	100
1	B	457/510 (90%)	437 (96%)	20 (4%)	0	100	100
1	C	457/510 (90%)	437 (96%)	20 (4%)	0	100	100
1	D	457/510 (90%)	437 (96%)	20 (4%)	0	100	100
All	All	1828/2040 (90%)	1748 (96%)	80 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/459 (83%)	378 (100%)	1 (0%)	94	98
1	B	379/459 (83%)	378 (100%)	1 (0%)	94	98
1	C	379/459 (83%)	378 (100%)	1 (0%)	94	98
1	D	379/459 (83%)	378 (100%)	1 (0%)	94	98
All	All	1516/1836 (83%)	1512 (100%)	4 (0%)	95	98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	630	GLN
1	B	630	GLN
1	C	630	GLN
1	D	630	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	636	GLN
1	C	458	GLN
1	D	543	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	580	ASN
1	B	630	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	801	1	14,14,15	0.16	0	15,19,21	0.55	0
2	NAG	A	802	1	14,14,15	0.33	0	15,19,21	0.57	1 (6%)
2	NAG	A	803	1	14,14,15	0.26	0	15,19,21	0.29	0
2	NAG	B	801	1	14,14,15	0.16	0	15,19,21	0.55	0
2	NAG	B	802	1	14,14,15	0.33	0	15,19,21	0.57	1 (6%)
2	NAG	B	803	1	14,14,15	0.24	0	15,19,21	0.28	0
2	NAG	C	801	1	14,14,15	0.15	0	15,19,21	0.55	0
2	NAG	C	802	1	14,14,15	0.34	0	15,19,21	0.58	1 (6%)
2	NAG	C	803	1	14,14,15	0.25	0	15,19,21	0.29	0
2	NAG	D	801	1	14,14,15	0.15	0	15,19,21	0.56	0
2	NAG	D	802	1	14,14,15	0.34	0	15,19,21	0.57	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	D	803	1	14,14,15	0.24	0	15,19,21	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	801	1	-	0/6/23/26	0/1/1/1
2	NAG	A	802	1	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1	-	0/6/23/26	0/1/1/1
2	NAG	B	801	1	-	0/6/23/26	0/1/1/1
2	NAG	B	802	1	-	0/6/23/26	0/1/1/1
2	NAG	B	803	1	-	0/6/23/26	0/1/1/1
2	NAG	C	801	1	-	0/6/23/26	0/1/1/1
2	NAG	C	802	1	-	0/6/23/26	0/1/1/1
2	NAG	C	803	1	-	0/6/23/26	0/1/1/1
2	NAG	D	801	1	-	0/6/23/26	0/1/1/1
2	NAG	D	802	1	-	0/6/23/26	0/1/1/1
2	NAG	D	803	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	802	NAG	C1-O5-C5	2.04	115.14	112.14
2	B	802	NAG	C1-O5-C5	2.05	115.16	112.14
2	A	802	NAG	C1-O5-C5	2.06	115.17	112.14
2	C	802	NAG	C1-O5-C5	2.09	115.21	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	NAG	1	0
2	B	801	NAG	1	0
2	C	801	NAG	1	0
2	D	801	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.