



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:13 PM GMT

PDB ID : 3T9N
Title : Crystal structure of a membrane protein
Authors : Yang, M.; Zhang, X.; Ge, J.; Wang, J.
Deposited on : 2011-08-03
Resolution : 3.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1000 (3.56-3.36)
Clashscore	102246	1090 (3.56-3.36)
Ramachandran outliers	100387	1057 (3.56-3.36)
Sidechain outliers	100360	1058 (3.56-3.36)
RSRZ outliers	91569	1005 (3.56-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div>7%</div> <div>46% 37% 10% • 6%</div> </div>
1	B	282	<div> <div>9%</div> <div>45% 36% 12% • 6%</div> </div>
1	C	282	<div> <div>8%</div> <div>44% 37% 12% • 6%</div> </div>
1	D	282	<div> <div>9%</div> <div>43% 37% 13% • 6%</div> </div>
1	E	282	<div> <div>10%</div> <div>43% 37% 13% • 6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	282	
1	G	282	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LMT	B	283	-	-	-	X
2	LMT	F	283	-	-	-	X

2 Entry composition ⓘ


There are 2 unique types of molecules in this entry. The entry contains 14789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Small-conductance mechanosensitive channel.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	0	0
			2116	1382	341	383	10			
1	B	264	Total	C	N	O	S	0	0	0
			2099	1373	339	377	10			
1	C	264	Total	C	N	O	S	0	0	0
			2099	1373	339	377	10			
1	D	265	Total	C	N	O	S	0	0	0
			2108	1378	340	380	10			
1	E	264	Total	C	N	O	S	0	0	0
			2099	1373	339	377	10			
1	F	264	Total	C	N	O	S	0	0	0
			2099	1373	339	377	10			
1	G	264	Total	C	N	O	S	0	0	0
			2099	1373	339	377	10			

- Molecule 2 is DODECYL-BETA-D-MALTOSIDE (three-letter code: LMT) (formula: C₂₄H₄₆O₁₁).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	F	1	Total	C	O	0	0
			35	24	11		
2	B	1	Total	C	O	0	0
			35	24	11		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Small-conductance mechanosensitive channel




- Molecule 1: Small-conductance mechanosensitive channel





- Molecule 1: Small-conductance mechanosensitive channel







• Molecule 1: Small-conductance mechanosensitive channel




• Molecule 1: Small-conductance mechanosensitive channel




• Molecule 1: Small-conductance mechanosensitive channel





• Molecule 1: Small-conductance mechanosensitive channel



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	95.03Å 138.63Å 214.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.74 – 3.46 38.74 – 3.46	Depositor EDS
% Data completeness (in resolution range)	95.4 (38.74-3.46) 95.4 (38.74-3.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 3.48Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.253 , 0.288 0.251 , 0.282	Depositor DCC
R_{free} test set	1801 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	110.0	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 94.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 36116 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	14789	wwPDB-VP
Average B, all atoms (Å ²)	136.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: LMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	0/2150	0.75	6/2894 (0.2%)
1	B	0.52	0/2133	0.72	4/2871 (0.1%)
1	C	0.51	0/2133	0.73	4/2871 (0.1%)
1	D	0.55	1/2142 (0.0%)	0.89	3/2883 (0.1%)
1	E	0.54	0/2133	0.69	1/2871 (0.0%)
1	F	0.51	0/2133	0.73	5/2871 (0.2%)
1	G	0.53	0/2133	0.94	6/2871 (0.2%)
All	All	0.53	1/14957 (0.0%)	0.78	29/20132 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	244	GLN	CG-CD	5.09	1.62	1.51

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	250	ARG	NE-CZ-NH2	-23.17	108.72	120.30
1	G	250	ARG	NE-CZ-NH1	20.29	130.44	120.30
1	D	155	ARG	NE-CZ-NH1	-19.53	110.53	120.30
1	D	155	ARG	NE-CZ-NH2	19.11	129.86	120.30
1	G	250	ARG	CD-NE-CZ	10.23	137.92	123.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	211	SER	Peptide
1	A	212	ARG	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2116	0	2222	128	2
1	B	2099	0	2212	127	0
1	C	2099	0	2212	140	0
1	D	2108	0	2218	166	2
1	E	2099	0	2212	142	5
1	F	2099	0	2212	125	3
1	G	2099	0	2212	133	4
2	B	35	0	46	3	1
2	F	35	0	46	10	0
All	All	14789	0	15592	817	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 817 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:273:THR:HG22	1:G:275:ILE:HB	1.24	1.17
1:A:273:THR:HG22	1:D:275:ILE:HB	1.11	1.08
1:C:275:ILE:HB	1:F:273:THR:HG22	1.35	1.07
1:A:269:TYR:OH	1:D:191:ASP:HB2	1.58	1.04
1:D:267:PHE:HB2	1:D:269:TYR:CE1	1.93	1.04

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:202:GLN:OE1	1:E:46:TYR:OH[3_545]	1.91	0.29
1:E:278:GLU:OE1	1:G:212:ARG:NH1[4_545]	1.95	0.25
1:E:278:GLU:OE1	1:G:212:ARG:CZ[4_545]	2.07	0.13
1:F:212:ARG:NH1	2:B:283:LMT:O6B[3_555]	2.11	0.09
1:D:199:GLU:N	1:E:43:TYR:OH[3_545]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/282 (93%)	230 (88%)	19 (7%)	13 (5%)	3	25
1	B	260/282 (92%)	233 (90%)	16 (6%)	11 (4%)	3	31
1	C	260/282 (92%)	232 (89%)	15 (6%)	13 (5%)	3	25
1	D	261/282 (93%)	232 (89%)	16 (6%)	13 (5%)	3	25
1	E	260/282 (92%)	233 (90%)	15 (6%)	12 (5%)	3	28
1	F	260/282 (92%)	232 (89%)	16 (6%)	12 (5%)	3	28
1	G	260/282 (92%)	232 (89%)	16 (6%)	12 (5%)	3	28
All	All	1823/1974 (92%)	1624 (89%)	113 (6%)	86 (5%)	3	28

5 of 86 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	131	VAL
1	A	178	ASP
1	A	218	GLY
1	B	131	VAL
1	B	178	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/250 (94%)	194 (83%)	41 (17%)	2	13
1	B	233/250 (93%)	193 (83%)	40 (17%)	2	14
1	C	233/250 (93%)	192 (82%)	41 (18%)	2	12
1	D	234/250 (94%)	191 (82%)	43 (18%)	2	10
1	E	233/250 (93%)	191 (82%)	42 (18%)	2	11
1	F	233/250 (93%)	192 (82%)	41 (18%)	2	12
1	G	233/250 (93%)	192 (82%)	41 (18%)	2	12
All	All	1634/1750 (93%)	1345 (82%)	289 (18%)	2	12

5 of 289 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	67	LEU
1	D	264	ASN
1	G	158	SER
1	D	95	LEU
1	D	179	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	89	ASN
1	G	162	HIS
1	F	89	ASN
1	D	89	ASN
1	G	89	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LMT	B	283	-	36,36,36	0.57	1 (2%)	47,47,47	1.03	5 (10%)
2	LMT	F	283	-	36,36,36	0.50	1 (2%)	47,47,47	1.29	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LMT	B	283	-	-	0/21/61/61	0/2/2/2
2	LMT	F	283	-	-	0/21/61/61	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	283	LMT	O1'-C1'	2.01	1.43	1.40
2	F	283	LMT	O1'-C1'	2.03	1.43	1.40

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	283	LMT	C3'-C4'-C5'	-2.39	105.44	110.84
2	F	283	LMT	C3B-C4B-C5B	-2.28	106.22	110.20
2	F	283	LMT	O2B-C2B-C1B	-2.15	105.30	110.02
2	B	283	LMT	C1B-O5B-C5B	-2.06	109.75	113.75
2	B	283	LMT	C3B-C4B-C5B	2.17	113.99	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	283	LMT	3	1
2	F	283	LMT	10	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	266/282 (94%)	0.34	21 (7%)	15 15	65, 120, 204, 296	0
1	B	264/282 (93%)	0.29	24 (9%)	11 11	62, 121, 206, 315	0
1	C	264/282 (93%)	0.38	22 (8%)	14 14	69, 124, 206, 324	0
1	D	265/282 (93%)	0.43	25 (9%)	11 11	62, 120, 231, 381	0
1	E	264/282 (93%)	0.40	27 (10%)	9 9	70, 126, 244, 426	0
1	F	264/282 (93%)	0.42	23 (8%)	13 13	62, 126, 231, 318	0
1	G	264/282 (93%)	0.19	20 (7%)	17 16	63, 129, 226, 322	0
All	All	1851/1974 (93%)	0.35	162 (8%)	12 13	62, 124, 225, 426	0

The worst 5 of 162 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	281	THR	27.3
1	F	280	LYS	13.5
1	C	178	ASP	13.2
1	G	277	SER	10.0
1	E	279	LYS	9.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	LMT	F	283	35/35	0.76	0.54	7.61	68,158,220,252	0
2	LMT	B	283	35/35	0.68	0.48	1.88	67,180,255,265	0

6.5 Other polymers [i](#)

There are no such residues in this entry.