



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 29, 2016 – 07:28 PM EDT

PDB ID : 5TBM
Title : Crystal structure of PT2385 bound to HIF2a-B*:ARNT-B* complex
Authors : Du, X.
Deposited on : 2016-09-12
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	115	<div> <div>8%</div> <div> <div></div> <div>71%</div> <div>17%</div> <div>• • 8%</div> </div> </div>
2	B	117	<div> <div>3%</div> <div> <div></div> <div>62%</div> <div>29%</div> <div>• 6%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endothelial PAS domain-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	106	Total	C	N	O	S	0	1	0
			865	549	140	167	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	234	GLY	-	expression tag	UNP Q99814
A	235	GLU	-	expression tag	UNP Q99814
A	236	PHE	-	expression tag	UNP Q99814
A	237	LEU	ILE	conflict	UNP Q99814
A	238	GLY	PRO	conflict	UNP Q99814
A	247	GLU	ARG	engineered mutation	UNP Q99814


- Molecule 2 is a protein called Aryl hydrocarbon receptor nuclear translocator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	110	Total	C	N	O	S	0	0	0
			917	583	161	167	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	351	GLY	-	expression tag	UNP P27540
B	352	GLU	-	expression tag	UNP P27540
B	353	PHE	-	expression tag	UNP P27540
B	354	LEU	MET	conflict	UNP P27540
B	355	GLY	SER	conflict	UNP P27540
B	362	ARG	GLU	engineered mutation	UNP P27540

- Molecule 3 is 3-[[[(1S)-2,2-difluoro-1-hydroxy-7-(methylsulfonyl)-2,3-dihydro-1H-inden-4-yl]oxy]-5-fluorobenzonitrile (three-letter code: 79A) (formula: C₁₇H₁₂F₃NO₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
3	A	1	26	17	3	1	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	12	Total	O	0	0
			12	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endothelial PAS domain-containing protein 1



- Molecule 2: Aryl hydrocarbon receptor nuclear translocator



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	73.40Å 84.10Å 41.40Å 90.00° 106.24° 90.00°	Depositor
Resolution (Å)	28.89 – 1.85 28.89 – 1.51	Depositor EDS
% Data completeness (in resolution range)	98.6 (28.89-1.85) 66.7 (28.89-1.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 1.51Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.237 , 0.300 0.239 , 0.241	Depositor DCC
R_{free} test set	1016 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.521	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1824	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 79A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	1/888 (0.1%)	0.98	3/1198 (0.3%)
2	B	1.03	1/939 (0.1%)	1.02	2/1269 (0.2%)
All	All	0.98	2/1827 (0.1%)	1.00	5/2467 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	278	TYR	CB-CG	-5.21	1.43	1.51
2	B	458	ILE	C-O	5.13	1.33	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	409	ARG	NE-CZ-NH1	7.27	123.94	120.30
1	A	275	ARG	NE-CZ-NH1	-6.16	117.22	120.30
2	B	409	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	319	LEU	CA-CB-CG	5.92	128.93	115.30
1	A	272	LEU	CA-CB-CG	5.03	126.87	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	865	0	824	20	0
2	B	917	0	891	20	1
3	A	26	0	0	2	0
4	A	4	0	0	0	0
4	B	12	0	0	2	0
All	All	1824	0	1715	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:ASN:HB3	1:A:329:PRO:HD3	1.40	1.03
2:B:409:ARG:HD3	4:B:512:HOH:O	1.67	0.94
1:A:347:ILE:HG22	1:A:348:GLU:H	1.39	0.87
1:A:328:ASN:HB3	1:A:329:PRO:CD	2.15	0.76
1:A:326:ILE:CD1	1:A:338:MET:HG2	2.22	0.68
1:A:319:LEU:CD2	1:A:343:VAL:HG22	2.24	0.67
2:B:376:VAL:HG11	2:B:391:LEU:HD12	1.83	0.60
2:B:381:VAL:O	2:B:385:GLY:HA2	2.02	0.59
1:A:347:ILE:HG22	1:A:348:GLU:N	2.14	0.59
1:A:309:MET:HG3	3:A:401:79A:N1	2.17	0.59
2:B:418:LEU:O	2:B:421:GLN:NE2	2.28	0.58
1:A:326:ILE:HD12	1:A:338:MET:HG2	1.85	0.58
1:A:239:LEU:HA	1:A:342[A]:TYR:HE1	1.69	0.57
2:B:389:GLN:N	2:B:389:GLN:OE1	2.35	0.57
2:B:417:LYS:HE3	4:B:507:HOH:O	2.04	0.56
1:A:326:ILE:HD11	1:A:338:MET:HG2	1.87	0.56
1:A:335:GLN:O	1:A:336:CYS:HB3	2.06	0.54
1:A:319:LEU:HD22	1:A:343:VAL:HG22	1.90	0.53
2:B:366:ARG:HG2	2:B:374:THR:OG1	2.09	0.53
2:B:366:ARG:HD2	2:B:456:TYR:CD1	2.48	0.48
2:B:401:HIS:HB2	2:B:428:ARG:HB2	1.96	0.48
2:B:418:LEU:O	2:B:421:GLN:HG3	2.13	0.47
2:B:364:ILE:O	2:B:377:ASP:HB3	2.15	0.46
2:B:362:ARG:HA	2:B:461:ASN:O	2.15	0.46
2:B:431:SER:HB3	2:B:437:LEU:HD11	1.98	0.46
1:A:244:PHE:HB2	1:A:258:ASP:HB2	1.98	0.45
1:A:239:LEU:HD23	1:A:342[A]:TYR:OH	2.16	0.45
2:B:389:GLN:CD	2:B:389:GLN:H	2.19	0.45
1:A:296:LEU:HD22	3:A:401:79A:O2	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:HIS:O	1:A:271:GLU:HG2	2.16	0.44
1:A:269:PRO:O	1:A:272:LEU:HB2	2.19	0.43
1:A:347:ILE:CG2	1:A:348:GLU:H	2.21	0.43
2:B:395:ASN:OD1	2:B:396:ILE:N	2.52	0.43
2:B:414:GLN:O	2:B:415:VAL:C	2.57	0.42
2:B:420:GLY:HA2	2:B:445:THR:OG1	2.20	0.41
2:B:430:ARG:HB2	2:B:436:TRP:CZ3	2.55	0.41
2:B:448:ASN:HA	2:B:449:PRO:HD3	1.89	0.41
1:A:319:LEU:HD21	1:A:343:VAL:HG22	2.00	0.40
2:B:362:ARG:HD2	2:B:460:THR:CG2	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:410:ASP:OD2	2:B:410:ASP:OD2[2_554]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/115 (90%)	96 (93%)	6 (6%)	1 (1%)	19	6
2	B	108/117 (92%)	104 (96%)	4 (4%)	0	100	100
All	All	211/232 (91%)	200 (95%)	10 (5%)	1 (0%)	34	17

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	328	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/103 (94%)	92 (95%)	5 (5%)	29	11
2	B	105/110 (96%)	99 (94%)	6 (6%)	25	9
All	All	202/213 (95%)	191 (95%)	11 (5%)	27	10

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	GLU
1	A	278	TYR
1	A	328	ASN
1	A	346	GLU
1	A	347	ILE
2	B	361	THR
2	B	377	ASP
2	B	410	ASP
2	B	447	GLN
2	B	455	GLU
2	B	463	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	378	HIS
2	B	433	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	79A	A	401	-	25,28,28	2.37	5 (20%)	30,44,44	2.33	12 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	79A	A	401	-	-	0/12/27/27	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	79A	C4-S1	-8.42	1.66	1.77
3	A	401	79A	C6-C17	-5.45	1.30	1.44
3	A	401	79A	C1-C14	-2.99	1.36	1.40
3	A	401	79A	C16-S1	-2.35	1.63	1.74
3	A	401	79A	C3-C2	3.08	1.44	1.38

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	79A	O3-S1-C4	-3.86	103.39	108.12
3	A	401	79A	C5-C10-C9	-3.70	118.75	123.54
3	A	401	79A	O2-S1-C4	-3.61	103.69	108.12
3	A	401	79A	O1-C1-C14	-3.15	111.33	117.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	79A	C1-C14-C13	-2.77	118.34	120.97
3	A	401	79A	O3-S1-O2	-2.06	113.23	117.73
3	A	401	79A	C2-C1-C14	2.21	123.79	120.62
3	A	401	79A	C15-C14-C13	2.62	113.61	111.00
3	A	401	79A	F1-C10-C9	2.81	122.01	118.22
3	A	401	79A	F3-C11-C12	3.01	116.94	112.07
3	A	401	79A	O2-S1-C16	4.53	116.18	108.54
3	A	401	79A	O3-S1-C16	5.20	117.31	108.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	79A	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	106/115 (92%)	0.42	9 (8%) 13 13	28, 42, 64, 103	0
2	B	110/117 (94%)	-0.22	4 (3%) 46 44	28, 38, 61, 77	0
All	All	216/232 (93%)	0.09	13 (6%) 25 24	28, 40, 64, 103	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	347	ILE	6.1
1	A	348	GLU	5.0
2	B	359	GLN	3.4
1	A	342[A]	TYR	3.1
1	A	339	CYS	3.0
1	A	346	GLU	2.9
1	A	278	TYR	2.8
1	A	298	THR	2.7
1	A	314	GLY	2.7
1	A	315	GLY	2.5
2	B	433	ASN	2.4
2	B	454	ILE	2.1
2	B	434	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	79A	A	401	26/26	0.91	0.16	0.91	24,32,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.