



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2016 – 05:43 AM EST

PDB ID : 5TCS
Title : Crystal structure of a Dwarf Ndc80 Tetramer
Authors : Valverde, R.; Harrison, S.C.
Deposited on : 2016-09-15
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028320
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028320


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3170 (2.88-2.80)
Clashscore	102246	3658 (2.88-2.80)
Ramachandran outliers	100387	3591 (2.88-2.80)
Sidechain outliers	100360	3594 (2.88-2.80)
RSRZ outliers	91569	3184 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	<div> <div>83%</div> <div>12%</div> <div>• •</div> </div>
2	B	215	<div> <div>7%</div> <div>90%</div> <div>7%</div> <div>•</div> </div>
3	C	100	<div> <div>37%</div> <div>88%</div> <div>9%</div> <div>•</div> </div>
4	D	115	<div> <div>14%</div> <div>85%</div> <div>14%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MG	B	501	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11358 atoms, of which 5650 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinetochore protein NDC80.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	266	Total	C	H	N	O	S	Se	0	0	0
			4501	1437	2264	375	419	1	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	111	SER	-	expression tag	UNP P40460
A	112	ASN	-	expression tag	UNP P40460
A	113	ALA	-	expression tag	UNP P40460

- Molecule 2 is a protein called Kinetochore protein NUF2.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
2	B	211	Total	C	H	N	O	S	Se	0	0	0
			3420	1093	1691	280	344	4	8			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	SER	-	expression tag	UNP P33895
B	-16	ASN	-	expression tag	UNP P33895
B	-15	ALA	-	expression tag	UNP P33895
B	-14	SER	-	expression tag	UNP P33895
B	-13	ILE	-	expression tag	UNP P33895
B	-12	PHE	-	expression tag	UNP P33895
B	-11	LYS	-	expression tag	UNP P33895
B	-10	ASP	-	expression tag	UNP P33895
B	-9	LEU	-	expression tag	UNP P33895
B	-8	GLU	-	expression tag	UNP P33895
B	-7	ALA	-	expression tag	UNP P33895
B	-6	LEU	-	expression tag	UNP P33895
B	-5	SER	-	expression tag	UNP P33895

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	PHE	-	expression tag	UNP P33895
B	-3	GLN	-	expression tag	UNP P33895
B	-2	SER	-	expression tag	UNP P33895
B	-1	ASN	-	expression tag	UNP P33895
B	0	ALA	-	expression tag	UNP P33895

- Molecule 3 is a protein called Kinetochore protein SPC24.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	97	Total	C	H	N	O	Se	0	0	0
			1626	511	815	141	158	1			

- Molecule 4 is a protein called Kinetochore protein SPC25.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
4	D	114	Total	C	H	N	O	S	Se	0	0	0
			1771	559	880	164	164	1	3			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	19	Total	O	0	0
			19	19		
6	B	15	Total	O	0	0
			15	15		
6	D	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Kinetochores protein NDC80




- Molecule 2: Kinetochores protein NUF2



- Molecule 3: Kinetochores protein SPC24



- Molecule 4: Kinetochores protein SPC25



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	169.38Å 186.59Å 122.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.75 – 2.83 49.51 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.3 (41.75-2.83) 99.3 (49.51-2.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, R_{free}	0.239 , 0.266 0.234 , 0.265	Depositor DCC
R_{free} test set	2332 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	80.2	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 58.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11358	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2275	0.41	0/3055
2	B	0.27	0/1746	0.41	0/2338
3	C	0.24	0/820	0.46	0/1101
4	D	0.25	0/904	0.44	0/1220
All	All	0.26	0/5745	0.42	0/7714

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	147	HIS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2237	2264	2264	28	1
2	B	1729	1691	1693	15	1
3	C	811	815	815	6	1
4	D	891	880	880	9	0
5	B	1	0	0	0	0
6	A	19	0	0	0	0
6	B	15	0	0	0	0
6	D	5	0	0	1	0
All	All	5708	5650	5652	48	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:157:HIS:HA	4:D:174:ALA:HB2	1.79	0.63
3:C:194:LEU:O	3:C:194:LEU:HG	2.00	0.61
1:A:668:GLU:OE1	3:C:33:ARG:NH1	2.38	0.56
3:C:190:ASP:OD2	3:C:210:ARG:NH2	2.40	0.54
1:A:657:LYS:NZ	3:C:27:ASP:OD2	2.35	0.54
4:D:26:ALA:O	6:D:301:HOH:O	2.18	0.54
2:B:33:ILE:O	2:B:33:ILE:HG22	2.09	0.52
1:A:253:ILE:HD13	1:A:270:ARG:HH12	1.75	0.52
1:A:267:ARG:NH2	1:A:318:ASP:OD2	2.41	0.52
1:A:115:ASP:HB3	1:A:116:PRO:HD3	1.92	0.50
1:A:159:GLN:NE2	1:A:208:SER:O	2.44	0.50
1:A:147:HIS:HB2	1:A:148:PRO:CD	2.41	0.50
1:A:272:GLU:HA	2:B:147:MSE:HE1	1.95	0.49
4:D:182:GLY:O	4:D:193:ARG:NH2	2.42	0.48
2:B:32:ASN:HB3	2:B:40:TYR:CZ	2.48	0.48
1:A:670:GLU:HG2	4:D:27:ARG:CZ	2.44	0.48
1:A:275:VAL:HB	2:B:147:MSE:HE1	1.96	0.47
1:A:160:LYS:O	1:A:164:ILE:HD13	2.15	0.46
1:A:238:LYS:CE	2:B:67:GLU:HG2	2.45	0.46
2:B:35:ARG:O	2:B:37:THR:N	2.44	0.45
2:B:33:ILE:HG12	2:B:40:TYR:OH	2.17	0.45
2:B:34:SER:O	2:B:37:THR:OG1	2.26	0.45
1:A:148:PRO:C	1:A:151:ILE:HG22	2.37	0.44
3:C:192:LEU:HD12	3:C:193:PRO:HD2	2.00	0.44
2:B:79:ASN:O	2:B:79:ASN:OD1	2.36	0.44
1:A:157:PRO:HG3	1:A:215:TRP:CE2	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:649:VAL:CG2	1:A:650:ILE:N	2.80	0.43
2:B:431:GLU:HG3	3:C:12:VAL:HG13	2.00	0.43
1:A:248:THR:O	1:A:251:ILE:HG22	2.19	0.43
4:D:185:HIS:HB3	4:D:186:PRO:HD3	2.01	0.43
4:D:188:LEU:HD21	4:D:217:LEU:HD23	2.01	0.42
1:A:121:ASP:OD1	1:A:123:ASN:N	2.53	0.42
1:A:246:GLN:O	1:A:249:GLN:HG2	2.18	0.42
2:B:450:MSE:O	2:B:451:GLN:HB2	2.19	0.42
1:A:115:ASP:CB	1:A:116:PRO:HD3	2.50	0.42
4:D:9:ASP:O	4:D:12:ARG:HB2	2.20	0.42
2:B:87:VAL:HG11	2:B:108:MSE:HE1	2.02	0.42
1:A:653:THR:HA	4:D:14:MSE:HE1	2.02	0.42
1:A:152:LYS:HG2	1:A:153:PHE:HD2	1.84	0.41
1:A:275:VAL:HB	2:B:147:MSE:CE	2.50	0.41
1:A:266:GLN:O	1:A:270:ARG:HG2	2.20	0.41
2:B:74:GLN:HG2	2:B:74:GLN:O	2.20	0.41
1:A:148:PRO:C	1:A:151:ILE:CG2	2.89	0.41
1:A:238:LYS:HE2	2:B:67:GLU:HG2	2.02	0.41
1:A:148:PRO:HB2	1:A:151:ILE:CG2	2.50	0.41
1:A:147:HIS:HB3	1:A:164:ILE:HG12	2.02	0.41
1:A:234:MSE:HG3	1:A:238:LYS:HZ1	1.86	0.40
4:D:150:LEU:HD12	4:D:162:VAL:HG21	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:-1:ASN:H	2:B:135:GLU:OE2[4_545]	1.52	0.08
1:A:130:GLU:OE2	3:C:4:LYS:HZ3[3_655]	1.56	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/277 (95%)	234 (89%)	30 (11%)	0	100	100
2	B	209/215 (97%)	198 (95%)	11 (5%)	0	100	100
3	C	95/100 (95%)	92 (97%)	3 (3%)	0	100	100
4	D	112/115 (97%)	104 (93%)	8 (7%)	0	100	100
All	All	680/707 (96%)	628 (92%)	52 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/261 (98%)	255 (100%)	1 (0%)	93	98
2	B	198/193 (103%)	197 (100%)	1 (0%)	92	98
3	C	92/93 (99%)	92 (100%)	0	100	100
4	D	95/92 (103%)	94 (99%)	1 (1%)	80	95
All	All	641/639 (100%)	638 (100%)	3 (0%)	92	98

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	248	THR
2	B	40	TYR
4	D	141	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/277 (94%)	0.52	4 (1%) 76 69	36, 69, 118, 248	0
2	B	203/215 (94%)	0.89	14 (6%) 20 12	27, 62, 158, 203	0
3	C	96/100 (96%)	1.83	37 (38%) 0 0	52, 124, 190, 214	0
4	D	111/115 (96%)	0.95	16 (14%) 3 2	64, 100, 163, 185	0
All	All	671/707 (94%)	0.89	71 (10%) 8 4	27, 75, 172, 248	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	150	SER	20.0
1	A	147	HIS	13.0
2	B	72	HIS	11.7
2	B	74	GLN	9.8
2	B	71	GLY	8.9
3	C	196	ASN	8.7
2	B	75	GLU	8.1
2	B	70	ASP	7.8
3	C	194	LEU	7.1
2	B	69	GLY	6.8
3	C	179	VAL	6.3
3	C	191	ILE	6.2
3	C	180	LEU	6.1
4	D	185	HIS	5.8
4	D	157	HIS	5.7
4	D	156	VAL	5.7
3	C	186	ASP	5.6
2	B	35	ARG	5.3
3	C	181	ILE	5.2
3	C	198	LEU	5.1
3	C	202	TYR	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	221	LEU	4.7
3	C	204	THR	4.7
1	A	149	ILE	4.6
2	B	73	LEU	4.4
3	C	195	ASP	4.4
3	C	192	LEU	4.3
3	C	213	LYS	4.3
3	C	174	LEU	4.2
2	B	76	GLU	4.2
3	C	187	GLY	4.1
3	C	207	ILE	3.9
3	C	183	ARG	3.8
4	D	198	HIS	3.8
3	C	212	GLY	3.7
3	C	188	ASN	3.6
4	D	27	ARG	3.4
3	C	206	TYR	3.4
3	C	190	ASP	3.4
3	C	170	VAL	3.3
4	D	155	ASP	3.3
4	D	28	GLN	3.2
3	C	184	LYS	3.0
3	C	211	LEU	3.0
3	C	172	LEU	3.0
4	D	167	SER	3.0
2	B	67	GLU	2.9
3	C	171	ILE	2.8
4	D	187	ALA	2.8
3	C	205	LYS	2.8
4	D	219	ALA	2.6
2	B	5	GLN	2.6
3	C	182	ASN	2.6
3	C	209	GLU	2.6
3	C	178	GLN	2.5
3	C	208	TRP	2.4
4	D	200	LEU	2.4
3	C	193	PRO	2.4
3	C	199	SER	2.4
4	D	140	LEU	2.3
4	D	12	ARG	2.3
3	C	197	ASN	2.3
2	B	447	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	210	LEU	2.3
3	C	189	ILE	2.2
3	C	201	PHE	2.2
4	D	209	PHE	2.2
2	B	68	THR	2.1
3	C	164	LEU	2.0
1	A	163	ILE	2.0
2	B	-11	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MG	B	501	1/1	0.97	0.66	20.20	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.