



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:30 PM GMT

PDB ID : 4TMW  
Title : Translation initiation factor eIF5B (517-858) from *C. thermophilum*, bound to GTP and Sodium  
Authors : Kuhle, B.; Ficner, F.  
Deposited on : 2014-06-02  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1665 (1.58-1.54)
Clashscore	102246	1014 (1.56-1.56)
Ramachandran outliers	100387	1704 (1.58-1.54)
Sidechain outliers	100360	1702 (1.58-1.54)
RSRZ outliers	91569	1668 (1.58-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	<div> <div>5%</div> <div>86%</div> <div>12%</div> <div>..</div> </div>
1	B	345	<div> <div>4%</div> <div>88%</div> <div>12%</div> <div>.</div> </div>

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called eIF5B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	0	29	0
			2911	1827	520	552	12			
1	B	345	Total	C	N	O	S	0	20	0
			2849	1796	507	533	13			

- Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0
3	A	1	Total Na 1 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	353	Total O 366 366	0	13
5	B	388	Total O 392 392	0	4


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: eIF5B



#### • Molecule 1: eIF5B



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.61Å 116.45Å 66.24Å 90.00° 101.15° 90.00°	Depositor
Resolution (Å)	43.37 – 1.55 46.44 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.8 (43.37-1.55) 99.1 (46.44-1.55)	Depositor EDS
$R_{merge}$	1.00	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.18 (at 1.55Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1690)	Depositor
R, $R_{free}$	0.157 , 0.184 0.159 , 0.184	Depositor DCC
$R_{free}$ test set	5919 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.3	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 118389 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6586	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, NA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.88	3/2953 (0.1%)	0.95	15/3981 (0.4%)
1	B	0.83	2/2890 (0.1%)	0.86	7/3895 (0.2%)
All	All	0.85	5/5843 (0.1%)	0.91	22/7876 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	527[A]	CYS	CB-SG	-8.22	1.68	1.82
1	A	527[B]	CYS	CB-SG	-8.22	1.68	1.82
1	A	788	PRO	N-CD	5.32	1.55	1.47
1	B	844	PRO	N-CD	5.20	1.55	1.47
1	B	524	PRO	N-CD	5.14	1.55	1.47

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	779[A]	ARG	NE-CZ-NH2	-8.24	116.18	120.30
1	A	779[B]	ARG	NE-CZ-NH2	-8.24	116.18	120.30
1	A	779[A]	ARG	NE-CZ-NH1	7.63	124.11	120.30
1	A	779[B]	ARG	NE-CZ-NH1	7.63	124.11	120.30
1	A	806	ARG	NE-CZ-NH1	7.61	124.11	120.30
1	A	847	ASP	O-C-N	-7.38	110.88	122.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	789	ILE	O-C-N	-7.18	111.21	122.70
1	A	803[A]	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	A	803[B]	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	B	682	ARG	NE-CZ-NH2	-7.15	116.72	120.30
1	A	847	ASP	CA-C-N	6.92	132.43	117.20
1	B	806	ARG	NE-CZ-NH1	6.87	123.74	120.30
1	A	789	ILE	CA-C-N	6.83	132.24	117.20
1	A	803[A]	ARG	NE-CZ-NH2	-6.52	117.04	120.30
1	A	803[B]	ARG	NE-CZ-NH2	-6.52	117.04	120.30
1	A	737	ARG	NE-CZ-NH1	-5.82	117.39	120.30
1	B	843	GLY	C-N-CD	5.58	140.13	128.40
1	A	787	GLY	C-N-CD	5.56	140.08	128.40
1	B	523	SER	C-N-CD	5.39	139.71	128.40
1	B	775	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	B	605	ARG	NE-CZ-NH2	-5.26	117.67	120.30
1	B	526	CYS	CA-C-N	5.18	128.60	117.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	558	GLN	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2911	0	2928	40	0
1	B	2849	0	2904	25	0
2	A	32	0	12	0	0
2	B	32	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	366	0	0	3	0
5	B	392	0	0	10	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6586	0	5856	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:790:LYS:HE2	1:B:852:GLU:OE1	1.55	1.05
1:A:580:ASP:OD2	1:A:582:LYS:NZ	2.09	0.86
1:A:790[B]:LYS:CG	1:A:848[B]:GLU:OE2	2.36	0.73
1:A:779[B]:ARG:NE	1:A:790[B]:LYS:HG2	2.03	0.72
1:A:849:GLU:O	1:A:853:GLU:OE1	2.07	0.71
1:B:737[A]:ARG:NH2	5:B:1232:HOH:O	2.25	0.70
1:A:559[A]:GLN:NE2	1:A:754[A]:GLU:OE2	2.26	0.69
1:B:803[B]:ARG:NH1	5:B:1222:HOH:O	2.25	0.67
1:A:517:ASN:HD21	1:A:519:ASP:HB2	1.59	0.67
1:B:639[A]:ARG:NH2	5:B:1162:HOH:O	2.28	0.66
1:A:779[B]:ARG:CZ	1:A:790[B]:LYS:HG2	2.26	0.65
1:B:514:GLY:O	1:B:517:ASN:ND2	2.30	0.65
1:A:779[B]:ARG:HE	1:A:790[B]:LYS:HG2	1.64	0.63
1:A:517:ASN:ND2	1:A:519:ASP:HB2	2.13	0.63
1:A:558:GLN:NE2	5:A:1226[A]:HOH:O	2.31	0.62
1:B:680:ARG:NH1	5:B:1001:HOH:O	2.18	0.61
1:A:781[B]:VAL:HG22	1:A:855:VAL:HG11	1.82	0.60
1:A:779[B]:ARG:NH1	1:A:790[B]:LYS:HG2	2.15	0.60
1:A:842:VAL:HG22	1:A:851:LEU:HD12	1.83	0.59
1:B:749[B]:GLN:HG2	5:B:1374:HOH:O	2.04	0.57
1:B:657:LYS:NZ	5:B:1308:HOH:O	2.32	0.57
1:B:790:LYS:HE2	1:B:852:GLU:CD	2.23	0.56
1:A:838[B]:ARG:NH1	1:A:854:GLU:O	2.40	0.55
1:A:781[A]:VAL:HG22	1:A:840:LEU:HB2	1.88	0.54
1:B:781[B]:VAL:HG13	1:B:840:LEU:HB2	1.89	0.54
1:B:681[B]:ASN:OD1	1:B:685:GLN:NE2	2.37	0.53
5:A:1327:HOH:O	1:B:568:GLU:HG3	2.09	0.53
1:B:781[A]:VAL:HG22	1:B:840:LEU:HB2	1.93	0.51
1:A:637:ARG:HH11	1:A:637:ARG:HG3	1.76	0.50
1:B:560:ILE:HG23	1:B:593[B]:ILE:HG23	1.92	0.50
1:A:790[B]:LYS:CG	1:A:848[B]:GLU:CD	2.80	0.49
1:A:781[B]:VAL:HG13	1:A:840:LEU:HB2	1.95	0.49
1:A:527[A]:CYS:SG	1:A:612:CYS:HB3	2.53	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:527[A]:CYS:SG	1:A:608:GLY:O	2.72	0.48
1:A:802:MET:HE2	1:A:802:MET:HB3	1.70	0.47
1:B:645:VAL:HB	1:B:712:LEU:HD23	1.97	0.47
1:A:644[A]:VAL:HG13	1:A:733:LEU:HD11	1.96	0.47
1:B:781[B]:VAL:CG1	1:B:840:LEU:HB2	2.45	0.47
1:B:848:GLU:O	1:B:852:GLU:HG3	2.15	0.47
1:B:527[A]:CYS:HB2	1:B:612:CYS:HB3	1.97	0.46
1:A:560:ILE:HG23	1:A:593:ILE:HG23	1.98	0.46
1:A:779[B]:ARG:HH11	1:A:790[B]:LYS:HG2	1.79	0.46
1:A:637:ARG:HG3	1:A:637:ARG:NH1	2.31	0.45
1:B:625:LEU:HD22	1:B:689:GLN:HB3	1.99	0.45
1:B:781[B]:VAL:HG11	1:B:851:LEU:HB3	1.99	0.45
1:B:756:LYS:HD2	5:B:1212:HOH:O	2.17	0.44
1:B:600[B]:SER:OG	5:B:1002:HOH:O	2.19	0.44
1:A:657[A]:LYS:HA	1:A:657[A]:LYS:HD3	1.81	0.44
1:A:625:LEU:HD22	1:A:689:GLN:HB3	2.00	0.44
1:A:685:GLN:HG2	5:A:1243:HOH:O	2.16	0.44
1:A:781[B]:VAL:HG21	1:A:852:GLU:HA	2.01	0.43
1:B:680:ARG:HD2	5:B:1001:HOH:O	2.19	0.42
1:A:790[B]:LYS:HG2	1:A:848[B]:GLU:OE2	2.17	0.42
1:A:848[B]:GLU:O	1:A:852:GLU:HG3	2.20	0.42
1:A:544:ARG:HA	1:A:802:MET:CE	2.50	0.42
1:B:559:GLN:NE2	1:B:754:GLU:OE2	2.53	0.41
1:A:779[B]:ARG:NE	1:A:790[B]:LYS:CG	2.78	0.41
1:B:688[B]:LEU:HD23	5:B:1168:HOH:O	2.21	0.40
1:A:515:HIS:O	1:A:515:HIS:ND1	2.55	0.40
1:A:783:CYS:SG	1:A:855:VAL:HG22	2.61	0.40
1:A:851:LEU:HD23	1:A:851:LEU:HA	1.82	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/345 (107%)	365 (99%)	5 (1%)	0	100	100
1	B	363/345 (105%)	360 (99%)	3 (1%)	0	100	100
All	All	733/690 (106%)	725 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/295 (108%)	315 (99%)	4 (1%)	76	51
1	B	314/295 (106%)	307 (98%)	7 (2%)	60	27
All	All	633/590 (107%)	622 (98%)	11 (2%)	68	38

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	748	LEU
1	A	782	LEU
1	A	821	GLN
1	A	851	LEU
1	B	704	LYS
1	B	737[A]	ARG
1	B	737[B]	ARG
1	B	782	LEU
1	B	790	LYS
1	B	821	GLN
1	B	845	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	517	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GTP	A	901	3,4	25,34,34	1.08	2 (8%)	34,54,54	1.87	8 (23%)
2	GTP	B	901	3,4	25,34,34	1.07	2 (8%)	34,54,54	1.87	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	A	901	3,4	-	0/18/38/38	0/3/3/3
2	GTP	B	901	3,4	-	0/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	GTP	C5-C4	2.66	1.46	1.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	GTP	C5-C4	2.79	1.46	1.40
2	B	901	GTP	C6-C5	3.02	1.47	1.41
2	A	901	GTP	C6-C5	3.28	1.47	1.41

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	GTP	C5-C6-N1	-4.80	117.02	123.59
2	A	901	GTP	C5-C6-N1	-4.41	117.56	123.59
2	A	901	GTP	C6-C5-C4	-3.48	116.74	120.90
2	A	901	GTP	C2'-C1'-N9	-3.35	109.18	114.29
2	A	901	GTP	N3-C2-N1	-3.21	122.56	127.44
2	B	901	GTP	C6-C5-C4	-3.07	117.23	120.90
2	B	901	GTP	N3-C2-N1	-3.06	122.79	127.44
2	B	901	GTP	C4-C5-N7	-3.03	106.69	109.48
2	B	901	GTP	C2'-C1'-N9	-2.83	109.97	114.29
2	B	901	GTP	PA-O3A-PB	-2.74	125.03	132.73
2	A	901	GTP	C4-C5-N7	-2.73	106.97	109.48
2	A	901	GTP	PA-O3A-PB	-2.68	125.22	132.73
2	B	901	GTP	PB-O3B-PG	-2.11	125.58	132.67
2	A	901	GTP	PB-O3B-PG	-2.03	125.87	132.67
2	B	901	GTP	O3G-PG-O2G	2.01	115.04	107.38
2	A	901	GTP	C6-N1-C2	5.07	122.97	115.94
2	B	901	GTP	C6-N1-C2	5.29	123.29	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	343/345 (99%)	0.00	16 (4%) 35 36	12, 20, 50, 95	0
1	B	345/345 (100%)	-0.04	14 (4%) 41 43	12, 20, 45, 73	0
All	All	688/690 (99%)	-0.02	30 (4%) 38 39	12, 20, 46, 95	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	857	SER	6.8
1	A	855	VAL	6.4
1	A	856	GLU	6.2
1	A	515	HIS	5.5
1	A	853	GLU	5.5
1	A	517	ASN	5.2
1	B	857	SER	4.4
1	A	516	MET	4.1
1	B	514	GLY	3.7
1	A	854	GLU	3.7
1	A	519	ASP	3.6
1	B	858	ASP	3.2
1	A	851	LEU	3.2
1	B	707	ALA	3.1
1	B	515	HIS	2.9
1	B	853	GLU	2.8
1	B	761	PHE	2.8
1	B	709	TYR	2.7
1	A	688[A]	LEU	2.7
1	A	805	LEU	2.7
1	B	654	TYR	2.6
1	B	760	GLY	2.5
1	B	708	ARG	2.5
1	B	759	GLU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	582	LYS	2.3
1	A	654	TYR	2.3
1	A	518	LYS	2.2
1	A	808	LYS	2.2
1	B	856	GLU	2.1
1	B	673	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	B	902	1/1	0.99	0.09	0.12	15,15,15,15	0
3	NA	A	902	1/1	0.98	0.07	-0.22	15,15,15,15	0
2	GTP	A	901	32/32	0.98	0.07	-0.55	11,14,17,19	0
4	MG	A	903	1/1	0.99	0.08	-0.65	13,13,13,13	0
2	GTP	B	901	32/32	0.98	0.07	-0.74	12,14,17,18	0
4	MG	B	903	1/1	0.99	0.06	-1.84	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.