



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 14, 2016 – 10:04 PM EDT

PDB ID : 4TN0  
Title : Crystal Structure of the C-terminal Periplasmic Domain of Phospho-ethanolamine Transferase EptC from Campylobacter jejuni  
Authors : Fage, C.D.; Brown, D.; Boll, J.M.; Keatinge-Clay, A.T.; Trent, M.S.  
Deposited on : 2014-06-02  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	<div> <div></div> <div>83% 11% 5%</div> </div>
1	B	325	<div> <div></div> <div>84% 10% 6%</div> </div>
1	C	325	<div> <div></div> <div>82% 12% 5%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UPF0141 protein yjdB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	P	S	0	0	0
			2467	1574	398	483	1	11			
1	B	307	Total	C	N	O	P	S	0	0	0
			2456	1568	394	482	1	11			
1	C	308	Total	C	N	O	P	S	0	0	0
			2467	1574	398	483	1	11			

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	MET	-	initiating methionine	UNP A3ZEY5
A	513	GLY	-	expression tag	UNP A3ZEY5
A	514	GLY	-	expression tag	UNP A3ZEY5
A	515	GLY	-	expression tag	UNP A3ZEY5
A	516	GLY	-	expression tag	UNP A3ZEY5
A	517	GLY	-	expression tag	UNP A3ZEY5
A	518	GLY	-	expression tag	UNP A3ZEY5
A	519	HIS	-	expression tag	UNP A3ZEY5
A	520	HIS	-	expression tag	UNP A3ZEY5
A	521	HIS	-	expression tag	UNP A3ZEY5
A	522	HIS	-	expression tag	UNP A3ZEY5
A	523	HIS	-	expression tag	UNP A3ZEY5
A	524	HIS	-	expression tag	UNP A3ZEY5
A	525	HIS	-	expression tag	UNP A3ZEY5
A	526	HIS	-	expression tag	UNP A3ZEY5
B	202	MET	-	initiating methionine	UNP A3ZEY5
B	513	GLY	-	expression tag	UNP A3ZEY5
B	514	GLY	-	expression tag	UNP A3ZEY5
B	515	GLY	-	expression tag	UNP A3ZEY5
B	516	GLY	-	expression tag	UNP A3ZEY5
B	517	GLY	-	expression tag	UNP A3ZEY5
B	518	GLY	-	expression tag	UNP A3ZEY5
B	519	HIS	-	expression tag	UNP A3ZEY5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	520	HIS	-	expression tag	UNP A3ZEY5
B	521	HIS	-	expression tag	UNP A3ZEY5
B	522	HIS	-	expression tag	UNP A3ZEY5
B	523	HIS	-	expression tag	UNP A3ZEY5
B	524	HIS	-	expression tag	UNP A3ZEY5
B	525	HIS	-	expression tag	UNP A3ZEY5
B	526	HIS	-	expression tag	UNP A3ZEY5
C	202	MET	-	initiating methionine	UNP A3ZEY5
C	513	GLY	-	expression tag	UNP A3ZEY5
C	514	GLY	-	expression tag	UNP A3ZEY5
C	515	GLY	-	expression tag	UNP A3ZEY5
C	516	GLY	-	expression tag	UNP A3ZEY5
C	517	GLY	-	expression tag	UNP A3ZEY5
C	518	GLY	-	expression tag	UNP A3ZEY5
C	519	HIS	-	expression tag	UNP A3ZEY5
C	520	HIS	-	expression tag	UNP A3ZEY5
C	521	HIS	-	expression tag	UNP A3ZEY5
C	522	HIS	-	expression tag	UNP A3ZEY5
C	523	HIS	-	expression tag	UNP A3ZEY5
C	524	HIS	-	expression tag	UNP A3ZEY5
C	525	HIS	-	expression tag	UNP A3ZEY5
C	526	HIS	-	expression tag	UNP A3ZEY5

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	7	Total O 7 7	0	0
3	B	10	Total O 10 10	0	0
3	C	1	Total O 1 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: UPF0141 protein yjdB



- Molecule 1: UPF0141 protein yjdB



- Molecule 1: UPF0141 protein yjdB



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.85Å 183.14Å 121.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.58 – 2.40 54.58 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (54.58-2.40) 99.6 (54.58-2.40)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.59 (at 2.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.215 , 0.256 0.213 , 0.255	Depositor DCC
$R_{free}$ test set	2700 reflections (5.08%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.4	Xtriage
Anisotropy	0.729	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 35.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7411	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/2512	0.62	1/3396 (0.0%)
1	B	0.48	0/2501	0.62	1/3382 (0.0%)
1	C	0.45	0/2512	0.60	1/3396 (0.0%)
All	All	0.46	0/7525	0.62	3/10174 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	508	LEU	CA-CB-CG	5.78	128.59	115.30
1	C	425	LEU	CA-CB-CG	5.34	127.59	115.30
1	B	482	LEU	CA-CB-CG	5.00	126.81	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2467	0	2390	19	0
1	B	2456	0	2377	19	0
1	C	2467	0	2390	25	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
3	A	7	0	0	0	0
3	B	10	0	0	0	0
3	C	1	0	0	0	0
All	All	7411	0	7157	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:413:LYS:HA	1:A:416:LYS:HE3	1.62	0.82
1:B:505:ASN:OD1	1:B:507:LYS:HG3	1.88	0.73
1:A:217:THR:O	1:A:347:GLN:NE2	2.27	0.68
1:C:503:LEU:HA	1:C:508:LEU:HD22	1.80	0.64
1:C:212:ARG:HG3	1:C:298:THR:HG23	1.80	0.63
1:C:217:THR:O	1:C:347:GLN:NE2	2.31	0.63
1:A:212:ARG:HD3	1:A:298:THR:HG22	1.80	0.63
1:B:217:THR:O	1:B:347:GLN:NE2	2.34	0.60
1:B:256:PHE:HB2	1:B:456:ALA:HB3	1.84	0.60
1:C:294:VAL:O	1:C:298:THR:HB	2.02	0.59
1:A:219:LYS:HB2	1:A:347:GLN:HG3	1.86	0.58
1:A:387:SER:O	1:A:391:ILE:HG12	2.05	0.56
1:B:500:GLU:OE1	1:B:500:GLU:N	2.33	0.56
1:C:298:THR:HG21	1:C:492:VAL:HG22	1.88	0.54
1:A:367:PRO:HB2	1:A:369:GLU:OE1	2.09	0.53
1:B:210:ALA:HA	1:B:493:LYS:O	2.08	0.53
1:B:500:GLU:HA	1:B:508:LEU:HA	1.90	0.52
1:B:503:LEU:HA	1:B:508:LEU:HD22	1.93	0.51
1:B:502:ASP:HB3	1:B:505:ASN:HB3	1.93	0.50
1:A:488:GLY:HA3	1:A:502:ASP:OD1	2.11	0.49
1:C:307:ASN:HB2	1:C:333:LEU:HD21	1.95	0.49
1:A:477:LEU:HD21	1:A:508:LEU:HD11	1.95	0.48
1:B:270:LEU:HB3	1:B:271:PRO:HD3	1.95	0.48
1:C:316:CYS:HA	1:C:319:LEU:HD13	1.96	0.47
1:B:202:MET:HG2	1:B:203:PHE:N	2.29	0.47
1:A:206:ILE:HG12	1:A:289:GLU:HB2	1.97	0.47
1:C:212:ARG:HD3	1:C:298:THR:O	2.14	0.47
1:A:256:PHE:HB2	1:A:456:ALA:HB3	1.95	0.46
1:B:298:THR:HG21	1:B:492:VAL:HG22	1.97	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:206:ILE:O	1:C:297:LYS:NZ	2.47	0.46
1:C:324:LYS:HE2	1:C:324:LYS:HB3	1.75	0.46
1:C:304:TRP:CZ3	1:C:351:VAL:HG11	2.51	0.46
1:C:270:LEU:HB3	1:C:271:PRO:HD3	1.97	0.46
1:C:253:VAL:HG12	1:C:459:TRP:HB2	1.99	0.45
1:C:405:LEU:HD22	1:C:422:LEU:HD21	1.97	0.45
1:A:270:LEU:HB3	1:A:271:PRO:HD3	1.99	0.45
1:B:405:LEU:HD22	1:B:422:LEU:HD21	2.00	0.44
1:B:464:LYS:HG2	1:B:464:LYS:O	2.16	0.44
1:C:212:ARG:HG3	1:C:298:THR:CG2	2.47	0.44
1:C:387:SER:O	1:C:391:ILE:HG12	2.18	0.44
1:C:212:ARG:NH2	1:C:347:GLN:OE1	2.45	0.44
1:C:256:PHE:CD2	1:C:473:LYS:HA	2.54	0.43
1:A:298:THR:HG21	1:A:492:VAL:HG22	2.00	0.43
1:A:472:HIS:CD2	1:A:508:LEU:HD12	2.54	0.43
1:A:468:LEU:O	1:A:472:HIS:ND1	2.52	0.42
1:C:472:HIS:CG	1:C:508:LEU:HD23	2.54	0.42
1:B:216:HIS:HB2	1:B:347:GLN:HE22	1.84	0.42
1:B:239:TYR:CZ	1:B:241:LYS:HB3	2.54	0.42
1:C:505:ASN:HA	1:C:506:PRO:HD3	1.82	0.42
1:C:381:GLU:OE1	1:C:384:LYS:HD2	2.19	0.42
1:C:367:PRO:HG2	1:C:370:PHE:HD2	1.85	0.42
1:C:502:ASP:HB3	1:C:505:ASN:HB3	2.02	0.42
1:A:458:PHE:CZ	1:A:460:SER:HB2	2.55	0.42
1:B:293:ASP:O	1:B:297:LYS:HG3	2.20	0.41
1:A:467:ASN:O	1:A:471:GLU:HG3	2.20	0.41
1:B:416:LYS:H	1:B:416:LYS:HG3	1.39	0.41
1:A:242:ASN:HB3	1:A:373:PHE:CZ	2.56	0.41
1:B:219:LYS:NZ	1:B:491:ASN:OD1	2.54	0.41
1:C:242:ASN:HB3	1:C:373:PHE:CZ	2.56	0.41
1:C:206:ILE:HG12	1:C:289:GLU:HB2	2.03	0.40
1:A:304:TRP:CZ3	1:A:351:VAL:HG11	2.57	0.40
1:A:481:ASN:ND2	1:A:501:TYR:HD2	2.20	0.40
1:B:273:MET:SD	1:B:425:LEU:HD21	2.62	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/325 (93%)	292 (96%)	11 (4%)	0	100	100
1	B	302/325 (93%)	292 (97%)	10 (3%)	0	100	100
1	C	303/325 (93%)	295 (97%)	8 (3%)	0	100	100
All	All	908/975 (93%)	879 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/285 (96%)	271 (99%)	3 (1%)	80	92
1	B	273/285 (96%)	269 (98%)	4 (2%)	72	87
1	C	274/285 (96%)	267 (97%)	7 (3%)	54	74
All	All	821/855 (96%)	807 (98%)	14 (2%)	68	85

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	HIS
1	A	284	SER
1	A	329	LEU
1	B	329	LEU
1	B	416	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	425	LEU
1	B	439	LEU
1	C	211	TYR
1	C	217	THR
1	C	257	ASP
1	C	324	LYS
1	C	329	LEU
1	C	425	LEU
1	C	442	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	281	ASN
1	C	343	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPO	A	266	1,2	7,10,11	0.66	0	10,14,16	1.30	1 (10%)
1	TPO	B	266	1,2	7,10,11	0.72	0	10,14,16	1.62	2 (20%)
1	TPO	C	266	1,2	7,10,11	0.92	0	10,14,16	1.73	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	266	1,2	-	0/8/11/13	0/0/0/0
1	TPO	B	266	1,2	-	0/8/11/13	0/0/0/0
1	TPO	C	266	1,2	-	0/8/11/13	0/0/0/0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	266	TPO	O3P-P-OG1	-2.84	98.13	106.62
1	C	266	TPO	CG2-CB-CA	-2.53	108.08	113.15
1	B	266	TPO	O-C-CA	-2.45	119.00	125.69
1	B	266	TPO	CG2-CB-CA	-2.19	108.77	113.15
1	A	266	TPO	CG2-CB-CA	-2.01	109.13	113.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/325 (94%)	-0.15	3 (0%) 84 83	26, 49, 73, 86	0
1	B	306/325 (94%)	-0.15	1 (0%) 94 94	26, 45, 76, 83	0
1	C	307/325 (94%)	-0.11	1 (0%) 94 94	30, 48, 76, 87	0
All	All	920/975 (94%)	-0.14	5 (0%) 91 91	26, 48, 75, 87	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	321	TYR	5.6
1	A	321	TYR	3.6
1	A	202	MET	2.2
1	B	504	LEU	2.2
1	A	220	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	B	266	11/12	0.95	0.13	-	33,40,50,55	0
1	TPO	A	266	11/12	0.95	0.12	-	32,36,41,46	0
1	TPO	C	266	11/12	0.94	0.10	-	41,43,55,57	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	601	1/1	0.99	0.11	-1.32	33,33,33,33	1
2	ZN	B	601	1/1	0.96	0.10	-4.74	34,34,34,34	1
2	ZN	C	601	1/1	0.97	0.04	-7.91	38,38,38,38	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.