



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:26 PM GMT

PDB ID : 1TOK  
Title : Maleic acid-bound structure of SRHEPT mutant of E. coli aspartate aminotransferase  
Authors : Chow, M.A.; McElroy, K.E.; Corbett, K.D.; Berger, J.M.; Kirsch, J.F.  
Deposited on : 2004-06-14  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	<div> <div></div> <div>85%12%•</div> </div>
1	B	396	<div> <div>2%</div> <div>80%15%••</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6463 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	396	Total	C	N	O	P	S	0	0	0
			3082	1942	535	591	1	13			
1	B	396	Total	C	N	O	P	S	0	0	0
			3082	1942	535	591	1	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	THR	ALA	ENGINEERED	UNP P00509
A	13	THR	PRO	ENGINEERED	UNP P00509
A	34	ASP	ASN	ENGINEERED	UNP P00509
A	109	SER	THR	ENGINEERED	UNP P00509
A	258	LLP	LYS	MODIFIED RESIDUE	UNP P00509
A	261	ALA	GLY	ENGINEERED	UNP P00509
A	285	GLY	SER	ENGINEERED	UNP P00509
A	297	SER	ASN	ENGINEERED	UNP P00509
B	12	THR	ALA	ENGINEERED	UNP P00509
B	13	THR	PRO	ENGINEERED	UNP P00509
B	34	ASP	ASN	ENGINEERED	UNP P00509
B	109	SER	THR	ENGINEERED	UNP P00509
B	258	LLP	LYS	MODIFIED RESIDUE	UNP P00509
B	261	ALA	GLY	ENGINEERED	UNP P00509
B	285	GLY	SER	ENGINEERED	UNP P00509
B	297	SER	ASN	ENGINEERED	UNP P00509

- Molecule 2 is MALEIC ACID (three-letter code: MAE) (formula: C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	4	4		
2	B	1	Total	C	O	0	0
			8	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	154	Total	O	0	0
			154	154		
3	B	129	Total	O	0	0
			129	129		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate aminotransferase



- Molecule 1: Aspartate aminotransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.14 Å   140.14 Å   81.08 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	30.00 – 1.85 48.58 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-1.85) 99.9 (48.58-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.04 (at 1.86 Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.194 , 0.227 0.195 , 0.225	Depositor DCC
$R_{free}$ test set	3876 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.7	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.7	EDS
Estimated twinning fraction	0.023 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 77185 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6463	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, MAE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/3117	0.77	2/4221 (0.0%)
1	B	0.56	0/3117	0.82	12/4221 (0.3%)
All	All	0.55	0/6234	0.79	14/8442 (0.2%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	400	LEU	CA-CB-CG	9.56	137.28	115.30
1	B	241	ARG	NE-CZ-NH1	7.43	124.02	120.30
1	B	113	ARG	NE-CZ-NH2	-7.31	116.64	120.30
1	A	112	LEU	CB-CG-CD1	6.93	122.79	111.00
1	B	144	LYS	CD-CE-NZ	6.60	126.87	111.70
1	B	276	ASP	CB-CG-OD2	6.03	123.73	118.30
1	B	114	VAL	CG1-CB-CG2	5.85	120.25	110.90
1	B	113	ARG	NE-CZ-NH1	5.77	123.19	120.30
1	B	241	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	B	15	ASP	CB-CG-OD2	5.54	123.29	118.30
1	A	250	LEU	CA-CB-CG	5.41	127.74	115.30
1	B	119	LEU	CB-CG-CD1	5.22	119.88	111.00
1	B	112	LEU	CB-CG-CD1	5.18	119.80	111.00
1	B	117	ASP	CB-CG-OD2	5.08	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3082	0	3021	40	0
1	B	3082	0	3020	51	0
2	A	8	0	2	1	0
2	B	8	0	2	0	0
3	A	154	0	0	3	0
3	B	129	0	0	10	0
All	All	6463	0	6045	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:ASN:C	1:A:137:ASN:HD22	1.61	1.03
1:B:393:THR:HG21	3:B:442:HOH:O	1.65	0.96
1:A:366:THR:HG22	1:A:369:GLN:H	1.33	0.94
1:B:393:THR:HG22	1:B:396:ASN:H	1.36	0.88
1:A:79:PHE:O	1:A:83:THR:HG23	1.83	0.79
1:A:57:GLU:OE2	1:A:301:HIS:HE1	1.64	0.78
1:A:294:ASN:HD21	1:B:294:ASN:HD21	1.30	0.76
1:A:301:HIS:HD2	3:A:457:HOH:O	1.68	0.76
1:A:137:ASN:ND2	1:A:137:ASN:C	2.37	0.74
1:B:203:GLU:HG3	3:B:469:HOH:O	1.87	0.73
1:B:87:LEU:O	1:B:241:ARG:HD2	1.87	0.73
1:B:78:GLU:OE1	1:B:81:ARG:NH2	2.17	0.72
1:B:168:LEU:HD21	1:B:173:LEU:HD23	1.73	0.71
1:A:99:ARG:HD2	1:A:274:ALA:O	1.91	0.70
1:B:409:LEU:CD1	3:B:501:HOH:O	2.41	0.69
1:A:388:ASN:HD21	1:A:390:ALA:HB3	1.58	0.69
1:A:99:ARG:HD3	1:A:275:ALA:O	1.93	0.67
1:B:230:ARG:HD3	3:B:470:HOH:O	1.96	0.65
1:A:312:ASN:HD22	1:A:315:LEU:H	1.44	0.65
1:A:12:THR:HG22	1:B:286:GLN:NE2	2.12	0.65
1:A:12:THR:HG22	1:B:286:GLN:CD	2.17	0.65

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:348:ARG:CD	3:B:535:HOH:O	2.46	0.64
1:A:144:LYS:NZ	1:A:148:ASN:HD21	1.95	0.63
1:B:19:GLY:O	1:B:23:LEU:HD23	2.01	0.60
1:B:334:ARG:NH2	1:B:360:PHE:O	2.34	0.60
1:B:210:GLN:HE21	1:B:214:GLU:HG3	1.68	0.59
1:B:23:LEU:CD2	3:B:474:HOH:O	2.50	0.59
1:B:69:ASN:HD22	1:B:70:TYR:N	2.03	0.57
3:A:488:HOH:O	1:B:12:THR:HG22	2.05	0.57
1:A:5:MET:N	1:A:7:GLU:OE2	2.38	0.56
1:A:126:LYS:NZ	1:A:129:ARG:HD3	2.21	0.56
1:B:348:ARG:HD2	3:B:535:HOH:O	2.05	0.56
1:B:393:THR:HG22	1:B:396:ASN:N	2.16	0.55
1:B:367:LYS:HE3	3:B:467:HOH:O	2.07	0.55
1:A:144:LYS:HZ3	1:A:148:ASN:HD21	1.54	0.54
1:B:339:ASN:O	1:B:342:GLN:HB2	2.06	0.54
1:B:350:PHE:HB3	1:B:353:ILE:HD12	1.88	0.53
1:B:405:VAL:HA	1:B:409:LEU:HD11	1.91	0.52
1:B:69:ASN:HD22	1:B:70:TYR:H	1.56	0.52
1:B:110:GLY:O	1:B:114:VAL:HG13	2.09	0.52
1:A:189:HIS:CD2	1:A:194:ASN:H	2.27	0.52
1:B:374:ARG:HG3	1:B:380:TYR:CE2	2.45	0.52
1:A:83:THR:HB	1:A:256:TYR:OH	2.10	0.51
1:B:324:THR:HB	1:B:327:ARG:NH2	2.26	0.51
1:A:258:LLP:H4'1	2:A:410:MAE:C3	2.40	0.51
1:B:392:MET:HG2	1:B:400:LEU:HD11	1.92	0.51
1:A:327:ARG:NH1	3:A:523:HOH:O	2.40	0.50
1:A:12:THR:CG2	1:B:286:GLN:CD	2.80	0.49
1:A:405:VAL:HA	1:A:409:LEU:HD22	1.96	0.48
1:A:22:ASP:OD1	1:A:25:ARG:NH2	2.47	0.47
1:A:41:LYS:HG3	1:A:391:GLY:HA2	1.96	0.47
1:B:404:ILE:O	1:B:409:LEU:HG	2.14	0.47
1:B:43:GLU:HG3	1:B:44:THR:HG23	1.97	0.47
1:A:264:ASN:HD22	1:A:264:ASN:C	2.16	0.47
1:A:312:ASN:ND2	1:A:315:LEU:H	2.11	0.46
1:A:187:LEU:HD13	1:A:188:PHE:N	2.31	0.46
1:B:113:ARG:HD2	1:B:117:ASP:OD2	2.15	0.45
1:A:189:HIS:HD2	1:A:194:ASN:H	1.62	0.45
1:B:194:ASN:HA	1:B:195:PRO:HA	1.82	0.45
1:B:292:ARG:HA	1:B:296:SER:HA	1.99	0.45
1:A:312:ASN:HD22	1:A:315:LEU:HB2	1.82	0.44
1:B:162:ASP:HB2	1:B:169:ASP:HB2	1.99	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:ASN:HB2	1:A:167:THR:HG22	1.99	0.44
1:B:220:LEU:HD13	1:B:251:ILE:HB	1.99	0.44
1:B:256:TYR:HA	1:B:259:ASN:HD21	1.82	0.43
1:A:162:ASP:OD2	1:A:165:ASN:ND2	2.48	0.43
1:A:187:LEU:C	1:A:187:LEU:HD13	2.37	0.43
1:A:256:TYR:HA	1:A:259:ASN:HD21	1.84	0.43
1:B:35:LEU:HB3	1:B:387:VAL:HG13	2.01	0.43
1:B:113:ARG:NH2	3:B:423:HOH:O	2.48	0.43
1:A:347:ASN:OD1	1:A:347:ASN:N	2.51	0.43
1:B:335:GLN:O	1:B:338:VAL:HG22	2.19	0.42
1:B:112:LEU:HD13	1:B:253:ALA:CB	2.50	0.42
1:B:203:GLU:HA	1:B:206:GLN:HE21	1.83	0.42
1:A:61:LEU:HD12	1:B:61:LEU:HD12	2.02	0.42
1:A:194:ASN:HA	1:A:195:PRO:HA	1.90	0.42
1:B:158:TYR:CD1	1:B:173:LEU:HD13	2.55	0.41
1:B:393:THR:HG23	1:B:395:ASP:H	1.85	0.41
1:B:320:GLU:O	1:B:324:THR:HG22	2.21	0.41
1:A:43:GLU:H	1:A:43:GLU:CD	2.24	0.41
1:B:210:GLN:NE2	1:B:214:GLU:HG3	2.34	0.41
1:B:23:LEU:HD21	3:B:474:HOH:O	2.19	0.40
1:A:112:LEU:HD13	1:A:253:ALA:CB	2.51	0.40
1:A:126:LYS:HZ2	1:A:129:ARG:HD3	1.86	0.40
1:B:112:LEU:HB3	1:B:146:VAL:HG11	2.04	0.40
1:B:212:SER:OG	1:B:247:HIS:CE1	2.75	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/396 (99%)	385 (98%)	8 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	393/396 (99%)	382 (97%)	11 (3%)	0	100	100
All	All	786/792 (99%)	767 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/319 (100%)	295 (92%)	24 (8%)	17	4
1	B	319/319 (100%)	294 (92%)	25 (8%)	16	3
All	All	638/638 (100%)	589 (92%)	49 (8%)	16	3

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	ARG
1	A	50	LEU
1	A	61	LEU
1	A	62	GLU
1	A	83	THR
1	A	87	LEU
1	A	112	LEU
1	A	126	LYS
1	A	137	ASN
1	A	152	LEU
1	A	178	ASN
1	A	202	LEU
1	A	223	PHE
1	A	250	LEU
1	A	259	ASN
1	A	262	LEU
1	A	264	ASN
1	A	272	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	327	ARG
1	A	334	ARG
1	A	341	LEU
1	A	366	THR
1	A	370	VAL
1	A	409	LEU
1	B	13	THR
1	B	20	LEU
1	B	41	LYS
1	B	61	LEU
1	B	69	ASN
1	B	87	LEU
1	B	112	LEU
1	B	119	LEU
1	B	137	ASN
1	B	173	LEU
1	B	179	GLU
1	B	220	LEU
1	B	223	PHE
1	B	240	LEU
1	B	259	ASN
1	B	273	VAL
1	B	294	ASN
1	B	324	THR
1	B	342	GLN
1	B	367	LYS
1	B	371	LEU
1	B	373	LEU
1	B	393	THR
1	B	400	LEU
1	B	409	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	84	GLN
1	A	137	ASN
1	A	148	ASN
1	A	175	ASN
1	A	189	HIS
1	A	206	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	226	GLN
1	A	259	ASN
1	A	264	ASN
1	A	301	HIS
1	A	312	ASN
1	A	328	GLN
1	A	331	GLN
1	A	335	GLN
1	A	339	ASN
1	A	357	ASN
1	A	388	ASN
1	B	63	ASN
1	B	69	ASN
1	B	84	GLN
1	B	96	ASN
1	B	137	ASN
1	B	148	ASN
1	B	175	ASN
1	B	206	GLN
1	B	210	GLN
1	B	226	GLN
1	B	247	HIS
1	B	259	ASN
1	B	294	ASN
1	B	328	GLN
1	B	331	GLN
1	B	339	ASN
1	B	357	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	LLP	A	258	1	23,24,25	2.09	6 (26%)	28,32,34	1.69	4 (14%)
1	LLP	B	258	1	23,24,25	2.28	6 (26%)	28,32,34	1.95	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	258	1	-	0/15/17/19	0/1/1/1
1	LLP	B	258	1	-	0/15/17/19	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	258	LLP	O3-C3	-5.71	1.23	1.37
1	A	258	LLP	O3-C3	-5.20	1.24	1.37
1	B	258	LLP	CG-CB	2.09	1.61	1.52
1	A	258	LLP	C6-N1	2.24	1.39	1.34
1	A	258	LLP	CD-CE	2.29	1.58	1.51
1	B	258	LLP	CD-CE	2.76	1.60	1.51
1	B	258	LLP	C4-C4'	3.45	1.52	1.46
1	A	258	LLP	C4'-NZ	3.51	1.38	1.27
1	B	258	LLP	C4'-NZ	3.53	1.38	1.27
1	A	258	LLP	C4-C4'	3.84	1.53	1.46
1	A	258	LLP	CE-NZ	4.57	1.56	1.46
1	B	258	LLP	CE-NZ	5.22	1.57	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	LLP	C4-C4'-NZ	-3.35	106.43	125.06
1	A	258	LLP	C5-C6-N1	-3.23	118.26	123.86
1	B	258	LLP	C4-C4'-NZ	-3.18	107.36	125.06
1	B	258	LLP	O-C-CA	-2.21	119.74	125.49
1	B	258	LLP	OP2-P-OP4	-2.20	100.22	106.56
1	A	258	LLP	O-C-CA	-2.17	119.83	125.49
1	B	258	LLP	OP4-C5'-C5	4.77	116.88	108.99
1	A	258	LLP	OP4-C5'-C5	5.29	117.74	108.99

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	258	LLP	CD-CE-NZ	7.07	122.56	110.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	258	LLP	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MAE	A	410	-	1,7,7	0.51	0	0,8,8	0.00	-
2	MAE	B	410	-	1,7,7	0.45	0	0,8,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAE	A	410	-	-	0/0/5/5	0/0/0/0
2	MAE	B	410	-	-	0/0/5/5	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	410	MAE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	395/396 (99%)	0.15	5 (1%) 79 79	17, 22, 31, 40	0
1	B	395/396 (99%)	-0.19	6 (1%) 76 76	14, 24, 35, 43	0
All	All	790/792 (99%)	-0.02	11 (1%) 78 78	14, 23, 34, 43	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	ASN	4.0
1	B	371	LEU	3.9
1	B	409	LEU	2.9
1	B	210	GLN	2.7
1	B	246	MET	2.4
1	A	164	GLU	2.4
1	A	291	ILE	2.3
1	A	210	GLN	2.2
1	B	28	GLU	2.1
1	B	90	LYS	2.1
1	A	13	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	A	258	24/25	0.95	0.12	-	21,23,32,34	0
1	LLP	B	258	24/25	0.96	0.10	-	17,22,30,34	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MAE	B	410	8/8	0.94	0.11	-0.02	24,27,29,29	0
2	MAE	A	410	8/8	0.97	0.07	-1.26	20,24,26,26	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.