



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:28 PM GMT

PDB ID : 1TTA
Title : THE X-RAY CRYSTAL STRUCTURE REFINEMENTS OF NORMAL HUMAN TRANSTHYRETIN AND THE AMYLOIDOGENIC VAL30MET VARIANT TO 1.7 ANGSTROMS RESOLUTION
Authors : Hamilton, J.A.; Steinrauf, L.K.; Braden, B.C.
Deposited on : 1992-11-02
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	127	<div> <div>11%</div> <div>68%</div> <div>20%</div> <div>10%</div> <div>.</div> </div>
1	B	127	<div> <div>12%</div> <div>67%</div> <div>22%</div> <div>11%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSTHYRETIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	127	Total	C	N	O	S	16	10	0
			988	627	161	196	4			
1	B	127	Total	C	N	O	S	22	8	0
			982	623	161	196	2			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	105	Total	O	0	2
			107	107		
2	B	79	Total	O	0	0
			79	79		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRANSTHYRETIN



• Molecule 1: TRANSTHYRETIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	43.80Å 86.22Å 65.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 1.70 5.04 – 1.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (5.00-1.70) 73.6 (5.04-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 1.70Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.168 , (Not available) 0.168 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.59 , 136.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 19882 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2156	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.12 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3748e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	3/1080 (0.3%)	1.64	11/1467 (0.7%)
1	B	1.11	2/1060 (0.2%)	1.42	7/1442 (0.5%)
All	All	1.17	5/2140 (0.2%)	1.54	18/2909 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	THR	C-N	9.61	1.50	1.33
1	B	1	GLY	C-N	7.21	1.48	1.34
1	A	63[A]	GLU	CD-OE2	6.95	1.33	1.25
1	A	63[B]	GLU	CD-OE2	6.95	1.33	1.25
1	B	63	GLU	CD-OE2	6.26	1.32	1.25

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	34	ARG	NE-CZ-NH1	15.57	128.08	120.30
1	A	74[A]	ASP	CB-CG-OD1	13.86	130.78	118.30
1	A	74[B]	ASP	CB-CG-OD1	13.86	130.78	118.30
1	A	74[A]	ASP	CB-CG-OD2	-13.85	105.84	118.30
1	A	74[B]	ASP	CB-CG-OD2	-13.85	105.84	118.30
1	B	74	ASP	CB-CG-OD1	8.25	125.72	118.30
1	A	9	LYS	N-CA-C	-7.30	91.29	111.00
1	B	34	ARG	CD-NE-CZ	6.46	132.64	123.60
1	B	114	TYR	CB-CG-CD1	-6.46	117.13	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	18	ASP	CB-CG-OD2	6.25	123.93	118.30
1	B	34	ARG	NE-CZ-NH2	-5.85	117.38	120.30
1	A	62	GLU	CB-CG-CD	5.73	129.68	114.20
1	A	21	ARG	CG-CD-NE	5.68	123.73	111.80
1	B	42	GLU	CA-CB-CG	5.55	125.60	113.40
1	A	103	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	66	GLU	CA-CB-CG	5.18	124.79	113.40
1	A	27	ASN	CB-CG-OD1	-5.07	111.47	121.60
1	A	39	ASP	CB-CG-OD2	-5.00	113.80	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1	GLY	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	988	0	953	78	0
1	B	982	0	961	45	0
2	A	107	0	0	23	0
2	B	79	0	0	12	0
All	All	2156	0	1914	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

All (123) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48[B]:LYS:NZ	2:A:327:HOH:O	1.57	1.28
1:A:7:GLU:HB2	1:A:10[B]:CYS:SG	1.76	1.24
1:A:74[B]:ASP:CG	2:A:714:HOH:O	1.76	1.21

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:PRO:O	2:A:640:HOH:O	1.66	1.13
1:A:76:LYS:HE2	1:A:80[B]:LYS:HE3	1.34	1.10
1:B:2:PRO:HD3	1:B:50:SER:HB3	1.33	1.10
1:A:3:THR:HG23	1:A:62:GLU:CB	1.85	1.07
1:A:3:THR:HG23	1:A:62:GLU:HB2	1.29	1.07
1:B:72:GLU:OE2	2:B:792:HOH:O	1.73	1.06
1:A:7:GLU:CB	1:A:10[B]:CYS:SG	2.44	1.06
1:A:76:LYS:HE2	1:A:80[B]:LYS:CE	1.90	1.01
1:A:8:SER:HA	2:A:752:HOH:O	1.63	0.97
1:A:74[B]:ASP:OD1	2:A:714:HOH:O	1.76	0.96
1:A:4:GLY:HA2	2:A:614:HOH:O	1.63	0.96
1:A:7:GLU:HB2	1:A:10[B]:CYS:HG	1.33	0.94
1:A:8:SER:CA	2:A:752:HOH:O	2.19	0.91
1:A:8:SER:CB	2:A:752:HOH:O	2.24	0.84
1:B:72:GLU:CD	2:B:792:HOH:O	2.14	0.80
1:A:4:GLY:CA	2:A:614:HOH:O	2.21	0.80
1:B:27[A]:ASN:ND2	2:B:704:HOH:O	2.14	0.79
1:A:4:GLY:CA	1:A:63[B]:GLU:HG3	2.14	0.77
1:B:72:GLU:HG2	2:B:792:HOH:O	1.85	0.76
1:B:85:SER:OG	2:B:711:HOH:O	2.04	0.75
1:B:13:MET:SD	1:B:15[B]:LYS:HG2	2.26	0.75
1:B:2:PRO:HG2	1:B:48:LYS:O	1.88	0.74
1:A:63[B]:GLU:HG2	2:A:721:HOH:O	1.87	0.74
1:A:74[B]:ASP:OD2	2:A:714:HOH:O	1.93	0.73
1:A:9:LYS:NZ	1:A:61:GLU:OE1	2.21	0.73
1:B:72:GLU:CG	2:B:792:HOH:O	2.36	0.73
1:A:76:LYS:CE	1:A:80[B]:LYS:HE3	2.16	0.72
1:B:5:THR:HG21	1:B:44:PHE:O	1.88	0.72
1:B:5:THR:HG22	1:B:45:ALA:HB2	1.71	0.72
1:A:8:SER:HB2	1:A:61:GLU:N	2.05	0.72
1:A:8:SER:HB2	1:A:61:GLU:H	1.54	0.70
1:A:7:GLU:HB3	1:A:10[B]:CYS:SG	2.30	0.70
1:A:3:THR:CG2	1:A:62:GLU:HB2	2.15	0.70
1:A:3:THR:HG23	1:A:62:GLU:HB3	1.71	0.70
1:A:117:SER:HB2	1:B:117[A]:SER:HB2	1.73	0.69
1:A:3:THR:OG1	1:A:62:GLU:CG	2.40	0.69
1:A:3:THR:OG1	1:A:62:GLU:HG2	1.92	0.69
1:A:2:PRO:O	1:A:62:GLU:HG2	1.92	0.68
1:A:119[B]:THR:HG22	1:B:115:SER:OG	1.94	0.67
1:A:4:GLY:HA2	1:A:63[B]:GLU:HG3	1.77	0.65
1:A:99:ASP:OD1	2:A:724:HOH:O	2.13	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:ARG:HD2	2:B:411:HOH:O	1.97	0.64
1:A:7:GLU:HB3	1:A:57:GLY:HA2	1.80	0.64
1:B:2:PRO:CD	1:B:50:SER:HB3	2.22	0.63
1:A:5:THR:OG1	2:A:790:HOH:O	2.15	0.62
1:B:2:PRO:HD2	2:B:779:HOH:O	1.97	0.62
1:A:76:LYS:HE2	1:A:80[B]:LYS:HE2	1.80	0.62
1:A:2:PRO:HD2	1:A:8:SER:HA	1.81	0.61
1:B:13:MET:SD	1:B:15[B]:LYS:CG	2.88	0.61
1:A:8:SER:CB	1:A:61:GLU:H	2.11	0.61
1:B:124:ASN:HB2	1:B:125:PRO:CD	2.30	0.61
1:B:2:PRO:HD3	1:B:50:SER:CB	2.22	0.61
1:B:124:ASN:HB2	1:B:125:PRO:HD2	1.83	0.61
1:A:6:GLY:O	1:A:59:THR:O	2.21	0.59
1:B:4:GLY:HA3	1:B:58:LEU:HA	1.85	0.58
1:A:3:THR:CG2	1:A:62:GLU:CB	2.72	0.58
1:B:5:THR:HG22	1:B:45:ALA:CB	2.32	0.57
1:B:13:MET:HB2	1:B:104:ARG:NH2	2.19	0.57
1:A:23:SER:HB2	1:A:24:PRO:HD2	1.87	0.57
1:A:125:PRO:O	1:A:126:LYS:CB	2.53	0.56
1:A:104[B]:ARG:HH12	1:A:104[B]:ARG:HH21	1.48	0.54
1:A:115:SER:HB2	1:B:119[A]:THR:HG23	1.89	0.54
1:B:74:ASP:OD1	2:B:709:HOH:O	2.19	0.54
1:A:60:THR:HB	2:A:752:HOH:O	2.08	0.54
1:B:124:ASN:CB	1:B:125:PRO:CD	2.85	0.54
1:B:5:THR:HG23	2:B:705:HOH:O	2.09	0.53
1:B:8:SER:HB3	1:B:60:THR:HG22	1.91	0.52
1:B:8:SER:OG	1:B:9:LYS:N	2.41	0.52
1:A:7:GLU:HA	1:A:59:THR:O	2.09	0.52
1:A:3:THR:OG1	1:A:62:GLU:HG3	2.10	0.51
1:A:2:PRO:HG2	1:A:8:SER:HA	1.92	0.50
1:A:36:ALA:HB2	1:A:42:GLU:HG3	1.94	0.49
1:B:62:GLU:OE1	1:B:62:GLU:HA	2.12	0.49
1:B:124:ASN:H	1:B:125:PRO:HD2	1.77	0.49
1:B:34:ARG:NH2	1:B:66:GLU:O	2.45	0.49
1:A:8:SER:OG	2:A:752:HOH:O	2.19	0.48
1:A:4:GLY:N	1:A:63[B]:GLU:HG3	2.28	0.48
1:A:2:PRO:CD	1:A:8:SER:HA	2.43	0.48
1:A:36:ALA:HB2	1:A:42:GLU:OE2	2.14	0.47
1:A:8:SER:HB3	1:A:11:PRO:HD3	1.95	0.47
1:B:21[B]:ARG:HH21	1:B:82:LEU:HD11	1.80	0.47
1:B:55:LEU:HD13	1:B:58:LEU:HD11	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:5:THR:O	1:B:5:THR:HG23	2.16	0.45
1:B:13:MET:HB2	1:B:104:ARG:CZ	2.47	0.45
1:A:27:ASN:ND2	2:A:209:HOH:O	2.49	0.45
1:B:5:THR:CG2	1:B:45:ALA:HB2	2.46	0.45
1:A:15:LYS:NZ	2:A:765:HOH:O	2.33	0.45
1:B:102:PRO:HB2	1:B:122:VAL:HG12	1.98	0.45
1:A:6:GLY:O	1:A:59:THR:N	2.49	0.44
1:B:92:GLU:HB2	2:B:700:HOH:O	2.18	0.44
1:A:66:GLU:HG2	2:A:717[A]:HOH:O	2.17	0.44
1:A:16:VAL:C	1:A:17[B]:LEU:HD12	2.38	0.44
1:A:48[A]:LYS:HE2	2:A:327:HOH:O	2.17	0.43
1:A:125:PRO:O	1:A:126:LYS:HB3	2.19	0.41
1:A:4:GLY:HA2	1:A:63[B]:GLU:CG	2.49	0.41
1:B:31:HIS:HD2	2:B:502:HOH:O	2.03	0.41
1:A:6:GLY:O	1:A:59:THR:C	2.59	0.41
1:B:8:SER:C	1:B:10:CYS:H	2.25	0.41
1:A:3:THR:CG2	2:A:665:HOH:O	2.70	0.40
1:B:124:ASN:N	1:B:125:PRO:HD2	2.36	0.40
1:A:23:SER:HB2	1:A:24:PRO:CD	2.51	0.40
1:A:2:PRO:O	1:A:3:THR:OG1	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/127 (106%)	126 (93%)	3 (2%)	6 (4%)	3	0
1	B	133/127 (105%)	123 (92%)	7 (5%)	3 (2%)	8	1
All	All	268/254 (106%)	249 (93%)	10 (4%)	9 (3%)	5	0

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	SER
1	A	125	PRO
1	A	126	LYS
1	B	125	PRO
1	A	2	PRO
1	B	9	LYS
1	A	10[A]	CYS
1	A	10[B]	CYS
1	B	2	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/105 (110%)	105 (91%)	10 (9%)	13	2
1	B	113/105 (108%)	102 (90%)	11 (10%)	10	2
All	All	228/210 (109%)	207 (91%)	21 (9%)	12	2

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	7	GLU
1	A	10[A]	CYS
1	A	10[B]	CYS
1	A	27	ASN
1	A	39	ASP
1	A	62	GLU
1	A	92	GLU
1	A	117	SER
1	A	127	GLU
1	B	8	SER
1	B	10	CYS
1	B	39	ASP
1	B	55	LEU
1	B	62	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	82	LEU
1	B	85	SER
1	B	86	PRO
1	B	104	ARG
1	B	117[A]	SER
1	B	117[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	ASN
1	A	56	HIS
1	B	90	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	126/127 (99%)	0.21	14 (11%) 7 8	11, 16, 41, 50	28 (22%)
1	B	125/127 (98%)	0.33	15 (12%) 6 6	11, 20, 47, 55	29 (23%)
All	All	251/254 (98%)	0.27	29 (11%) 6 7	11, 18, 46, 55	57 (22%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	GLY	15.9
1	A	1	GLY	10.8
1	A	5	THR	9.8
1	A	3	THR	8.8
1	A	9	LYS	8.0
1	B	5	THR	7.1
1	A	8	SER	6.9
1	B	2	PRO	6.8
1	B	8	SER	6.7
1	A	126	LYS	6.5
1	A	4	GLY	6.4
1	A	2	PRO	5.8
1	B	10	CYS	5.8
1	B	9	LYS	5.3
1	B	125	PRO	5.0
1	B	3	THR	4.6
1	B	7	GLU	4.2
1	A	6	GLY	3.8
1	B	101	GLY	3.8
1	B	102	PRO	3.7
1	A	7	GLU	3.7
1	A	125	PRO	2.8
1	B	124	ASN	2.7
1	B	4	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	10[A]	CYS	2.6
1	B	6	GLY	2.4
1	B	37	ALA	2.4
1	A	124	ASN	2.1
1	A	37	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.