



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:33 PM GMT

PDB ID : 3U1O
Title : THREE DIMENSIONAL STRUCTURE OF DE NOVO DESIGNED CYSTEINE ESTERASE ECH19, Northeast Structural Genomics Consortium Target OR49
Authors : Kuzin, A.; Su, M.; Lew, S.; Forouhar, F.; Seetharaman, J.; Daya, P.; Xiao, R.; Ciccocanti, C.; Richter, F.; Everett, J.K.; Acton, T.B.; Baker, D.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2011-09-30
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	<div> <div>2%</div> <div>78%</div> <div>18%</div> <div>• •</div> </div>
1	B	419	<div> <div>%</div> <div>76%</div> <div>19%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	421	-	-	-	X
2	SO4	A	422	-	-	X	X
2	SO4	A	423	-	-	-	X
2	SO4	A	424	-	-	-	X
3	NA	A	426	-	-	-	X
3	NA	B	421	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called De Novo design cysteine esterase ECH19.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	Se	0	1	0
			3295	2128	542	607	2	16			
1	B	407	Total	C	N	O	S	Se	0	4	0
			3288	2126	538	606	2	16			

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	3	Total	Na	0	0
			3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	229	Total	O	0	2
			231	231		
4	A	210	Total	O	0	0
			210	210		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:**
-
- 2% 78% 18%
- MSE G2 M11 M17 G18 R19 V22 T23 K32 L50 S51 E52 Q66 T61 E62 V65 M66 Q67 T68 L73 P74 K78 F87 E91 Q92 L93 K101 E102 I103 Q104 I115 S118 V119 G149 K150 E154 V164 G168 I177
- R175 S176 Y177 Q180 T186 K192 W207 M211 N214 N217 F218 S219 T220 K221 F226 G227 K228 S229 N230 P236 W242 M247 N248 N249 S250 T251 K254 V255 K261 K264 G268 D278 L284 M288 E313 A317 R322 G323 V324
- P325 L326 V331 L334 T340 K341 L350 K351 M352 E355 T362 W369 W372 K398 Q399 R402 I403 L404 K405 R406 A407 M408 R409 G410 S411 L412 GLU HIS HIS HIS HIS HIS HIS

- Chain B:
-
- 76% 19%
- •
- | Residue | Category |
|---------|----------|
| E154 | Yellow |
| E321 | Green |
| R322 | Green |
| G323 | Green |
| V324 | Green |
| F325 | Green |
| L326 | Orange |
| V331 | Yellow |
| L334 | Green |
| R335 | Green |
| T340 | Green |
| K341 | Green |
| D342 | Green |
| M352 | Yellow |
| H358 | Green |
| K359 | Green |
| M360 | Green |
| T361 | Green |
| T362 | Green |
| Y365 | Green |
| W369 | Red |
| Y384 | Green |
| G385 | Green |
| Q386 | Green |
| Q390 | Orange |
| K398 | Green |
| Q399 | Green |
| R402 | Green |
| I403 | Green |
| L404 | Green |
| K405 | Orange |
| R406 | Green |
| A407 | Green |
| M408 | Green |
| R409 | Red |
| GLY | Grey |
| SER | Grey |
| LEU | Grey |
| GLU | Grey |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |
| HIS | Grey |
| MSE | Grey |
| GLY | Grey |
| S3 | Green |
| S12 | Yellow |
| M17 | Green |
| G18 | Green |
| R19 | Yellow |
| V22 | Yellow |
| T23 | Orange |
| L27 | Yellow |
| K32 | Yellow |
| K40 | Yellow |
| A41 | Green |
| E42 | Green |
| L50 | Yellow |
| L53 | Yellow |
| G59 | Red |
| G60 | Yellow |
| T61 | Yellow |
| T68 | Yellow |
| L73 | Yellow |
| P74 | Yellow |
| S77 | Yellow |
| K78 | Yellow |
| G82 | Yellow |
| L86 | Yellow |
| K90 | Yellow |
| L93 | Yellow |
| E102 | Yellow |
| I115 | Yellow |
| S118 | Yellow |
| V119 | Yellow |
| E137 | Orange |
| Y138 | Yellow |
| L146 | Orange |
| G149 | Yellow |
| V152 | Orange |

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	109.14Å 129.20Å 72.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.03 – 2.49 35.03 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.6 (35.03-2.49) 99.7 (35.03-2.49)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.201 , 0.262 0.188 , 0.249	Depositor DCC
R_{free} test set	1802 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	7 of 36184 reflections (0.019%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7058	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.51 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.4017e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3372	0.58	0/4551
1	B	0.44	0/3376	0.58	0/4555
All	All	0.43	0/6748	0.58	0/9106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3295	0	3233	62	0
1	B	3288	0	3232	69	0
2	A	25	0	0	5	0
2	B	5	0	0	0	0
3	A	3	0	0	0	0
3	B	1	0	0	0	0
4	A	210	0	0	1	0
4	B	231	0	0	2	0
All	All	7058	0	6465	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:MSE:HE1	1:A:211:MSE:SE	2.11	1.00
1:B:207:MSE:HE1	1:B:211:MSE:SE	2.16	0.96
1:A:398:LYS:HE3	1:A:402:ARG:HH21	1.34	0.91
1:B:23:THR:HG21	1:B:322:ARG:HH22	1.42	0.83
1:A:404:LEU:HD22	1:A:408:MSE:HE3	1.62	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/419 (98%)	396 (97%)	13 (3%)	1 (0%)	52	75
1	B	409/419 (98%)	397 (97%)	10 (2%)	2 (0%)	34	55
All	All	819/838 (98%)	793 (97%)	23 (3%)	3 (0%)	39	61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	409	ARG
1	B	230	ASN
1	B	359	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/342 (103%)	336 (96%)	16 (4%)	34	59
1	B	353/342 (103%)	331 (94%)	22 (6%)	23	41
All	All	705/684 (103%)	667 (95%)	38 (5%)	29	49

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	23	THR
1	B	53	LEU
1	B	390[A]	GLN
1	B	40	LYS
1	B	137[A]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	420	-	4,4,4	0.17	0	6,6,6	0.19	0
2	SO4	A	421	-	4,4,4	0.20	0	6,6,6	0.08	0
2	SO4	A	422	-	4,4,4	0.23	0	6,6,6	0.21	0
2	SO4	A	423	-	4,4,4	0.21	0	6,6,6	0.16	0
2	SO4	A	424	-	4,4,4	0.21	0	6,6,6	0.18	0
2	SO4	B	420	-	4,4,4	0.12	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	420	-	-	0/0/0/0	0/0/0/0
2	SO4	A	421	-	-	0/0/0/0	0/0/0/0
2	SO4	A	422	-	-	0/0/0/0	0/0/0/0
2	SO4	A	423	-	-	0/0/0/0	0/0/0/0
2	SO4	A	424	-	-	0/0/0/0	0/0/0/0
2	SO4	B	420	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	421	SO4	1	0
2	A	422	SO4	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/419 (94%)	-0.30	8 (2%) 68 72	9, 15, 39, 84	0
1	B	391/419 (93%)	-0.30	6 (1%) 76 79	9, 16, 37, 60	0
All	All	786/838 (93%)	-0.30	14 (1%) 71 75	9, 15, 38, 84	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	410	GLY	9.5
1	A	412	LEU	9.3
1	A	411	SER	8.4
1	A	2	GLY	7.0
1	B	409	ARG	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	A	423	5/5	0.85	0.38	11.69	90,93,94,94	0
2	SO4	A	421	5/5	0.91	0.61	11.09	92,94,95,97	0
2	SO4	A	422	5/5	0.82	0.41	8.20	94,97,99,99	0
2	SO4	A	424	5/5	0.74	0.34	6.44	95,97,99,100	0
3	NA	B	421	1/1	0.90	0.25	5.63	39,39,39,39	0
3	NA	A	426	1/1	0.96	0.25	5.46	36,36,36,36	0
3	NA	A	425	1/1	0.86	0.18	1.49	32,32,32,32	0
3	NA	A	427	1/1	0.91	0.13	-0.45	41,41,41,41	0
2	SO4	A	420	5/5	0.98	0.09	-	50,54,56,58	0
2	SO4	B	420	5/5	0.99	0.10	-	39,48,51,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.