



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 08:06 PM GMT

PDB ID : 4U67
Title : Crystal structure of the large ribosomal subunit (50S) of *Deinococcus radiodurans* containing a three residue insertion in L22
Authors : Wekselman, I.; Zimmerman, E.; Rozenberg, H.; Bashan, A.; Yonath, A.
Deposited on : 2014-07-28
Resolution : 3.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026982
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



























Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1010 (3.82-3.50)
Clashscore	102246	1125 (3.82-3.50)
Ramachandran outliers	100387	1079 (3.82-3.50)
Sidechain outliers	100360	1078 (3.82-3.50)
RSRZ outliers	91569	1017 (3.82-3.50)
RNA backbone	2183	1066 (4.52-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>5%</div> <div>45%</div> <div>39%</div> <div>11%</div> <div>5%</div> </div>
2	B	211	<div> <div>51%</div> <div>40%</div> <div>7%</div> <div>.</div> </div>
3	C	205	<div> <div>5%</div> <div>39%</div> <div>46%</div> <div>10%</div> <div>5%</div> </div>
4	D	180	<div> <div>13%</div> <div>53%</div> <div>41%</div> <div>.</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	185	
6	G	174	
7	H	134	
8	I	156	
9	J	141	
10	K	116	
11	L	114	
12	M	166	
13	N	118	
14	O	100	
15	P	137	
16	Q	95	
17	R	114	
18	S	237	
19	T	91	
20	U	81	
21	V	67	
22	W	55	
23	Z	60	
24	1	55	
25	2	47	
26	3	66	
27	X	2880	
28	Y	123	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
29	MG	K	201	-	-	-	X
29	MG	X	2901	-	-	-	X
29	MG	X	2902	-	-	-	X
29	MG	X	2905	-	-	-	X
29	MG	X	2907	-	-	-	X
29	MG	X	2909	-	-	-	X
29	MG	X	2910	-	-	-	X
29	MG	X	2912	-	-	-	X
29	MG	X	2913	-	-	-	X
29	MG	X	2917	-	-	-	X
29	MG	X	2919	-	-	-	X
29	MG	X	2920	-	-	-	X
29	MG	X	2921	-	-	-	X
29	MG	X	2922	-	-	-	X
29	MG	X	2924	-	-	-	X
29	MG	X	2926	-	-	-	X
29	MG	X	2928	-	-	-	X
29	MG	X	2929	-	-	-	X
29	MG	X	2931	-	-	-	X
29	MG	X	2934	-	-	-	X
29	MG	X	2936	-	-	-	X
29	MG	X	2939	-	-	-	X
29	MG	X	2940	-	-	-	X
29	MG	X	2943	-	-	-	X
29	MG	X	2944	-	-	-	X
29	MG	X	2947	-	-	-	X

2 Entry composition [i](#)

There are 29 unique types of molecules in this entry. The entry contains 83768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			1987	1235	399	350	3			

- Molecule 2 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	205	Total	C	N	O	S	0	0	0
			1539	965	295	271	8			

- Molecule 3 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	194	Total	C	N	O	S	0	0	0
			1481	920	284	275	2			

- Molecule 4 is a protein called 50S ribosomal protein L5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	177	Total	C	N	O	S	0	0	0
			1400	892	247	254	7			

- Molecule 5 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	171	Total	C	N	O	S	0	0	0
			1286	812	237	236	1			

- Molecule 6 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	142	Total	C	N	O	S	0	0	0
			1114	704	209	198	3			

- Molecule 7 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	134	Total	C	N	O	S	0	0	0
			997	614	198	180	5			

- Molecule 8 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	I	134	Total	C	N	O		0	0	0
			1011	619	206	186				

- Molecule 9 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	J	136	Total	C	N	O	S	0	0	0
			1090	696	202	185	7			

- Molecule 10 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	K	113	Total	C	N	O	S	0	0	0
			878	541	178	157	2			

- Molecule 11 is a protein called 50S ribosomal protein L18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	L	104	Total	C	N	O		0	0	0
			779	476	161	142				

- Molecule 12 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	M	108	Total	C	N	O		0	0	0
			871	543	172	156				

- Molecule 13 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	N	117	Total	C	N	O	S	0	0	0
			978	608	210	159	1			

- Molecule 14 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
14	O	94	Total	C	N	O			
			741	465	139	137	0	0	0

- Molecule 15 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	P	130	Total	C	N	O	S			
			1038	655	205	176	2	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	110	VAL	-	insertion	UNP Q9RXJ7
P	111	PRO	-	insertion	UNP Q9RXJ7
P	112	ARG	-	insertion	UNP Q9RXJ7

- Molecule 16 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
16	Q	93	Total	C	N	O	S			
			726	458	136	130	2	0	0	0

- Molecule 17 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
17	R	110	Total	C	N	O	S			
			825	513	160	151	1	0	0	0

- Molecule 18 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
18	S	175	Total	C	N	O	S			
			1345	849	236	254	6	0	0	0

- Molecule 19 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
19	T	74	Total	C	N	O	S			
			556	351	107	97	1	0	0	0

- Molecule 20 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
20	U	72	Total	C	N	O			
			552	341	116	95	0	0	0

- Molecule 21 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
21	V	65	Total	C	N	O	S			
			525	322	106	95	2	0	0	0

- Molecule 22 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
22	W	55	Total	C	N	O	S			
			424	264	82	76	2	0	0	0

- Molecule 23 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
23	Z	56	Total	C	N	O	S			
			443	272	91	75	5	0	0	0

- Molecule 24 is a protein called 50S ribosomal protein L33.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
24	1	53	Total	C	N	O	S			
			431	274	80	76	1	0	0	0

- Molecule 25 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
25	2	46	Total	C	N	O	S			
			383	230	91	60	2	0	0	0

- Molecule 26 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
26	3	59	Total	C	N	O	S			
			462	290	95	73	4	0	0	0

- Molecule 27 is a RNA chain called 23s RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
27	X	2667	Total	C	N	O	P	0	0	0
			57254	25538	10574	18475	2667			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	1526	U	UNK	conflict	GB 11612676

- Molecule 28 is a RNA chain called 5s RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
28	Y	122	Total	C	N	O	P	0	0	0
			2601	1161	476	842	122			


- Molecule 29 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
29	X	50	Total	Mg	0	0
			50	50		
29	K	1	Total	Mg	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 50S ribosomal protein L2




• Molecule 2: 50S ribosomal protein L3




• Molecule 3: 50S ribosomal protein L4







• Molecule 4: 50S ribosomal protein L5



• Molecule 5: 50S ribosomal protein L6



• Molecule 6: 50S ribosomal protein L13




• Molecule 7: 50S ribosomal protein L14







• Molecule 8: 50S ribosomal protein L15




• Molecule 9: 50S ribosomal protein L16




• Molecule 10: 50S ribosomal protein L17




• Molecule 11: 50S ribosomal protein L18




• Molecule 12: 50S ribosomal protein L19









• Molecule 13: 50S ribosomal protein L20





• Molecule 14: 50S ribosomal protein L21



• Molecule 15: 50S ribosomal protein L22




• Molecule 16: 50S ribosomal protein L23




• Molecule 17: 50S ribosomal protein L24






• Molecule 18: 50S ribosomal protein L25











• Molecule 28: 5s RNA



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	169.72Å 412.59Å 696.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 3.65 49.45 – 3.63	Depositor EDS
% Data completeness (in resolution range)	95.9 (19.99-3.65) 95.2 (49.45-3.63)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 3.67Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.226 , 0.270 0.231 , 0.273	Depositor DCC
R_{free} test set	12953 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	108.4	Xtriage
Anisotropy	0.636	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.17 , 17.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 261123 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	83768	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	0/2025	0.71	1/2726 (0.0%)
2	B	0.53	0/1567	0.78	0/2105
3	C	0.45	0/1504	0.70	1/2036 (0.0%)
4	D	0.28	0/1419	0.51	0/1903
5	E	0.30	0/1308	0.51	0/1771
6	G	0.49	0/1138	0.75	1/1539 (0.1%)
7	H	0.53	0/1007	0.75	0/1352
8	I	0.47	0/1022	0.73	0/1366
9	J	0.47	0/1113	0.74	0/1486
10	K	0.61	0/886	0.84	1/1188 (0.1%)
11	L	0.32	0/785	0.59	0/1048
12	M	0.57	0/884	0.86	1/1186 (0.1%)
13	N	0.46	0/994	0.66	0/1323
14	O	0.40	0/750	0.72	0/1000
15	P	0.54	0/1052	0.79	0/1409
16	Q	0.42	0/737	0.70	1/988 (0.1%)
17	R	0.43	0/835	0.73	0/1121
18	S	0.30	0/1370	0.56	0/1862
19	T	0.41	0/563	0.63	0/747
20	U	0.41	0/556	0.66	0/741
21	V	0.34	0/529	0.54	0/704
22	W	0.35	0/426	0.58	0/568
23	Z	0.51	0/455	0.78	0/611
24	1	0.47	0/438	0.71	0/583
25	2	0.43	0/387	0.75	1/509 (0.2%)
26	3	0.49	0/468	0.86	1/614 (0.2%)
27	X	0.59	3/64113 (0.0%)	1.17	296/99999 (0.3%)
28	Y	0.40	0/2907	0.94	2/4529 (0.0%)
All	All	0.55	3/91238 (0.0%)	1.07	306/137014 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying

if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
6	G	0	4
7	H	0	1
8	I	0	2
10	K	0	1
17	R	0	2
20	U	0	1
26	3	0	1
All	All	0	15

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
27	X	1	G	OP3-P	-10.66	1.48	1.61
27	X	774	A	N7-C5	-6.22	1.35	1.39
27	X	542	A	N9-C4	-5.92	1.34	1.37

The worst 5 of 306 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
27	X	1468	A	C8-N9-C4	-15.21	99.72	105.80
27	X	774	A	C8-N9-C4	-13.39	100.45	105.80
27	X	774	A	N7-C8-N9	11.68	119.64	113.80
27	X	1333	G	N3-C4-N9	-11.32	119.21	126.00
27	X	1468	A	N7-C8-N9	11.23	119.41	113.80

There are no chirality outliers.

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18	THR	Peptide
2	B	122	PHE	Peptide
2	B	146	THR	Peptide
6	G	108	GLY	Peptide
6	G	34	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1987	0	2056	130	0
2	B	1539	0	1600	101	0
3	C	1481	0	1504	101	0
4	D	1400	0	1481	62	0
5	E	1286	0	1336	35	0
6	G	1114	0	1144	63	0
7	H	997	0	1046	63	0
8	I	1011	0	1047	54	0
9	J	1090	0	1125	56	0
10	K	878	0	930	37	0
11	L	779	0	820	40	0
12	M	871	0	894	49	0
13	N	978	0	1020	72	0
14	O	741	0	756	46	0
15	P	1038	0	1125	78	0
16	Q	726	0	753	35	0
17	R	825	0	881	51	0
18	S	1345	0	1372	41	0
19	T	556	0	579	28	0
20	U	552	0	604	29	0
21	V	525	0	546	14	0
22	W	424	0	470	15	0
23	Z	443	0	444	27	0
24	1	431	0	456	30	0
25	2	383	0	414	27	0
26	3	462	0	506	52	0
27	X	57254	0	28850	1328	0
28	Y	2601	0	1327	54	0
29	K	1	0	0	0	0
29	X	50	0	0	0	0
All	All	83768	0	55086	2361	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 2361 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:GLY:H	1:A:217:ARG:HB2	1.20	1.04
2:B:136:ARG:HB3	27:X:1673:C:H5'	1.36	1.04
27:X:571:U:HO2'	27:X:581:A:H8	1.12	0.98
27:X:517:A:H5''	27:X:518:A:H5'	1.45	0.95
2:B:116:VAL:HG22	2:B:136:ARG:HG3	1.49	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/275 (94%)	223 (86%)	34 (13%)	1 (0%)	39	80
2	B	203/211 (96%)	182 (90%)	20 (10%)	1 (0%)	34	77
3	C	192/205 (94%)	165 (86%)	25 (13%)	2 (1%)	19	66
4	D	175/180 (97%)	153 (87%)	21 (12%)	1 (1%)	30	74
5	E	169/185 (91%)	155 (92%)	13 (8%)	1 (1%)	30	74
6	G	140/174 (80%)	126 (90%)	14 (10%)	0	100	100
7	H	132/134 (98%)	123 (93%)	8 (6%)	1 (1%)	24	70
8	I	132/156 (85%)	101 (76%)	28 (21%)	3 (2%)	8	52
9	J	134/141 (95%)	115 (86%)	19 (14%)	0	100	100
10	K	111/116 (96%)	102 (92%)	9 (8%)	0	100	100
11	L	102/114 (90%)	86 (84%)	16 (16%)	0	100	100
12	M	106/166 (64%)	101 (95%)	5 (5%)	0	100	100
13	N	115/118 (98%)	103 (90%)	10 (9%)	2 (2%)	11	57
14	O	92/100 (92%)	81 (88%)	11 (12%)	0	100	100
15	P	128/137 (93%)	109 (85%)	17 (13%)	2 (2%)	12	58
16	Q	91/95 (96%)	78 (86%)	11 (12%)	2 (2%)	8	52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
17	R	108/114 (95%)	86 (80%)	22 (20%)	0	100	100
18	S	173/237 (73%)	152 (88%)	21 (12%)	0	100	100
19	T	72/91 (79%)	61 (85%)	9 (12%)	2 (3%)	6	47
20	U	70/81 (86%)	52 (74%)	14 (20%)	4 (6%)	2	28
21	V	63/67 (94%)	59 (94%)	4 (6%)	0	100	100
22	W	53/55 (96%)	49 (92%)	4 (8%)	0	100	100
23	Z	54/60 (90%)	48 (89%)	6 (11%)	0	100	100
24	1	51/55 (93%)	37 (72%)	11 (22%)	3 (6%)	2	27
25	2	44/47 (94%)	40 (91%)	4 (9%)	0	100	100
26	3	57/66 (86%)	46 (81%)	11 (19%)	0	100	100
All	All	3025/3380 (90%)	2633 (87%)	367 (12%)	25 (1%)	24	70

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
16	Q	6	ILE
13	N	94	VAL
15	P	127	ILE
24	1	9	ILE
24	1	10	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/216 (94%)	158 (78%)	44 (22%)	1	9
2	B	155/157 (99%)	137 (88%)	18 (12%)	7	37
3	C	154/163 (94%)	126 (82%)	28 (18%)	2	15
4	D	153/156 (98%)	140 (92%)	13 (8%)	13	53
5	E	136/144 (94%)	122 (90%)	14 (10%)	9	43
6	G	118/146 (81%)	98 (83%)	20 (17%)	2	18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
7	H	103/103 (100%)	88 (85%)	15 (15%)	4	27
8	I	101/121 (84%)	83 (82%)	18 (18%)	2	16
9	J	110/115 (96%)	86 (78%)	24 (22%)	1	9
10	K	90/93 (97%)	75 (83%)	15 (17%)	3	19
11	L	74/82 (90%)	53 (72%)	21 (28%)	0	4
12	M	94/134 (70%)	71 (76%)	23 (24%)	1	7
13	N	96/97 (99%)	81 (84%)	15 (16%)	3	23
14	O	75/79 (95%)	65 (87%)	10 (13%)	5	31
15	P	112/118 (95%)	92 (82%)	20 (18%)	2	16
16	Q	75/76 (99%)	59 (79%)	16 (21%)	1	10
17	R	91/95 (96%)	78 (86%)	13 (14%)	4	28
18	S	149/192 (78%)	126 (85%)	23 (15%)	3	24
19	T	55/67 (82%)	49 (89%)	6 (11%)	8	41
20	U	57/66 (86%)	43 (75%)	14 (25%)	1	7
21	V	53/55 (96%)	46 (87%)	7 (13%)	5	31
22	W	48/48 (100%)	45 (94%)	3 (6%)	22	65
23	Z	50/53 (94%)	43 (86%)	7 (14%)	4	29
24	1	46/48 (96%)	33 (72%)	13 (28%)	0	4
25	2	39/40 (98%)	26 (67%)	13 (33%)	0	2
26	3	46/52 (88%)	37 (80%)	9 (20%)	1	12
All	All	2482/2716 (91%)	2060 (83%)	422 (17%)	2	18

5 of 422 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
9	J	133	VAL
12	M	25	PRO
24	1	21	TYR
10	K	9	LYS
11	L	17	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
13	N	37	GLN
15	P	16	GLN
23	Z	44	HIS
13	N	41	ASN
13	N	66	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
27	X	2657/2880 (92%)	611 (22%)	39 (1%)
28	Y	121/123 (98%)	31 (25%)	2 (1%)
All	All	2778/3003 (92%)	642 (23%)	41 (1%)

5 of 642 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
27	X	4	C
27	X	13	A
27	X	14	A
27	X	15	G
27	X	17	G

5 of 41 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
27	X	1141	U
27	X	1391	A
27	X	2756	A
27	X	1182	U
27	X	1225	G

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 51 ligands modelled in this entry, 51 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/275 (94%)	0.05	14 (5%) 29 18	45, 101, 165, 255	0
2	B	205/211 (97%)	-0.53	0 100 100	26, 43, 97, 208	0
3	C	194/205 (94%)	-0.13	10 (5%) 31 19	38, 101, 172, 256	0
4	D	177/180 (98%)	0.37	24 (13%) 4 3	108, 173, 225, 261	0
5	E	171/185 (92%)	-0.26	3 (1%) 71 54	51, 120, 177, 236	0
6	G	142/174 (81%)	-0.13	4 (2%) 56 39	35, 72, 151, 204	0
7	H	134/134 (100%)	-0.42	0 100 100	31, 42, 85, 136	0
8	I	134/156 (85%)	0.42	10 (7%) 17 10	53, 125, 199, 245	0
9	J	136/141 (96%)	-0.01	4 (2%) 55 37	56, 88, 168, 221	0
10	K	113/116 (97%)	-0.58	0 100 100	25, 30, 61, 99	0
11	L	104/114 (91%)	0.62	18 (17%) 2 1	128, 160, 197, 232	0
12	M	108/166 (65%)	-0.49	1 (0%) 85 74	29, 37, 93, 143	0
13	N	117/118 (99%)	-0.37	1 (0%) 85 74	40, 78, 123, 213	0
14	O	94/100 (94%)	-0.24	1 (1%) 82 68	51, 96, 164, 191	0
15	P	130/137 (94%)	-0.35	3 (2%) 64 46	31, 51, 149, 168	0
16	Q	93/95 (97%)	-0.35	1 (1%) 82 68	45, 83, 145, 192	0
17	R	110/114 (96%)	0.17	7 (6%) 23 12	66, 97, 187, 238	0
18	S	175/237 (73%)	-0.26	3 (1%) 73 56	89, 141, 204, 258	0
19	T	74/91 (81%)	0.16	3 (4%) 41 25	70, 110, 148, 231	0
20	U	72/81 (88%)	0.77	10 (13%) 4 3	73, 123, 178, 252	0
21	V	65/67 (97%)	-0.18	2 (3%) 52 35	67, 111, 168, 222	0
22	W	55/55 (100%)	0.32	5 (9%) 11 6	72, 94, 147, 177	0
23	Z	56/60 (93%)	-0.57	0 100 100	30, 36, 87, 100	0
24	1	53/55 (96%)	0.84	9 (16%) 2 2	98, 127, 189, 275	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
25	2	46/47 (97%)	-0.20	1 (2%) 65 47	39, 69, 102, 127	0
26	3	59/66 (89%)	0.83	8 (13%) 4 3	79, 99, 176, 252	0
27	X	2667/2880 (92%)	-0.54	22 (0%) 87 76	25, 73, 176, 304	0
28	Y	122/123 (99%)	-0.50	3 (2%) 61 43	74, 150, 189, 279	0
All	All	5866/6383 (91%)	-0.30	167 (2%) 56 39	25, 87, 183, 304	0

The worst 5 of 167 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
20	U	28	GLY	10.8
27	X	1086	C	9.1
27	X	1525	A	5.9
27	X	1734	C	5.8
27	X	1524	C	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
29	MG	X	2926	1/1	0.91	0.83	83.37	32,32,32,32	0
29	MG	X	2921	1/1	0.99	0.70	43.54	51,51,51,51	0
29	MG	X	2907	1/1	0.90	1.23	28.02	42,42,42,42	0
29	MG	X	2917	1/1	0.93	1.05	26.93	55,55,55,55	0
29	MG	X	2912	1/1	0.93	0.59	20.49	27,27,27,27	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
29	MG	X	2947	1/1	0.94	0.62	18.49	49,49,49,49	0
29	MG	X	2944	1/1	0.86	1.39	17.81	55,55,55,55	0
29	MG	X	2943	1/1	0.97	0.32	14.87	31,31,31,31	0
29	MG	X	2913	1/1	0.97	0.52	14.45	27,27,27,27	0
29	MG	X	2910	1/1	0.96	0.53	14.23	39,39,39,39	0
29	MG	X	2905	1/1	0.98	0.45	13.28	27,27,27,27	0
29	MG	X	2924	1/1	0.93	0.66	12.66	77,77,77,77	0
29	MG	X	2902	1/1	0.96	0.55	11.91	33,33,33,33	0
29	MG	X	2931	1/1	0.99	0.39	11.54	25,25,25,25	0
29	MG	X	2934	1/1	0.97	0.44	11.17	42,42,42,42	0
29	MG	X	2919	1/1	0.97	0.50	8.23	59,59,59,59	0
29	MG	K	201	1/1	0.86	0.70	8.22	25,25,25,25	0
29	MG	X	2940	1/1	0.96	0.66	8.15	54,54,54,54	0
29	MG	X	2929	1/1	0.96	0.45	7.53	37,37,37,37	0
29	MG	X	2928	1/1	0.98	0.29	6.92	36,36,36,36	0
29	MG	X	2939	1/1	0.97	0.31	6.08	32,32,32,32	0
29	MG	X	2920	1/1	0.97	0.32	4.47	41,41,41,41	0
29	MG	X	2901	1/1	0.96	0.32	4.34	37,37,37,37	0
29	MG	X	2936	1/1	0.97	0.21	3.46	33,33,33,33	0
29	MG	X	2909	1/1	0.91	0.53	2.90	51,51,51,51	0
29	MG	X	2922	1/1	0.95	0.35	2.27	30,30,30,30	0
29	MG	X	2923	1/1	0.96	0.38	-	33,33,33,33	0
29	MG	X	2903	1/1	0.94	0.60	-	30,30,30,30	0
29	MG	X	2950	1/1	0.92	0.84	-	37,37,37,37	0
29	MG	X	2927	1/1	0.91	0.16	-	69,69,69,69	0
29	MG	X	2908	1/1	0.97	0.49	-	31,31,31,31	0
29	MG	X	2904	1/1	0.92	0.40	-	29,29,29,29	0
29	MG	X	2906	1/1	0.92	0.88	-	27,27,27,27	0
29	MG	X	2916	1/1	0.93	1.10	-	26,26,26,26	0
29	MG	X	2915	1/1	0.93	0.37	-	28,28,28,28	0
29	MG	X	2911	1/1	0.96	0.87	-	40,40,40,40	0
29	MG	X	2938	1/1	0.97	0.40	-	27,27,27,27	0
29	MG	X	2945	1/1	0.97	0.46	-	61,61,61,61	0
29	MG	X	2935	1/1	0.90	0.33	-	74,74,74,74	0
29	MG	X	2930	1/1	0.96	0.45	-	30,30,30,30	0
29	MG	X	2925	1/1	0.98	0.54	-	32,32,32,32	0
29	MG	X	2949	1/1	0.98	0.22	-	32,32,32,32	0
29	MG	X	2937	1/1	0.97	0.38	-	36,36,36,36	0
29	MG	X	2933	1/1	0.95	0.34	-	55,55,55,55	0
29	MG	X	2946	1/1	0.97	0.69	-	27,27,27,27	0
29	MG	X	2914	1/1	0.98	0.55	-	26,26,26,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
29	MG	X	2941	1/1	0.86	0.78	-	115,115,115,115	0
29	MG	X	2932	1/1	0.97	0.47	-	39,39,39,39	0
29	MG	X	2948	1/1	0.93	0.23	-	48,48,48,48	0
29	MG	X	2918	1/1	0.99	0.56	-	36,36,36,36	0
29	MG	X	2942	1/1	0.96	0.33	-	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.