



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 6, 2016 – 05:10 PM EDT

PDB ID : 4U6P
Title : Structural mechanism of error-free bypass of major benzo[a]pyrene adduct by human polymerase kappa
Authors : Jha, V.K.; Ling, H.
Deposited on : 2014-07-29
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 67%; height: 10px; background-color: green;"></div> <div style="width: 13%; height: 10px; background-color: yellow;"></div> <div style="width: 18%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 67% 13% • 18% </div> </div>
1	B	526	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 70%; height: 10px; background-color: green;"></div> <div style="width: 11%; height: 10px; background-color: yellow;"></div> <div style="width: 18%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 70% 11% • 18% </div> </div>
2	C	9	<div> <div style="width: 67%; height: 10px; background-color: green;"></div> <div style="width: 33%; height: 10px; background-color: yellow;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 67% 33% </div>
2	P	9	<div> <div style="width: 67%; height: 10px; background-color: green;"></div> <div style="width: 33%; height: 10px; background-color: yellow;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 67% 33% </div>
3	D	13	<div> <div style="width: 62%; height: 10px; background-color: green;"></div> <div style="width: 38%; height: 10px; background-color: yellow;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 62% 38% </div>
3	T	13	<div> <div style="width: 69%; height: 10px; background-color: green;"></div> <div style="width: 31%; height: 10px; background-color: yellow;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 69% 31% </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	A	602	-	-	-	X
4	MG	B	603	-	-	-	X
5	DCT	B	605	-	-	X	-
6	PGE	B	601	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7767 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase kappa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	431	Total	C	N	O	S	0	0	0
			3376	2135	597	625	19			
1	B	431	Total	C	N	O	S	0	0	0
			3324	2107	585	612	20			

- Molecule 2 is a DNA chain called DNA (5'-D(P*GP*CP*GP*GP*AP*TP*CP*AP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	P	0	0	0
			184	87	36	52	9			
2	P	9	Total	C	N	O	P	0	0	0
			184	87	36	52	9			


- Molecule 3 is a DNA chain called DNA (5'-D(*TP*AP*TP*GP*GP*TP*GP*AP*TP*CP*CP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	13	Total	C	N	O	P	0	0	0
			263	127	47	77	12			
3	T	13	Total	C	N	O	P	0	0	0
			263	127	47	77	12			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Mg	0	0
			3	3		
4	A	3	Total	Mg	0	0
			3	3		

- Molecule 5 is 2',3'-DIDEOXYCYTIDINE 5'-TRIPHOSPHATE (three-letter code: DCT) (formula: C₉H₁₆N₃O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			27	9	3	12	3		
5	B	1	Total	C	N	O	P	0	0
			27	9	3	12	3		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	54	Total 54	O 54	0	0
7	B	25	Total 25	O 25	0	0
7	C	6	Total 6	O 6	0	0
7	D	8	Total 8	O 8	0	0
7	P	4	Total 4	O 4	0	0
7	T	6	Total 6	O 6	0	0

Chain C:  67% 33%




- Molecule 2: DNA (5'-D(P*GP*CP*GP*GP*AP*TP*CP*AP*C)-3')

Chain P:  67% 33%



- Molecule 3: DNA (5'-D(*TP*AP*TP*GP*GP*TP*GP*AP*TP*CP*CP*GP*C)-3')

Chain D:  62% 38%



- Molecule 3: DNA (5'-D(*TP*AP*TP*GP*GP*TP*GP*AP*TP*CP*CP*GP*C)-3')

Chain T:  69% 31%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.15Å 129.29Å 167.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.23 – 2.59 46.23 – 2.59	Depositor EDS
% Data completeness (in resolution range)	97.5 (46.23-2.59) 97.6 (46.23-2.59)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.58Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.200 , 0.239 0.200 , 0.237	Depositor DCC
R_{free} test set	1064 reflections (2.40%)	DCC
Wilson B-factor (Å ²)	60.3	Xtriage
Anisotropy	0.863	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7767	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PGE, DCT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3429	0.57	0/4622
1	B	0.41	0/3377	0.56	0/4559
2	C	0.59	0/206	1.05	0/315
2	P	0.71	0/206	1.24	0/315
3	D	0.66	0/294	1.10	0/453
3	T	0.69	0/294	1.34	0/453
All	All	0.45	0/7806	0.69	0/10717

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3376	0	3396	48	0
1	B	3324	0	3303	43	0
2	C	184	0	99	3	0
2	P	184	0	99	2	0
3	D	263	0	146	5	0
3	T	263	0	146	3	0
4	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	3	0	0	0	0
5	A	27	0	12	3	0
5	B	27	0	12	11	0
6	B	10	0	14	1	0
7	A	54	0	0	5	0
7	B	25	0	0	1	0
7	C	6	0	0	1	0
7	D	8	0	0	1	0
7	P	4	0	0	1	0
7	T	6	0	0	0	0
All	All	7767	0	7227	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 101 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:605:DCT:C5'	5:B:605:DCT:H6	2.00	0.92
2:P:13:DC:O2	7:P:101:HOH:O	1.95	0.83
1:A:507:ARG:NH2	3:D:4:DT:OP1	2.13	0.81
5:B:605:DCT:H5''	5:B:605:DCT:H6	1.62	0.80
1:B:214:TRP:O	1:B:219:ARG:NH1	2.23	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	427/526 (81%)	417 (98%)	9 (2%)	1 (0%)	52 77
1	B	427/526 (81%)	411 (96%)	15 (4%)	1 (0%)	52 77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	854/1052 (81%)	828 (97%)	24 (3%)	2 (0%)	52 77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	87	ASP
1	A	33	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	361/472 (76%)	343 (95%)	18 (5%)	30 56
1	B	347/472 (74%)	340 (98%)	7 (2%)	63 85
All	All	708/944 (75%)	683 (96%)	25 (4%)	43 71

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	402	LEU
1	A	434	LEU
1	B	506	LEU
1	A	422	PHE
1	A	448	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DCT	A	604	4	21,28,28	0.65	0	24,43,43	1.50	3 (12%)
6	PGE	B	601	-	9,9,9	0.57	0	8,8,8	0.42	0
5	DCT	B	605	4	21,28,28	0.95	1 (4%)	24,43,43	1.38	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DCT	A	604	4	-	0/18/31/31	0/2/2/2
6	PGE	B	601	-	-	0/7/7/7	0/0/0/0
5	DCT	B	605	4	-	0/18/31/31	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	605	DCT	C2-N3	-2.76	1.32	1.38

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	605	DCT	C2'-C1'-N1	-3.26	106.12	112.49
5	B	605	DCT	O4'-C4'-C5'	-2.81	105.28	109.52
5	A	604	DCT	C3'-C4'-C5'	-2.07	109.42	115.77
5	B	605	DCT	C6-C5-C4	2.05	118.24	117.44
5	A	604	DCT	C2'-C3'-C4'	2.36	106.91	102.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	604	DCT	3	0
6	B	601	PGE	1	0
5	B	605	DCT	11	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/526 (81%)	0.21	6 (1%) 78 74	31, 54, 85, 116	0
1	B	431/526 (81%)	0.25	13 (3%) 54 47	36, 58, 87, 114	0
2	C	9/9 (100%)	-0.16	0 100 100	39, 48, 53, 61	0
2	P	9/9 (100%)	0.03	0 100 100	46, 52, 57, 59	0
3	D	13/13 (100%)	0.18	0 100 100	38, 50, 73, 78	0
3	T	13/13 (100%)	0.03	0 100 100	41, 56, 67, 68	0
All	All	906/1096 (82%)	0.22	19 (2%) 67 61	31, 56, 86, 116	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	31	LEU	4.8
1	A	223	ILE	4.2
1	B	214	TRP	3.6
1	B	121	PRO	3.5
1	A	86	VAL	3.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	PGE	B	601	10/10	0.70	0.45	7.91	55,65,76,96	0
4	MG	A	602	1/1	0.87	0.29	4.06	49,49,49,49	0
4	MG	B	603	1/1	0.97	0.23	3.63	56,56,56,56	0
4	MG	B	602	1/1	0.87	0.22	1.82	34,34,34,34	0
4	MG	A	601	1/1	0.96	0.18	-0.09	21,21,21,21	0
5	DCT	B	605	27/27	0.94	0.17	-0.30	52,62,82,85	0
5	DCT	A	604	27/27	0.94	0.19	-0.57	43,53,69,76	0
4	MG	B	604	1/1	0.85	0.14	-	75,75,75,75	0
4	MG	A	603	1/1	0.81	0.15	-	65,65,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.