



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 2, 2017 – 01:57 PM EST

PDB ID : 5U72
Title : Structure of human MR1-5OH-DCF in complex with human MAIT A-F7 TCR
Authors : Keller, A.N.; Rossjohn, J.
Deposited on : 2016-12-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div><div>6%</div><div>84%</div><div>11%</div><div>.</div></div>
1	C	271	<div><div>3%</div><div>87%</div><div>9%</div><div>.</div></div>
2	B	203	<div><div>%</div><div>90%</div><div>8%</div><div>.</div></div>
2	D	203	<div><div>6%</div><div>88%</div><div>9%</div><div>.</div></div>
3	E	245	<div><div>4%</div><div>91%</div><div>8%</div><div>.</div></div>
3	G	245	<div><div>%</div><div>90%</div><div>9%</div><div>..</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	99	
4	H	99	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	F	101	-	-	-	X
6	GOL	A	303	-	-	-	X
8	DMS	C	302	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 13241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2094	1340	357	386	11			
1	C	263	Total	C	N	O	S	0	1	0
			2171	1391	372	397	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	conflict	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	conflict	UNP Q95460

- Molecule 2 is a protein called MAIT T-cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	200	Total	C	N	O	S	0	0	0
			1555	981	250	315	9			
2	D	198	Total	C	N	O	S	0	0	0
			1529	967	246	307	9			


- Molecule 3 is a protein called MAIT T-cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	243	Total	C	N	O	S	0	0	0
			1906	1197	333	367	9			
3	E	243	Total	C	N	O	S	0	0	0
			1915	1205	334	367	9			

- Molecule 4 is a protein called Beta-2-microglobulin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	97	Total	C	N	O	S	0	0	0
			792	506	134	150	2			
4	F	97	Total	C	N	O	S	0	0	0
			800	511	135	152	2			

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O		0	0
			6	3	3			

- Molecule 7 is 5-hydroxydiclofenac (three-letter code: 7ZV) (formula: $C_{14}H_{11}Cl_2NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total	C	Cl	N	O	0	0
			20	14	2	1	3		

- Molecule 8 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	C	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	60	Total	O	0	0
			60	60		
9	B	74	Total	O	0	0
			74	74		
9	G	85	Total	O	0	0
			85	85		
9	H	9	Total	O	0	0
			9	9		
9	C	68	Total	O	0	0
			68	68		
9	D	55	Total	O	0	0
			55	55		
9	E	59	Total	O	0	0
			59	59		
9	F	27	Total	O	0	0
			27	27		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 1: Major histocompatibility complex class I-related gene protein




- Molecule 2: MAIT T-cell receptor alpha chain



- Molecule 2: MAIT T-cell receptor alpha chain




- Molecule 3: MAIT T-cell receptor beta chain









- Molecule 3: MAIT T-cell receptor beta chain



- Molecule 4: Beta-2-microglobulin



- Molecule 4: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	213.09Å 69.60Å 142.37Å 90.00° 103.72° 90.00°	Depositor
Resolution (Å)	36.81 – 2.50 36.81 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.7 (36.81-2.50) 98.7 (36.81-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.51Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.173 , 0.231 0.173 , 0.231	Depositor DCC
R_{free} test set	3452 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	38.9	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13241	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7ZV, GOL, DMS, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2156	0.48	0/2933
1	C	0.33	0/2235	0.51	1/3036 (0.0%)
2	B	0.34	0/1590	0.53	0/2157
2	D	0.35	0/1563	0.52	0/2122
3	E	0.31	0/1966	0.51	0/2674
3	G	0.35	0/1955	0.50	0/2658
4	F	0.32	0/823	0.49	0/1118
4	H	0.28	0/815	0.46	0/1109
All	All	0.33	0/13103	0.50	1/17807 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	246	LEU	CA-CB-CG	7.02	131.44	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2094	0	1941	21	0
1	C	2171	0	2047	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1555	0	1463	15	0
2	D	1529	0	1426	11	0
3	E	1915	0	1815	13	0
3	G	1906	0	1807	18	0
4	F	800	0	753	6	0
4	H	792	0	738	7	0
5	A	8	0	6	2	0
5	F	4	0	3	0	0
6	A	6	0	8	0	0
7	C	20	0	0	1	0
8	C	4	0	6	0	0
9	A	60	0	0	0	0
9	B	74	0	0	3	0
9	C	68	0	0	2	0
9	D	55	0	0	0	0
9	E	59	0	0	2	0
9	F	27	0	0	0	0
9	G	85	0	0	6	0
9	H	9	0	0	0	0
All	All	13241	0	12013	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ARG:HD3	3:G:54:THR:HG21	1.37	1.03
3:G:64:TYR:O	9:G:301:HOH:O	2.00	0.79
1:C:246:LEU:O	9:C:401:HOH:O	2.02	0.77
1:C:211[B]:TYR:HB3	1:C:259:GLU:HB2	1.77	0.66
3:E:243:ARG:NH2	9:E:301:HOH:O	2.28	0.66
1:A:151:LEU:HD22	2:B:51:LEU:HD12	1.77	0.65
1:C:247:ASP:OD1	1:C:247:ASP:N	2.30	0.64
4:H:39:LEU:HD12	4:H:68:THR:HG23	1.78	0.64
3:G:2:ALA:N	9:G:304:HOH:O	2.33	0.60
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.84	0.59
4:F:49:VAL:HA	4:F:68:THR:HG22	1.83	0.59
1:C:211[A]:TYR:HB2	1:C:259:GLU:HB2	1.84	0.59
1:A:46:ARG:NH1	4:H:53:ASP:OD2	2.35	0.59
2:D:28:GLY:HA3	2:D:93:SER:OG	2.04	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ALA:HB2	4:H:60:TRP:CE2	2.39	0.57
1:C:77:LEU:HD13	1:C:92:TYR:HB2	1.87	0.55
4:F:39:LEU:HD12	4:F:68:THR:HG23	1.90	0.54
2:B:28:GLY:HA3	2:B:93:SER:OG	2.08	0.54
1:A:43:LYS:HD2	1:A:62:TYR:HB3	1.91	0.53
2:B:108:LYS:NZ	2:B:140:GLN:OE1	2.42	0.53
3:E:194:ARG:N	3:E:194:ARG:HD2	2.24	0.53
1:A:2:THR:HG22	5:A:302:ACT:H3	1.91	0.52
2:B:162:ARG:NH1	2:B:165:ASP:OD1	2.43	0.52
2:B:1:GLY:HA2	2:B:27:SER:H	1.75	0.52
1:A:229:ASP:OD2	4:H:6:LYS:NZ	2.26	0.51
3:E:117:ASP:OD1	3:E:119:LYS:HG2	2.11	0.51
1:A:152:TYR:CE1	3:G:100:GLY:HA3	2.45	0.51
4:H:48:LYS:O	4:H:68:THR:HG22	2.10	0.51
2:B:177:LYS:NZ	9:B:301:HOH:O	2.22	0.51
1:A:252:ASN:HB2	1:A:254:TYR:CE2	2.46	0.50
3:G:206:ARG:HD2	9:G:351:HOH:O	2.12	0.49
1:C:149:GLU:OE2	3:E:100:GLY:HA2	2.13	0.49
3:E:155:HIS:HB3	3:E:216:TYR:HB2	1.95	0.49
1:A:154:LYS:HD3	2:B:51:LEU:HD11	1.96	0.48
2:D:26:THR:HG23	2:D:28:GLY:O	2.14	0.48
3:E:119:LYS:HE2	3:E:226:GLN:OE1	2.13	0.47
1:C:155:ASN:OD1	2:D:66:ARG:NH1	2.46	0.47
2:D:36:GLN:HB2	2:D:46:LEU:HD11	1.95	0.47
3:E:8:PRO:HD2	3:E:21:LEU:HD22	1.96	0.47
3:G:174:ASP:OD1	3:G:194:ARG:NH2	2.47	0.47
1:A:15:PRO:HG3	1:A:21:GLU:HB3	1.96	0.46
2:B:25:GLN:NE2	9:B:307:HOH:O	2.48	0.46
1:A:122:PHE:HB2	1:A:129:TRP:CH2	2.49	0.46
3:E:131:PRO:HD2	3:E:202:TRP:CZ2	2.51	0.46
3:G:8:PRO:HD2	3:G:21:LEU:HD22	1.98	0.46
1:C:104:GLY:O	9:C:402:HOH:O	2.21	0.46
3:G:117:ASP:OD2	3:G:119:LYS:HG2	2.17	0.45
3:G:128:VAL:HG23	3:G:238:ALA:HB3	1.98	0.45
2:D:121:LEU:HB3	3:E:129:PHE:HB3	1.99	0.45
1:A:214:TRP:CD1	1:A:244:ILE:HG23	2.53	0.44
4:F:14:PRO:HA	4:F:97:ARG:HH22	1.82	0.44
2:B:1:GLY:CA	2:B:27:SER:H	2.30	0.44
1:C:113:ALA:HB2	4:F:60:TRP:CE2	2.53	0.44
1:A:8:PHE:CE1	1:A:95:MET:HG3	2.52	0.43
3:E:181:GLN:O	3:E:187:SER:HB2	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:210:ARG:NE	9:G:302:HOH:O	2.26	0.43
2:B:162:ARG:HD2	2:B:162:ARG:HA	1.86	0.43
1:A:19:VAL:HA	1:A:20:PRO:HD3	1.87	0.43
1:A:164:TRP:CE2	5:A:301:ACT:H2	2.54	0.43
1:C:33:ILE:HB	1:C:50:MET:HG3	2.01	0.43
2:D:1:GLY:CA	2:D:27:SER:H	2.32	0.43
1:C:196:THR:HB	1:C:246:LEU:HD13	2.01	0.43
2:D:65:SER:OG	2:D:68:LYS:HB2	2.19	0.43
2:B:125:LYS:O	1:C:130:LEU:HD23	2.19	0.42
1:C:155:ASN:HB2	2:D:51:LEU:CD1	2.49	0.42
1:A:28:VAL:HG23	1:A:33:ILE:HD13	2.02	0.42
1:C:5:LEU:HD11	1:C:7:TYR:HE2	1.83	0.42
3:E:30:ASN:O	3:E:51:SER:HA	2.19	0.42
2:D:54:LEU:HD11	2:D:61:SER:HB3	2.02	0.42
1:C:231:LEU:HD13	4:F:10:TYR:CE1	2.55	0.42
4:F:97:ARG:HD2	4:F:97:ARG:HA	1.26	0.42
2:B:170:SER:OG	3:G:194:ARG:HD2	2.19	0.42
2:B:68:LYS:NZ	9:B:302:HOH:O	2.27	0.42
1:A:152:TYR:CD1	3:G:100:GLY:HA3	2.54	0.42
1:C:46:ARG:HA	1:C:46:ARG:HD3	1.80	0.42
3:E:188:ARG:NH2	9:E:302:HOH:O	2.35	0.42
1:A:40:THR:O	1:A:42:GLN:HG2	2.20	0.41
3:E:20:THR:OG1	3:E:77:ARG:HD3	2.20	0.41
2:B:166:PHE:CD2	3:G:141:LYS:HE2	2.55	0.41
2:B:199:SER:HA	2:B:200:PRO:HD3	1.81	0.41
1:C:34:THR:HB	1:C:43:LYS:HE2	2.01	0.41
2:D:199:SER:HA	2:D:200:PRO:HD3	1.88	0.41
3:G:19:MET:HB3	3:G:78:LEU:HB2	2.02	0.41
3:G:203:GLN:HA	3:G:243:ARG:O	2.20	0.41
4:H:37:VAL:HB	4:H:66:TYR:CZ	2.56	0.41
3:G:111:ARG:NE	9:G:316:HOH:O	2.54	0.41
3:G:80:SER:N	9:G:313:HOH:O	2.52	0.41
1:C:66:LEU:HD21	7:C:301:7ZV:C	2.51	0.40
1:A:6:ARG:NH1	4:H:58:LYS:HD3	2.36	0.40
2:D:23:THR:HG22	2:D:70:TYR:HB2	2.03	0.40
3:G:86:THR:HG23	3:G:113:THR:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/271 (93%)	248 (99%)	3 (1%)	0	100	100
1	C	256/271 (94%)	251 (98%)	5 (2%)	0	100	100
2	B	198/203 (98%)	193 (98%)	5 (2%)	0	100	100
2	D	194/203 (96%)	187 (96%)	7 (4%)	0	100	100
3	E	241/245 (98%)	233 (97%)	8 (3%)	0	100	100
3	G	241/245 (98%)	232 (96%)	9 (4%)	0	100	100
4	F	95/99 (96%)	93 (98%)	2 (2%)	0	100	100
4	H	95/99 (96%)	94 (99%)	1 (1%)	0	100	100
All	All	1571/1636 (96%)	1531 (98%)	40 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/241 (90%)	217 (100%)	1 (0%)	92	98
1	C	230/241 (95%)	223 (97%)	7 (3%)	48	76
2	B	173/180 (96%)	173 (100%)	0	100	100
2	D	167/180 (93%)	166 (99%)	1 (1%)	90	97
3	E	207/211 (98%)	206 (100%)	1 (0%)	92	98
3	G	206/211 (98%)	204 (99%)	2 (1%)	82	95

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	F	89/94 (95%)	86 (97%)	3 (3%)	44	72
4	H	87/94 (93%)	87 (100%)	0	100	100
All	All	1377/1452 (95%)	1362 (99%)	15 (1%)	80	94

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	PHE
3	G	117	ASP
3	G	194	ARG
1	C	73	PHE
1	C	76	GLU
1	C	221	ILE
1	C	245	GLU
1	C	246	LEU
1	C	247	ASP
1	C	268	GLN
2	D	91	LYS
3	E	194	ARG
4	F	20	SER
4	F	70	PHE
4	F	97	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	A	301	-	0,3,3	0.00	-	0,3,3	0.00	-
5	ACT	A	302	-	0,3,3	0.00	-	0,3,3	0.00	-
6	GOL	A	303	-	5,5,5	0.32	0	5,5,5	0.43	0
7	7ZV	C	301	-	18,21,21	0.34	0	26,29,29	0.40	0
8	DMS	C	302	-	3,3,3	0.62	0	3,3,3	0.36	0
5	ACT	F	101	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ACT	A	301	-	-	0/0/0/0	0/0/0/0
5	ACT	A	302	-	-	0/0/0/0	0/0/0/0
6	GOL	A	303	-	-	0/4/4/4	0/0/0/0
7	7ZV	C	301	-	-	0/6/8/8	0/2/2/2
8	DMS	C	302	-	-	0/0/0/0	0/0/0/0
5	ACT	F	101	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	301	ACT	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	302	ACT	1	0
7	C	301	7ZV	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	259/271 (95%)	0.14	17 (6%)	22 24	26, 43, 74, 108	0
1	C	263/271 (97%)	-0.01	9 (3%)	49 54	25, 37, 66, 106	1 (0%)
2	B	200/203 (98%)	-0.07	3 (1%)	76 79	23, 36, 63, 93	0
2	D	198/203 (97%)	0.15	13 (6%)	22 24	26, 45, 88, 127	0
3	E	243/245 (99%)	0.05	9 (3%)	45 50	26, 46, 85, 105	1 (0%)
3	G	243/245 (99%)	-0.12	2 (0%)	87 89	26, 39, 63, 103	1 (0%)
4	F	97/99 (97%)	0.18	3 (3%)	52 57	26, 44, 69, 92	0
4	H	97/99 (97%)	0.81	17 (17%)	2 2	38, 67, 97, 110	0
All	All	1600/1636 (97%)	0.08	73 (4%)	36 41	23, 42, 81, 127	3 (0%)

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	0	MET	6.0
1	C	251	SER	5.1
1	A	246	LEU	4.6
3	E	225	THR	4.2
2	B	179	ASP	4.0
2	D	149	SER	3.8
3	E	2	ALA	3.7
3	E	221	ASN	3.6
4	H	47	GLU	3.5
1	A	254	TYR	3.4
1	A	244	ILE	3.4
4	H	49	VAL	3.4
1	A	245	GLU	3.3
4	H	42	ASN	3.3
2	D	179	ASP	3.3
2	D	162	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	195	VAL	3.3
4	H	44	GLU	3.2
1	C	252	ASN	3.2
1	C	221	ILE	3.2
1	A	198	LEU	3.1
4	H	43	GLY	3.1
1	A	250	SER	3.1
4	H	78	TYR	3.0
3	E	224	TRP	3.0
1	A	199	PHE	2.9
1	A	248	PRO	2.9
3	E	206	ARG	2.9
4	H	77	GLU	2.8
2	D	128	ASP	2.8
1	C	194	GLY	2.8
2	B	1	GLY	2.8
4	H	40	LEU	2.8
3	G	185	ASN	2.8
1	A	225	ILE	2.8
1	A	243	SER	2.7
3	E	229	ALA	2.7
2	D	129	LYS	2.7
1	A	247	ASP	2.7
1	A	197	ALA	2.6
4	H	73	THR	2.6
1	C	191	THR	2.6
1	A	0	MET	2.6
4	H	87	LEU	2.5
1	A	224	GLU	2.5
4	H	68	THR	2.5
2	B	178	SER	2.5
4	H	48	LYS	2.4
4	H	94	LYS	2.4
1	A	103	ASP	2.4
2	D	165	ASP	2.4
1	A	223	GLN	2.4
4	H	45	ARG	2.4
3	E	228	ARG	2.4
4	H	46	ILE	2.4
2	D	178	SER	2.4
2	D	200	PRO	2.4
2	D	112	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	220	GLU	2.3
2	D	1	GLY	2.3
4	H	41	LYS	2.2
2	D	180	PHE	2.2
4	F	47	GLU	2.2
3	G	184	LEU	2.2
2	D	163	SER	2.2
4	H	79	ALA	2.2
2	D	130	SER	2.2
4	F	68	THR	2.2
1	C	193	PRO	2.1
3	E	223	GLU	2.1
1	C	246	LEU	2.1
4	F	44	GLU	2.0
3	E	227	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	ACT	F	101	4/4	0.95	0.22	5.64	48,48,53,58	0
8	DMS	C	302	4/4	0.93	0.25	5.42	57,69,75,76	0
6	GOL	A	303	6/6	0.90	0.25	4.31	48,52,55,58	0
7	7ZV	C	301	20/20	0.94	0.18	0.35	40,58,66,76	0
5	ACT	A	301	4/4	0.95	0.15	0.30	38,41,45,46	0
5	ACT	A	302	4/4	0.95	0.14	-0.37	53,55,57,58	0

6.5 Other polymers

There are no such residues in this entry.