



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 2, 2017 – 05:17 PM EST

PDB ID : 5UAL  
Title : Escherichia coli RNA polymerase and Rifampin complex, RpoB S531L mutant  
Authors : Molodtsov, V.; Scharf, N.T.; Stefan, M.A.; Garcia, G.A.; Murakami, K.S.  
Deposited on : 2016-12-19  
Resolution : 3.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1005 (4.24-3.52)
Clashscore	102246	1026 (4.20-3.56)
Ramachandran outliers	100387	1003 (4.22-3.54)
Sidechain outliers	100360	1043 (4.24-3.52)
RSRZ outliers	91569	1009 (4.24-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div><div>0.0%</div><div><div>44%</div><div>22%</div><div>•</div><div>31%</div></div></div>
1	B	329	<div><div>2%</div><div><div>39%</div><div>24%</div><div>•</div><div>35%</div></div></div>
1	G	329	<div><div>0.0%</div><div><div>41%</div><div>23%</div><div>•</div><div>32%</div></div></div>
1	H	329	<div><div>2%</div><div><div>37%</div><div>26%</div><div>•</div><div>35%</div></div></div>
2	C	1342	<div><div>5%</div><div><div>62%</div><div>34%</div><div>••</div></div></div>
2	I	1342	<div><div>5%</div><div><div>67%</div><div>29%</div><div>••</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	1407	<div><div><div></div><div></div><div></div><div></div></div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>52%27%•17%</div></div>
3	J	1407	<div><div><div></div><div></div><div></div><div></div></div><div>3%</div><div><div></div><div></div><div></div><div></div></div><div>53%25%•18%</div></div>
4	E	91	<div><div><div></div><div></div><div></div><div></div></div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>78%18%••</div></div>
4	K	91	<div><div><div></div><div></div><div></div><div></div></div><div>21%</div><div><div></div><div></div><div></div><div></div></div><div>56%27%•13%</div></div>
5	F	613	<div><div><div></div><div></div><div></div><div></div></div><div>3%</div><div><div></div><div></div><div></div><div></div></div><div>50%23%•24%</div></div>
5	L	613	<div><div><div></div><div></div><div></div><div></div></div><div>3%</div><div><div></div><div></div><div></div><div></div></div><div>53%21%•23%</div></div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 54898 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1753	1091	311	345	6			
1	B	214	Total	C	N	O	S	0	0	0
			1649	1029	290	324	6			
1	G	224	Total	C	N	O	S	0	0	0
			1730	1076	308	340	6			
1	H	215	Total	C	N	O	S	0	0	0
			1659	1037	291	325	6			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1332	Total	C	N	O	S	0	0	0
			10514	6600	1827	2043	44			
2	I	1328	Total	C	N	O	S	0	0	0
			10486	6583	1822	2038	43			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	531	LEU	SER	engineered mutation	UNP P0A8V2
I	531	LEU	SER	engineered mutation	UNP P0A8V2

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1166	Total	C	N	O	S	0	0	0
			9089	5714	1627	1702	46			
3	J	1155	Total	C	N	O	S	0	0	0
			9001	5659	1612	1684	46			


- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	89	Total	C	N	O	S	0	0	0
			691	421	129	140	1			
4	K	79	Total	C	N	O	S	0	0	0
			627	382	118	126	1			

- Molecule 5 is a protein called RNA polymerase sigma factor RpoD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	468	Total	C	N	O	S	0	0	0
			3813	2389	678	723	23			
5	L	469	Total	C	N	O	S	0	0	0
			3821	2393	679	726	23			

- Molecule 6 is RIFAMPICIN (three-letter code: RFP) (formula:  $C_{43}H_{58}N_4O_{12}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	N	O	0	0
			59	43	4	12		


- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	J	1	Total	Mg	0	0
			1	1		
7	D	1	Total	Mg	0	0
			1	1		


- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	J	2	Total 2	Zn 2	0	0
8	D	2	Total 2	Zn 2	0	0







• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 2: DNA-directed RNA polymerase subunit beta









• Molecule 2: DNA-directed RNA polymerase subunit beta






• Molecule 3: DNA-directed RNA polymerase subunit beta'












• Molecule 5: RNA polymerase sigma factor RpoD



• Molecule 5: RNA polymerase sigma factor RpoD





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	187.37Å 205.95Å 309.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.94 – 3.89 29.94 – 3.89	Depositor EDS
% Data completeness (in resolution range)	82.9 (29.94-3.89) 82.9 (29.94-3.89)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 3.86Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, $R_{free}$	0.230 , 0.286 0.225 , 0.284	Depositor DCC
$R_{free}$ test set	1975 reflections (2.16%)	DCC
Wilson B-factor (Å <sup>2</sup> )	168.3	Xtriage
Anisotropy	0.221	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 134.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	54898	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	210.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, RFP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.42	1/1774 (0.1%)	0.57	0/2405
1	B	0.34	0/1668	0.71	2/2260 (0.1%)
1	G	0.36	1/1751 (0.1%)	0.55	1/2373 (0.0%)
1	H	0.29	0/1678	0.62	0/2274
2	C	0.37	0/10681	0.57	0/14410
2	I	0.30	0/10653	0.53	0/14373
3	D	0.39	1/9229 (0.0%)	0.61	0/12459
3	J	0.34	0/9140	0.58	2/12341 (0.0%)
4	E	0.29	0/693	0.51	0/935
4	K	0.26	0/629	0.47	0/847
5	F	0.30	0/3864	0.53	0/5194
5	L	0.28	0/3872	0.50	0/5205
All	All	0.34	3/55632 (0.0%)	0.57	5/75076 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	2
2	I	0	2
3	D	0	2
3	J	0	2
5	F	0	1
5	L	0	1
All	All	0	10

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	29	GLU	C-N	10.64	1.54	1.34
1	G	29	GLU	C-N	9.19	1.51	1.34
3	D	898	CYS	CB-SG	-5.28	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	343	LEU	CA-CB-CG	12.07	143.06	115.30
1	B	9	LEU	C-N-CA	9.90	146.46	121.70
1	G	9	LEU	C-N-CA	5.58	135.64	121.70
3	J	857	LEU	CA-CB-CG	5.13	127.09	115.30
1	B	13	LEU	CA-CB-CG	5.13	127.09	115.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	109	ALA	Peptide
2	C	236	LYS	Peptide
3	D	1184	ASP	Peptide
3	D	1296	GLY	Peptide
5	F	601	PRO	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1753	0	1780	64	0
1	B	1649	0	1673	71	0
1	G	1730	0	1756	66	0
1	H	1659	0	1692	72	0
2	C	10514	0	10531	335	0
2	I	10486	0	10496	268	0
3	D	9089	0	9263	310	0
3	J	9001	0	9167	288	0
4	E	691	0	695	11	0
4	K	627	0	634	22	0
5	F	3813	0	3880	96	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	3821	0	3884	82	0
6	C	59	0	56	9	0
7	D	1	0	0	0	0
7	J	1	0	0	0	0
8	D	2	0	0	0	0
8	J	2	0	0	0	0
All	All	54898	0	55507	1505	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 1505 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:343:LEU:HD22	3:D:344:GLY:HA3	1.36	1.07
2:C:1271:GLY:HA2	3:D:343:LEU:HD21	1.44	0.99
2:C:1269:ARG:HG3	3:D:343:LEU:HD12	1.49	0.94
2:C:65:ASN:HB3	2:C:105:TYR:HB2	1.50	0.91
2:C:745:GLU:HG3	2:C:1017:GLN:HB3	1.52	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	225/329 (68%)	195 (87%)	26 (12%)	4 (2%)	11 54
1	B	210/329 (64%)	180 (86%)	25 (12%)	5 (2%)	7 49
1	G	222/329 (68%)	192 (86%)	24 (11%)	6 (3%)	6 46
1	H	211/329 (64%)	185 (88%)	21 (10%)	5 (2%)	7 49
2	C	1328/1342 (99%)	1220 (92%)	101 (8%)	7 (0%)	34 76

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	1324/1342 (99%)	1220 (92%)	97 (7%)	7 (0%)	34	76
3	D	1162/1407 (83%)	1070 (92%)	85 (7%)	7 (1%)	30	73
3	J	1151/1407 (82%)	1059 (92%)	80 (7%)	12 (1%)	19	64
4	E	87/91 (96%)	82 (94%)	5 (6%)	0	100	100
4	K	77/91 (85%)	75 (97%)	2 (3%)	0	100	100
5	F	462/613 (75%)	424 (92%)	37 (8%)	1 (0%)	52	86
5	L	463/613 (76%)	426 (92%)	36 (8%)	1 (0%)	52	86
All	All	6922/8222 (84%)	6328 (91%)	539 (8%)	55 (1%)	24	68

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	GLU
1	B	13	LEU
2	C	3	TYR
2	C	1159	VAL
3	D	10	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/286 (68%)	181 (93%)	13 (7%)	20	59
1	B	182/286 (64%)	172 (94%)	10 (6%)	27	66
1	G	191/286 (67%)	179 (94%)	12 (6%)	22	61
1	H	184/286 (64%)	175 (95%)	9 (5%)	31	68
2	C	1150/1157 (99%)	1050 (91%)	100 (9%)	13	48
2	I	1147/1157 (99%)	1045 (91%)	102 (9%)	12	47
3	D	970/1168 (83%)	871 (90%)	99 (10%)	9	40
3	J	960/1168 (82%)	864 (90%)	96 (10%)	9	41
4	E	72/75 (96%)	65 (90%)	7 (10%)	10	43

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	K	67/75 (89%)	63 (94%)	4 (6%)	24	63
5	F	417/540 (77%)	376 (90%)	41 (10%)	10	42
5	L	418/540 (77%)	378 (90%)	40 (10%)	10	43
All	All	5952/7024 (85%)	5419 (91%)	533 (9%)	12	47

5 of 533 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	F	417	ASP
2	I	90	VAL
4	K	58	LEU
5	F	471	LEU
1	G	9	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 64 such sidechains are listed below:

Mol	Chain	Res	Type
5	F	309	ASN
2	I	139	ASN
5	L	131	GLN
5	F	383	ASN
5	F	518	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 6 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	RFP	C	3001	-	62,63,63	2.47	13 (20%)	76,94,94	2.02	22 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	RFP	C	3001	-	-	0/60/85/85	0/1/5/5

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	3001	RFP	C12-C11	-6.54	1.36	1.54
6	C	3001	RFP	O7-C25	-4.93	1.37	1.44
6	C	3001	RFP	O6-C27	-2.87	1.37	1.43
6	C	3001	RFP	C6-C7	-2.41	1.35	1.39
6	C	3001	RFP	C26-C25	-2.16	1.49	1.53

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	3001	RFP	C24-C23-C22	-4.78	108.47	115.46
6	C	3001	RFP	O4-C11-C5	-4.61	122.23	131.67
6	C	3001	RFP	C12-O3-C6	-3.90	103.36	107.86
6	C	3001	RFP	C20-C19-C18	-3.37	118.62	126.06
6	C	3001	RFP	C18-C17-C16	-3.14	118.11	127.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	3001	RFP	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	227/329 (68%)	-0.30	3 (1%) 79 70	134, 178, 241, 363	0
1	B	214/329 (65%)	0.05	8 (3%) 45 34	141, 222, 328, 396	0
1	G	224/329 (68%)	-0.10	4 (1%) 71 61	175, 225, 285, 364	0
1	H	215/329 (65%)	0.20	8 (3%) 45 34	179, 247, 312, 401	0
2	C	1332/1342 (99%)	-0.10	64 (4%) 34 25	101, 169, 396, 542	0
2	I	1328/1342 (98%)	0.06	62 (4%) 35 26	139, 219, 353, 523	0
3	D	1166/1407 (82%)	-0.25	11 (0%) 85 79	93, 151, 275, 441	0
3	J	1155/1407 (82%)	-0.09	40 (3%) 48 36	128, 181, 311, 455	0
4	E	89/91 (97%)	-0.14	1 (1%) 82 74	144, 205, 237, 275	0
4	K	79/91 (86%)	0.92	19 (24%) 1 1	255, 318, 411, 416	0
5	F	468/613 (76%)	-0.15	17 (3%) 46 35	139, 222, 362, 481	0
5	L	469/613 (76%)	-0.15	16 (3%) 49 37	161, 231, 357, 487	0
All	All	6966/8222 (84%)	-0.08	253 (3%) 46 35	93, 196, 346, 542	0

The worst 5 of 253 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	311	CYS	14.1
2	C	252	SER	11.6
2	I	981	ALA	8.1
2	I	982	GLY	7.9
2	C	283	LYS	7.6

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	RFP	C	3001	59/59	0.95	0.21	-0.19	91,139,195,202	0
8	ZN	J	1503	1/1	0.98	0.14	-0.24	126,126,126,126	0
8	ZN	D	1502	1/1	0.97	0.07	-1.13	150,150,150,150	0
8	ZN	D	1503	1/1	0.99	0.12	-1.41	82,82,82,82	0
8	ZN	J	1502	1/1	0.96	0.05	-1.42	174,174,174,174	0
7	MG	J	1501	1/1	0.95	0.42	-	121,121,121,121	0
7	MG	D	1501	1/1	0.94	0.39	-	77,77,77,77	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.