



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 05:13 PM BST

PDB ID : 1UFG
Title : Solution structure of immunoglobulin like domain of mouse nuclear lamin
Authors : Kobayashi, N.; Kigawa, T.; Koshiba, S.; Inoue, M.; Yokoyama, S.; RIKEN
Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2003-05-29

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	151	

2 Ensemble composition and analysis

This entry contains 20 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:33-A:73, A:78-A:145 (109)	0.20	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 9, 12, 19
2	7, 8, 11, 16
3	6, 15, 18, 20
4	13, 14
Single-model clusters	10; 17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2256 atoms, of which 1115 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Lamin A.

Mol	Chain	Residues	Atoms						Trace
1	A	151	Total	C	H	N	O	S	0
			2256	692	1115	218	227	4	

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP P48678
A	2	SER	-	CLONING ARTIFACT	UNP P48678
A	3	SER	-	CLONING ARTIFACT	UNP P48678
A	4	GLY	-	CLONING ARTIFACT	UNP P48678
A	5	SER	-	CLONING ARTIFACT	UNP P48678
A	6	SER	-	CLONING ARTIFACT	UNP P48678
A	7	GLY	-	CLONING ARTIFACT	UNP P48678
A	146	SER	-	CLONING ARTIFACT	UNP P48678
A	147	GLY	-	CLONING ARTIFACT	UNP P48678
A	148	PRO	-	CLONING ARTIFACT	UNP P48678
A	149	SER	-	CLONING ARTIFACT	UNP P48678
A	150	SER	-	CLONING ARTIFACT	UNP P48678
A	151	GLY	-	CLONING ARTIFACT	UNP P48678

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Lamin A



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: Lamin A



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.7
CYANA	refinement	1.0.7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 10120
Number of chemical shift lists	1
Total number of shifts	1648
Number of shifts mapped to atoms	1648
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	858	854	854	49±7
All	All	17160	17080	17080	978

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

5 of 276 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:69:ILE:HD11	1:A:83:PHE:CZ	1.03	1.88	4	19
1:A:69:ILE:HD11	1:A:83:PHE:CE2	0.97	1.93	3	20
1:A:33:HIS:NE2	1:A:144:VAL:HG23	0.87	1.85	8	2
1:A:45:VAL:HG22	1:A:52:VAL:HG13	0.83	1.48	17	19
1:A:125:SER:HA	1:A:144:VAL:HG23	0.81	1.53	19	4

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/151 (72%)	87±2 (80±2%)	17±2 (16±2%)	5±1 (4±1%)	6	30
All	All	2180/3020 (72%)	1739 (80%)	345 (16%)	96 (4%)	6	30

5 of 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	110	THR	19
1	A	90	LYS	18
1	A	66	ASN	17
1	A	67	TRP	14
1	A	106	HIS	8

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/124 (73%)	72±3 (79±4%)	20±3 (21±4%)	4	33
All	All	1820/2480 (73%)	1430 (79%)	390 (21%)	4	33

5 of 49 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	36	THR	20
1	A	53	ARG	20
1	A	126	LEU	20
1	A	107	SER	20
1	A	66	ASN	19

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 83% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 10120

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1648
Number of shifts mapped to atoms	1648
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	23

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	141	-0.23 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	125	-0.10 ± 0.06	None needed (< 0.5 ppm)
$^{13}\text{C}'$	120	0.25 ± 0.07	None needed (< 0.5 ppm)
^{15}N	133	-1.39 ± 0.40	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 1223 atoms were assigned a chemical shift out of a possible 1360. 1 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	525/537 (98%)	214/214 (100%)	206/218 (94%)	105/105 (100%)
Sidechain	607/726 (84%)	375/427 (88%)	212/252 (84%)	20/47 (43%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	91/97 (94%)	47/51 (92%)	40/40 (100%)	4/6 (67%)
Overall	1223/1360 (90%)	636/692 (92%)	458/510 (90%)	129/158 (82%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	35	ARG	NE	112.49	92.63 – 76.73	17.5
1	A	53	ARG	NE	110.56	92.63 – 76.73	16.3
1	A	145	ARG	NE	109.96	92.63 – 76.73	15.9
1	A	70	ARG	NE	109.92	92.63 – 76.73	15.9
1	A	71	ARG	NE	65.38	92.63 – 76.73	-12.1
1	A	117	GLN	HB2	-0.12	3.30 – 0.80	-8.7
1	A	46	ASP	HB2	0.63	4.07 – 1.37	-7.7
1	A	117	GLN	HG2	0.48	3.67 – 0.97	-6.8
1	A	96	THR	HG21	-0.27	2.29 – -0.01	-6.1
1	A	96	THR	HG22	-0.27	2.29 – -0.01	-6.1
1	A	96	THR	HG23	-0.27	2.29 – -0.01	-6.1
1	A	81	TYR	HE2	5.32	7.86 – 5.56	-6.0
1	A	81	TYR	HE1	5.32	7.86 – 5.56	-6.0
1	A	64	MET	HG2	0.37	4.23 – 0.63	-5.7
1	A	52	VAL	HG11	-0.61	2.13 – -0.47	-5.6
1	A	52	VAL	HG12	-0.61	2.13 – -0.47	-5.6
1	A	52	VAL	HG13	-0.61	2.13 – -0.47	-5.6
1	A	81	TYR	HD2	5.28	8.44 – 5.44	-5.5
1	A	81	TYR	HD1	5.28	8.44 – 5.44	-5.5
1	A	18	LYS	CB	42.34	41.68 – 23.88	5.4
1	A	132	ASN	HA	2.73	6.52 – 2.82	-5.2
1	A	81	TYR	HB2	0.98	4.76 – 1.06	-5.2
1	A	62	GLN	NE2	121.10	120.91 – 102.81	5.1

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

