



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:38 PM GMT

PDB ID : 3UFS
Title : Structure of rat nitric oxide synthase heme domain in complex with 6-(((3R,4R)-4-((5-(3-chloro-5-fluorophenyl)pentyl)oxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2011-11-01
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	8664 (2.00-1.96)
Clashscore	102246	9905 (2.00-1.96)
Ramachandran outliers	100387	9792 (2.00-1.96)
Sidechain outliers	100360	9791 (2.00-1.96)
RSRZ outliers	91569	8679 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>24%</div> <div>83%</div> <div>12%</div> <div>..</div> </div>
1	B	422	<div> <div>10%</div> <div>86%</div> <div>10%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	860	-	-	-	X
5	ACT	B	860	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7141 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	2	0
			3324	2128	568	607	21			
1	B	411	Total	C	N	O	S	0	1	0
			3348	2142	574	610	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 6-[[[(3R,4R)-4-{[5-(3-CHLORO-5-FLUOROPHENYL)PENTYL]OXY}PYRROLIDIN-3-YL]METHYL]-4-METHYLPYRIDIN-2-AMINE (three-letter code: HW6) (formula: C₂₂H₂₉ClF₂N₃O).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	Cl	F	N	O	0	0
			28	22	1	1	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	Cl	F	N	O	0	0
			28	22	1	1	3	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	116	Total	O	0	0
			116	116		
7	B	168	Total	O	0	0
			168	168		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.83Å 111.28Å 164.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.64 – 1.97 37.64 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.0 (37.64-1.97) 99.0 (37.64-1.97)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.193 , 0.233 0.220 , 0.254	Depositor DCC
R_{free} test set	3325 reflections (5.19%)	DCC
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.297	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 67337 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7141	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HW6, HEM, ZN, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/3423	0.69	0/4644
1	B	0.77	0/3444	0.72	1/4669 (0.0%)
All	All	0.73	0/6867	0.70	1/9313 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	489	ASP	CB-CG-OD2	5.94	123.65	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3324	0	3235	40	0
1	B	3348	0	3264	28	0
2	A	43	0	30	1	0
2	B	43	0	30	1	0
3	A	17	0	15	2	0
3	B	17	0	15	0	0
4	A	28	0	29	2	0
4	B	28	0	29	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0
7	A	116	0	0	0	0
7	B	168	0	0	1	0
All	All	7141	0	6653	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:645:LYS:NZ	1:A:645:LYS:HB2	1.87	0.89
1:A:645:LYS:HB2	1:A:645:LYS:HZ1	1.48	0.76
1:A:380:ARG:HD3	1:A:400:GLU:OE1	1.95	0.66
1:A:317:HIS:O	1:A:320:SER:HB3	1.99	0.63
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.84	0.59
1:B:353:GLN:H	1:B:353:GLN:HE21	1.50	0.59
1:A:306:TRP:CE2	1:B:336:MET:HE3	2.39	0.58
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.87	0.57
1:A:371:ARG:HG3	1:A:371:ARG:HH21	1.71	0.56
1:A:299:ARG:HG3	1:A:318:LEU:HD21	1.88	0.55
1:A:371:ARG:CG	1:A:371:ARG:HH21	2.20	0.55
1:A:486:LYS:HD2	1:A:499:VAL:HG11	1.89	0.55
1:B:504:ILE:O	1:B:508:GLN:HG2	2.09	0.53
1:A:517:PHE:CB	1:A:560:LYS:HE3	2.40	0.52
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.91	0.52
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.46	0.51
1:A:517:PHE:HB2	1:A:560:LYS:HE3	1.93	0.51
1:A:336:MET:HE3	1:B:306:TRP:CD1	2.46	0.51
1:A:337:LEU:HD21	4:A:800:HW6:H25	1.93	0.51
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.00	0.50
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	1.94	0.50
1:A:609:GLU:O	1:A:613:LYS:HG2	2.11	0.50
1:A:336:MET:CE	3:A:760:H4B:H9	2.43	0.49
1:A:588:TYR:CD1	1:A:593:ILE:HD11	2.47	0.49
1:A:336:MET:HE2	3:A:760:H4B:H9	1.95	0.49
1:A:628:GLN:NE2	1:B:632:GLU:OE2	2.46	0.48
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.94	0.48
1:B:548:HIS:NE2	1:B:632:GLU:OE1	2.43	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:645:LYS:CB	1:A:645:LYS:NZ	2.63	0.47
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.50	0.47
1:A:371:ARG:CG	1:A:371:ARG:NH2	2.78	0.46
1:B:353:GLN:H	1:B:353:GLN:NE2	2.14	0.46
1:B:350:THR:HG22	1:B:352:ASP:HB3	1.97	0.45
1:B:445:HIS:C	1:B:445:HIS:CD2	2.90	0.44
1:B:475:TRP:HB2	1:B:523:LEU:HB3	1.99	0.44
1:A:460:THR:O	1:A:583:PRO:HD2	2.18	0.44
1:A:475:TRP:CZ2	1:A:531:PRO:HG3	2.53	0.44
2:A:750:HEM:CMC	2:A:750:HEM:HBC2	2.48	0.43
1:A:674:ALA:HB3	1:A:695:MET:HB3	2.01	0.43
1:B:486:LYS:HB2	1:B:486:LYS:HE2	1.71	0.43
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.54	0.43
1:A:507:GLN:O	1:A:507:GLN:HG2	2.18	0.43
1:B:675:ASP:O	1:B:679:ILE:HG12	2.18	0.43
1:B:478:GLN:HA	1:B:566:ALA:O	2.19	0.42
1:B:680:VAL:HA	1:B:681:PRO:HD3	1.91	0.42
1:A:492:THR:HG21	1:A:496:PRO:HG3	2.00	0.42
1:A:475:TRP:CE2	1:A:531:PRO:HG3	2.54	0.42
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.52	0.42
1:A:626:LYS:HB3	1:B:687:ILE:HD12	2.00	0.42
1:B:496:PRO:HB2	1:B:602:SER:O	2.19	0.42
1:B:391:THR:O	1:B:392:SER:CB	2.66	0.42
1:A:569:ASN:H	1:A:569:ASN:HD22	1.68	0.42
1:B:460:THR:O	1:B:583:PRO:HD2	2.19	0.42
1:A:455:LEU:HD12	1:A:587:TRP:HB3	2.02	0.42
1:A:330:ILE:HD11	1:B:696:LEU:HB3	2.02	0.42
1:A:485:TYR:HB3	1:A:514:ARG:NH1	2.35	0.41
1:A:378:MET:HA	1:A:378:MET:CE	2.50	0.41
1:B:507:GLN:NE2	7:B:129:HOH:O	2.53	0.41
1:A:587:TRP:O	4:A:800:HW6:H1	2.19	0.41
1:B:525:GLN:HG3	1:B:529:ASN:O	2.20	0.41
1:B:595:VAL:O	1:B:599:CYS:HB2	2.20	0.41
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.09	0.41
1:A:525:GLN:HG3	1:A:529:ASN:O	2.21	0.41
1:B:537:PRO:HA	1:B:538:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/422 (96%)	394 (97%)	11 (3%)	0	100	100
1	B	408/422 (97%)	400 (98%)	8 (2%)	0	100	100
All	All	813/844 (96%)	794 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/377 (97%)	347 (95%)	18 (5%)	31	22
1	B	367/377 (97%)	359 (98%)	8 (2%)	60	59
All	All	732/754 (97%)	706 (96%)	26 (4%)	44	36

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	319	LYS
1	A	320	SER
1	A	370	LYS
1	A	375	LYS
1	A	378	MET
1	A	380	ARG
1	A	390	SER
1	A	507	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	523	LEU
1	A	527	ASN
1	A	547	ARG
1	A	569	ASN
1	A	601[A]	ASN
1	A	601[B]	ASN
1	A	620	LYS
1	A	645	LYS
1	A	713	THR
1	A	715	VAL
1	B	336	MET
1	B	350	THR
1	B	353	GLN
1	B	392	SER
1	B	454	ASN
1	B	486	LYS
1	B	540	LEU
1	B	547	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	440	ASN
1	A	454	ASN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	628	GLN
1	A	697	ASN
1	B	353	GLN
1	B	385	ASN
1	B	425	GLN
1	B	454	ASN
1	B	507	GLN
1	B	601	ASN
1	B	605	ASN
1	B	642	GLN
1	B	697	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	750	1	30,50,50	2.31	9 (30%)	24,82,82	2.78	12 (50%)
3	H4B	A	760	-	13,18,18	0.83	0	11,26,26	2.74	6 (54%)
4	HW6	A	800	-	27,30,30	1.09	2 (7%)	31,40,40	2.39	11 (35%)
5	ACT	A	860	-	1,3,3	1.12	0	0,3,3	0.00	-
2	HEM	B	750	1	30,50,50	2.28	9 (30%)	24,82,82	2.61	12 (50%)
3	H4B	B	760	-	13,18,18	1.10	0	11,26,26	2.70	6 (54%)
4	HW6	B	800	-	27,30,30	1.04	2 (7%)	31,40,40	2.25	9 (29%)
5	ACT	B	860	-	1,3,3	1.57	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	HW6	A	800	-	-	0/13/23/23	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	HW6	B	800	-	-	0/13/23/23	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3D-C4D	-6.61	1.43	1.51
2	A	750	HEM	C3B-C4B	-6.12	1.46	1.51
2	B	750	HEM	C3D-C4D	-5.97	1.43	1.51
2	B	750	HEM	C3B-C4B	-5.89	1.46	1.51
2	B	750	HEM	C2C-C1C	-4.34	1.44	1.52
2	A	750	HEM	C2C-C1C	-3.43	1.46	1.52
2	B	750	HEM	C2B-C1B	-2.60	1.43	1.51
2	A	750	HEM	C2D-C1D	-2.21	1.44	1.51
2	A	750	HEM	C2B-C1B	-2.00	1.45	1.51
2	A	750	HEM	CMB-C2B	2.08	1.58	1.53
2	A	750	HEM	FE-NC	2.12	2.04	1.95
2	A	750	HEM	C3B-CAB	2.36	1.55	1.51
2	B	750	HEM	CAA-C2A	2.37	1.56	1.52
2	B	750	HEM	C1C-NC	2.38	1.38	1.36
2	B	750	HEM	FE-NC	2.46	2.05	1.95
4	B	800	HW6	C24-C23	2.53	1.42	1.37
4	A	800	HW6	C24-C23	2.56	1.42	1.37
2	B	750	HEM	C3C-CAC	2.70	1.56	1.51
2	B	750	HEM	CMA-C3A	3.19	1.58	1.51
4	A	800	HW6	C25-CL2	3.31	1.82	1.74
4	B	800	HW6	C25-CL2	3.32	1.82	1.74
2	A	750	HEM	C1C-NC	3.88	1.40	1.36

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CBA-CAA-C2A	-6.02	101.74	112.53
4	A	800	HW6	F23-C23-C22	-4.35	112.42	118.22
2	B	750	HEM	CBA-CAA-C2A	-4.14	105.12	112.53
2	B	750	HEM	CAA-C2A-C1A	-4.05	122.61	127.01
2	A	750	HEM	CBD-CAD-C3D	-3.84	102.36	113.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	800	HW6	C05-C06-N01	-3.62	118.88	122.96
3	A	760	H4B	N3-C2-N1	-3.25	120.21	125.53
2	B	750	HEM	CBD-CAD-C3D	-3.17	104.31	113.55
4	A	800	HW6	C05-C06-N01	-3.14	119.41	122.96
3	B	760	H4B	N3-C2-N1	-2.96	120.67	125.53
4	A	800	HW6	C21-C22-C23	-2.96	116.39	118.84
2	B	750	HEM	C3C-CAC-CBC	-2.96	119.92	124.46
4	A	800	HW6	C26-C25-C24	-2.93	117.76	121.69
4	B	800	HW6	C26-C25-CL2	-2.61	115.90	119.14
4	B	800	HW6	C26-C25-C24	-2.59	118.21	121.69
4	A	800	HW6	C26-C25-CL2	-2.48	116.06	119.14
2	A	750	HEM	C3B-C4B-NB	-2.37	107.09	111.63
2	B	750	HEM	C3B-C4B-NB	-2.29	107.25	111.63
2	A	750	HEM	CAA-CBA-CGA	-2.24	108.64	112.75
4	A	800	HW6	C12-C13-C14	-2.24	104.44	113.90
2	A	750	HEM	C3B-CAB-CBB	-2.17	121.13	124.46
4	B	800	HW6	C25-C24-C23	2.09	119.37	117.73
4	A	800	HW6	N02-C02-N01	2.20	120.51	116.50
3	B	760	H4B	C2-N1-C8A	2.24	119.56	114.54
3	B	760	H4B	N2-C2-N3	2.37	121.13	117.20
3	A	760	H4B	C2-N1-C8A	2.38	119.88	114.54
4	B	800	HW6	C10-O09-C3'	2.38	119.59	113.95
3	B	760	H4B	C4A-C8A-N8	2.41	121.27	118.43
2	B	750	HEM	C3B-C4B-CHC	2.60	126.82	123.16
2	A	750	HEM	CMD-C2D-C3D	2.70	126.31	114.35
2	B	750	HEM	C2D-C3D-C4D	2.74	106.15	101.50
2	B	750	HEM	CMD-C2D-C3D	2.77	126.62	114.35
2	A	750	HEM	C2D-C3D-C4D	2.93	106.46	101.50
4	B	800	HW6	F23-C23-C24	2.94	122.13	118.22
3	A	760	H4B	N2-C2-N3	2.97	122.11	117.20
4	A	800	HW6	C25-C26-C21	3.07	122.74	119.72
3	A	760	H4B	C4A-C8A-N8	3.71	122.80	118.43
4	B	800	HW6	C25-C26-C21	3.80	123.45	119.72
2	A	750	HEM	C3B-C4B-CHC	3.82	128.54	123.16
2	B	750	HEM	CAD-C3D-C2D	3.88	124.37	113.22
4	A	800	HW6	F23-C23-C24	3.94	123.46	118.22
2	A	750	HEM	CAD-C3D-C4D	4.12	126.99	112.47
3	B	760	H4B	C4-N3-C2	4.12	121.66	115.94
2	B	750	HEM	CMB-C2B-C3B	4.15	126.90	116.53
2	B	750	HEM	CMC-C2C-C3C	4.18	126.95	116.53
2	A	750	HEM	CMB-C2B-C3B	4.33	127.33	116.53
3	A	760	H4B	C4-N3-C2	4.40	122.05	115.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CMC-C2C-C3C	4.45	127.65	116.53
2	A	750	HEM	CAD-C3D-C2D	4.63	126.54	113.22
3	A	760	H4B	C4-C4A-C8A	4.72	118.84	114.56
2	B	750	HEM	CAD-C3D-C4D	4.79	129.37	112.47
4	B	800	HW6	C24-C25-CL2	5.42	125.88	119.14
4	A	800	HW6	C24-C25-CL2	5.66	126.17	119.14
3	B	760	H4B	C4-C4A-C8A	6.03	120.02	114.56
4	A	800	HW6	C02-N01-C06	6.62	122.93	118.23
4	B	800	HW6	C02-N01-C06	7.10	123.28	118.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	1	0
3	A	760	H4B	2	0
4	A	800	HW6	2	0
2	B	750	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	1.29	100 (24%) ⓘ ⓘ	27, 53, 95, 122	0
1	B	411/422 (97%)	0.74	43 (10%) ⓘ ⓘ	26, 42, 67, 97	0
All	All	818/844 (96%)	1.01	143 (17%) ⓘ ⓘ	26, 47, 88, 122	0

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	716	TRP	8.8
1	B	300	PHE	8.8
1	A	715	VAL	8.5
1	B	350	THR	8.4
1	B	348	VAL	8.2
1	A	488	PRO	8.0
1	A	351	LYS	6.9
1	A	352	ASP	6.6
1	A	355	PHE	6.1
1	A	350	THR	5.9
1	B	351	LYS	5.8
1	B	352	ASP	5.7
1	A	486	LYS	5.1
1	B	718	GLY	5.1
1	B	619	ARG	5.0
1	A	491	SER	4.9
1	A	506	ILE	4.9
1	A	593	ILE	4.7
1	A	386	LYS	4.6
1	A	714	HIS	4.5
1	A	503	GLU	4.3
1	A	567	VAL	4.3
1	A	300	PHE	4.3
1	A	712	ASN	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	678	TRP	4.2
1	A	619	ARG	4.2
1	A	507	GLN	4.1
1	A	489	ASP	4.1
1	A	391	THR	4.0
1	A	415	CYS	3.9
1	A	388	ILE	3.9
1	A	493	LEU	3.8
1	A	392	SER	3.8
1	A	713	THR	3.8
1	A	706	TYR	3.7
1	A	679	ILE	3.7
1	A	551	PHE	3.7
1	B	677	VAL	3.7
1	A	469	LYS	3.7
1	A	677	VAL	3.7
1	A	514	ARG	3.7
1	A	487	GLN	3.6
1	B	349	ARG	3.5
1	B	620	LYS	3.5
1	A	490	GLY	3.5
1	A	505	CYS	3.4
1	B	310	VAL	3.4
1	A	299	ARG	3.4
1	A	511	LYS	3.4
1	A	682	PRO	3.4
1	B	389	GLU	3.3
1	A	479	LEU	3.3
1	A	552	ASP	3.3
1	A	393	THR	3.3
1	A	385	ASN	3.3
1	A	389	GLU	3.3
1	A	353	GLN	3.3
1	A	504	ILE	3.3
1	B	567	VAL	3.2
1	A	591	THR	3.2
1	A	492	THR	3.2
1	A	676	TRP	3.1
1	A	609	GLU	3.1
1	B	691	PHE	3.1
1	A	470	HIS	3.1
1	A	390	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	679	ILE	3.1
1	A	322	LEU	3.0
1	B	479	LEU	3.0
1	B	566	ALA	3.0
1	A	681	PRO	3.0
1	A	584	PHE	2.9
1	A	595	VAL	2.9
1	B	713	THR	2.9
1	A	680	VAL	2.9
1	B	680	VAL	2.9
1	A	683	MET	2.9
1	B	591	THR	2.9
1	A	485	TYR	2.9
1	B	681	PRO	2.8
1	B	353	GLN	2.8
1	B	676	TRP	2.8
1	A	594	GLY	2.8
1	A	667	ARG	2.7
1	A	373	GLY	2.7
1	B	561	TRP	2.7
1	A	588	TYR	2.7
1	A	685	GLY	2.7
1	B	678	TRP	2.7
1	B	321	THR	2.6
1	A	494	GLY	2.6
1	A	321	THR	2.6
1	A	704	PHE	2.6
1	B	682	PRO	2.6
1	A	466	THR	2.6
1	A	570	MET	2.5
1	A	354	LEU	2.5
1	B	667	ARG	2.5
1	B	683	MET	2.5
1	A	620	LYS	2.5
1	A	711	TRP	2.5
1	A	384	VAL	2.4
1	B	480	ILE	2.4
1	B	375	LYS	2.4
1	A	566	ALA	2.4
1	A	370	LYS	2.4
1	A	568	SER	2.4
1	A	480	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	562	TYR	2.3
1	A	356	PRO	2.3
1	A	416	VAL	2.3
1	B	299	ARG	2.3
1	A	500	GLN	2.3
1	A	660	LYS	2.3
1	A	592	GLU	2.3
1	B	416	VAL	2.3
1	B	715	VAL	2.3
1	A	312	LEU	2.3
1	B	311	VAL	2.2
1	B	633	ILE	2.2
1	B	687	ILE	2.2
1	A	318	LEU	2.2
1	A	645	LYS	2.2
1	B	615	ASP	2.2
1	B	323	GLU	2.2
1	B	590	GLY	2.2
1	A	617	ASP	2.2
1	A	562	TYR	2.2
1	A	381	LEU	2.2
1	A	582	CYS	2.1
1	B	415	CYS	2.1
1	A	528	GLY	2.1
1	A	510	TRP	2.1
1	A	561	TRP	2.1
1	A	565	PRO	2.1
1	B	685	GLY	2.1
1	A	550	LYS	2.1
1	A	710	PRO	2.1
1	A	643	SER	2.0
1	A	512	ALA	2.0
1	A	382	GLU	2.0
1	A	572	LEU	2.0
1	A	369	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	ACT	A	860	4/4	0.94	0.25	5.00	53,53,53,54	0
5	ACT	B	860	4/4	0.96	0.17	3.74	47,47,49,49	0
4	HW6	A	800	28/28	0.79	0.30	1.68	43,58,65,66	0
2	HEM	B	750	43/43	0.98	0.20	1.03	22,30,45,56	0
4	HW6	B	800	28/28	0.91	0.22	1.00	38,49,57,58	0
2	HEM	A	750	43/43	0.97	0.23	0.82	28,31,42,51	0
3	H4B	B	760	17/17	0.96	0.19	0.60	32,36,39,41	0
3	H4B	A	760	17/17	0.95	0.19	0.14	37,40,45,47	0
6	ZN	A	900	1/1	0.99	0.09	-0.67	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.