



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 7, 2016 – 06:21 PM EDT

PDB ID : 4UHS  
Title : Crystal structure of the receiver domain of CpxR from E. coli (tetragonal form)  
Authors : Mechaly, A.E.; Alzari, P.M.A.  
Deposited on : 2015-03-25  
Resolution : 5.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 5.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1119 (6.22-3.60)
Clashscore	102246	1019 (6.22-3.66)
Ramachandran outliers	100387	1158 (6.22-3.60)
Sidechain outliers	100360	1136 (6.22-3.60)
RSRZ outliers	91569	1122 (6.22-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	<div> <div>2%</div> <div>88%</div> <div>7%</div> </div>
1	B	136	<div> <div>4%</div> <div>85%</div> <div>8%</div> <div>7%</div> </div>
1	C	136	<div> <div>4%</div> <div>86%</div> <div>6%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	B	1128	-	-	-	X



## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3003 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REGULATORY PROTEIN CPXR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	0	0
			1000	630	175	191	4			
1	B	126	Total	C	N	O	S	0	0	0
			1000	630	175	191	4			
1	C	126	Total	C	N	O	S	0	0	0
			1000	630	175	191	4			

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	LYS	-	EXPRESSION TAG	UNP P0AE88
A	125	LEU	-	EXPRESSION TAG	UNP P0AE88
A	126	ALA	-	EXPRESSION TAG	UNP P0AE88
A	127	ALA	-	EXPRESSION TAG	UNP P0AE88
A	128	ALA	-	EXPRESSION TAG	UNP P0AE88
A	129	LEU	-	EXPRESSION TAG	UNP P0AE88
A	130	GLU	-	EXPRESSION TAG	UNP P0AE88
A	131	HIS	-	EXPRESSION TAG	UNP P0AE88
A	132	HIS	-	EXPRESSION TAG	UNP P0AE88
A	133	HIS	-	EXPRESSION TAG	UNP P0AE88
A	134	HIS	-	EXPRESSION TAG	UNP P0AE88
A	135	HIS	-	EXPRESSION TAG	UNP P0AE88
A	136	HIS	-	EXPRESSION TAG	UNP P0AE88
B	124	LYS	-	EXPRESSION TAG	UNP P0AE88
B	125	LEU	-	EXPRESSION TAG	UNP P0AE88
B	126	ALA	-	EXPRESSION TAG	UNP P0AE88
B	127	ALA	-	EXPRESSION TAG	UNP P0AE88
B	128	ALA	-	EXPRESSION TAG	UNP P0AE88
B	129	LEU	-	EXPRESSION TAG	UNP P0AE88
B	130	GLU	-	EXPRESSION TAG	UNP P0AE88
B	131	HIS	-	EXPRESSION TAG	UNP P0AE88
B	132	HIS	-	EXPRESSION TAG	UNP P0AE88
B	133	HIS	-	EXPRESSION TAG	UNP P0AE88

*Continued on next page...*



*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	134	HIS	-	EXPRESSION TAG	UNP P0AE88
B	135	HIS	-	EXPRESSION TAG	UNP P0AE88
B	136	HIS	-	EXPRESSION TAG	UNP P0AE88
C	124	LYS	-	EXPRESSION TAG	UNP P0AE88
C	125	LEU	-	EXPRESSION TAG	UNP P0AE88
C	126	ALA	-	EXPRESSION TAG	UNP P0AE88
C	127	ALA	-	EXPRESSION TAG	UNP P0AE88
C	128	ALA	-	EXPRESSION TAG	UNP P0AE88
C	129	LEU	-	EXPRESSION TAG	UNP P0AE88
C	130	GLU	-	EXPRESSION TAG	UNP P0AE88
C	131	HIS	-	EXPRESSION TAG	UNP P0AE88
C	132	HIS	-	EXPRESSION TAG	UNP P0AE88
C	133	HIS	-	EXPRESSION TAG	UNP P0AE88
C	134	HIS	-	EXPRESSION TAG	UNP P0AE88
C	135	HIS	-	EXPRESSION TAG	UNP P0AE88
C	136	HIS	-	EXPRESSION TAG	UNP P0AE88

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0




### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN CPXR

Chain A: 





- Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN CPXR

Chain B: 



- Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN CPXR

Chain C: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.22Å 114.22Å 356.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.09 – 5.00 48.09 – 5.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.09-5.00) 99.9 (48.09-5.00)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 5.10Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.261 , 0.267 0.259 , 0.265	Depositor DCC
$R_{free}$ test set	437 reflections (4.56%)	DCC
Wilson B-factor (Å <sup>2</sup> )	257.3	Xtriage
Anisotropy	0.659	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 264.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.32$ , $\langle L^2 \rangle = 0.16$	Xtriage
Outliers	2 of 5454 reflections (0.037%)	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	3003	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	260.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	1/1011 (0.1%)	0.68	0/1365
1	B	0.44	0/1011	0.64	0/1365
1	C	0.51	1/1011 (0.1%)	0.72	1/1365 (0.1%)
All	All	0.50	2/3033 (0.1%)	0.68	1/4095 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	LEU	CB-CG	-7.39	1.31	1.52
1	C	84	LEU	CB-CG	-5.14	1.37	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	84	LEU	CD1-CG-CD2	5.74	127.71	110.50

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1000	0	1033	12	0
1	B	1000	0	1033	11	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1000	0	1033	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
All	All	3003	0	3099	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:84:LEU:N	1:C:84:LEU:HD13	1.98	0.78
1:C:98:LEU:HD22	1:C:107:LEU:HD13	1.67	0.75
1:A:84:LEU:HD13	1:A:84:LEU:N	2.02	0.74
1:B:86:ARG:NH2	1:B:99:PRO:HB3	2.11	0.65
1:B:99:PRO:O	1:B:102:PHE:HB3	1.99	0.62
1:A:84:LEU:HG	1:C:84:LEU:HG	1.82	0.61
1:A:91:GLU:HB2	1:C:88:LEU:HD11	1.82	0.60
1:A:91:GLU:CB	1:C:88:LEU:HD11	2.34	0.58
1:B:105:ARG:HH12	1:C:84:LEU:HB2	1.70	0.56
1:A:84:LEU:CD2	1:C:84:LEU:HG	2.38	0.54
1:A:84:LEU:HD21	1:C:84:LEU:HG	1.90	0.54
1:C:84:LEU:HD13	1:C:84:LEU:H	1.73	0.53
1:A:84:LEU:CG	1:C:84:LEU:HG	2.38	0.52
1:B:105:ARG:HH22	1:C:82:SER:HB3	1.76	0.51
1:A:84:LEU:HD21	1:C:84:LEU:CD2	2.40	0.51
1:C:82:SER:N	1:C:86:ARG:HH12	2.09	0.50
1:B:82:SER:O	1:B:85:ASP:HB2	2.11	0.49
1:B:105:ARG:HH12	1:C:84:LEU:CB	2.27	0.48
1:B:86:ARG:HH22	1:B:99:PRO:HB3	1.77	0.47
1:A:91:GLU:HG3	1:C:88:LEU:HD21	1.97	0.47
1:B:89:GLY:HA3	1:B:97:TYR:OH	2.16	0.46
1:A:84:LEU:HA	1:A:84:LEU:HD12	1.58	0.46
1:C:98:LEU:CD2	1:C:107:LEU:HD13	2.44	0.44
1:B:98:LEU:HD22	1:B:107:LEU:HD13	1.99	0.43
1:A:71:GLN:O	1:B:117:ARG:NH2	2.52	0.42
1:C:89:GLY:HA3	1:C:97:TYR:OH	2.19	0.42
1:A:6:LEU:CD1	1:A:49:LEU:HD12	2.51	0.41
1:B:105:ARG:HH22	1:C:82:SER:CB	2.35	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/136 (91%)	121 (98%)	3 (2%)	0	100	100
1	B	124/136 (91%)	123 (99%)	1 (1%)	0	100	100
1	C	124/136 (91%)	121 (98%)	3 (2%)	0	100	100
All	All	372/408 (91%)	365 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	111/120 (92%)	109 (98%)	2 (2%)	66	87
1	B	111/120 (92%)	111 (100%)	0	100	100
1	C	111/120 (92%)	109 (98%)	2 (2%)	66	87
All	All	333/360 (92%)	329 (99%)	4 (1%)	78	90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	GLU
1	A	84	LEU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	C	84	LEU
1	C	85	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	126/136 (92%)	-0.01	3 (2%) 62 54	125, 249, 289, 300	0
1	B	126/136 (92%)	0.25	5 (3%) 42 35	191, 277, 300, 300	0
1	C	126/136 (92%)	0.21	6 (4%) 34 28	142, 275, 295, 299	0
All	All	378/408 (92%)	0.15	14 (3%) 45 37	125, 272, 300, 300	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	55	PRO	4.8
1	B	56	LYS	4.3
1	A	56	LYS	3.6
1	C	28	VAL	2.9
1	B	71	GLN	2.8
1	C	29	ILE	2.8
1	C	27	ASN	2.7
1	C	2	ASN	2.6
1	C	56	LYS	2.4
1	A	41	LEU	2.4
1	B	100	LYS	2.2
1	B	72	THR	2.1
1	A	30	VAL	2.0
1	C	45	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	1128	1/1	0.86	0.80	1.84	300,300,300,300	0
2	MG	A	1128	1/1	0.90	0.29	0.36	125,125,125,125	0
2	MG	C	1128	1/1	0.96	0.21	-1.41	129,129,129,129	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.