



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:40 PM GMT

PDB ID : 3UKU
Title : Structure of Arp2/3 complex with bound inhibitor CK-869
Authors : Nolen, B.J.; Han, M.
Deposited on : 2011-11-09
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	
2	B	394	
3	C	372	
4	D	300	
5	E	178	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	168	<div> <div>%</div> <div> <div></div> <div>53%</div> <div>42%</div> <div>.</div> <div>..</div> </div> </div>
7	G	151	<div> <div>7%</div> <div> <div></div> <div>56%</div> <div>34%</div> <div>•</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	C69	A	501	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 13460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACTIN-LIKE PROTEIN 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	0	0	0
			3197	2053	534	595	15			

- Molecule 2 is a protein called ACTIN-LIKE PROTEIN 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	195	Total	C	N	O	S	0	0	0
			1528	979	261	284	4			

- Molecule 3 is a protein called Actin-related protein 2/3 complex subunit 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	341	Total	C	N	O	S	0	0	0
			2633	1669	460	485	19			

- Molecule 4 is a protein called Actin-related protein 2/3 complex subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	277	Total	C	N	O	S	0	0	0
			2217	1410	381	418	8			

- Molecule 5 is a protein called Actin-related protein 2/3 complex subunit 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	173	Total	C	N	O	S	0	0	0
			1403	901	234	259	9			


- Molecule 6 is a protein called Actin-related protein 2/3 complex subunit 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	166	Total	C	N	O	S	0	0	0
			1364	871	238	246	9			

- Molecule 7 is a protein called Actin-related protein 2/3 complex subunit 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	138	Total	C	N	O	S	0	0	0
			971	609	171	190	1			

- Molecule 8 is (2S)-2-(3-BROMOPHENYL)-3-(2,4-DIMETHOXYPHENYL)-1,3-THIAZOLIDIN-4-ONE (three-letter code: C69) (formula: C₁₇H₁₆BrNO₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total	Br	C	N	O	S	0	0
			23	1	17	1	3	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	25	Total	O	0	0
			25	25		
9	B	11	Total	O	0	0
			11	11		
9	C	37	Total	O	0	0
			37	37		
9	D	26	Total	O	0	0
			26	26		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	E	6	Total 6	O 6	0	0
9	F	15	Total 15	O 15	0	0
9	G	4	Total 4	O 4	0	0




• Molecule 3: Actin-related protein 2/3 complex subunit 1B

Chain C: 50% 40% 8%



• Molecule 4: Actin-related protein 2/3 complex subunit 2

Chain D: 47% 40% 5% 8%




• Molecule 5: Actin-related protein 2/3 complex subunit 3

Chain E: 5% 39% 54% 2%






• Molecule 6: Actin-related protein 2/3 complex subunit 4



• Molecule 7: Actin-related protein 2/3 complex subunit 5



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	111.11Å 129.66Å 203.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.75 43.51 – 2.74	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.75) 90.6 (43.51-2.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.73Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.257 , 0.301 0.240 , 0.283	Depositor DCC
R_{free} test set	3866 reflections (5.80%)	DCC
Wilson B-factor (Å ²)	55.9	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 50.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 76990 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13460	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C69

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/3278	0.66	0/4447
2	B	0.40	0/1557	0.64	0/2110
3	C	0.47	0/2702	0.73	0/3673
4	D	0.42	0/2265	0.66	0/3062
5	E	0.34	0/1437	0.60	0/1940
6	F	0.46	0/1386	0.70	0/1858
7	G	0.33	0/982	0.56	0/1330
All	All	0.42	0/13607	0.66	0/18420

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3197	0	3145	188	0
2	B	1528	0	1530	107	0
3	C	2633	0	2560	151	0
4	D	2217	0	2158	149	0
5	E	1403	0	1398	108	0
6	F	1364	0	1403	101	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	G	971	0	895	54	0
8	A	23	0	16	11	0
9	A	25	0	0	1	0
9	B	11	0	0	0	0
9	C	37	0	0	1	0
9	D	26	0	0	0	0
9	E	6	0	0	0	0
9	F	15	0	0	0	0
9	G	4	0	0	0	0
All	All	13460	0	13105	787	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 787 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:182:LEU:HD11	2:B:278:VAL:H	1.12	1.12
3:C:183:THR:HG22	3:C:185:TRP:H	1.19	1.07
4:D:197:GLN:HE21	4:D:199:LEU:HD11	1.14	1.07
6:F:4:THR:HG23	6:F:55:ARG:HE	1.25	1.01
5:E:167:GLN:HE21	5:E:172:SER:HB2	1.29	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/418 (94%)	317 (81%)	61 (16%)	15 (4%)	4	11
2	B	193/394 (49%)	150 (78%)	36 (19%)	7 (4%)	4	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	337/372 (91%)	302 (90%)	31 (9%)	4 (1%)	16	43
4	D	273/300 (91%)	219 (80%)	41 (15%)	13 (5%)	3	7
5	E	171/178 (96%)	141 (82%)	24 (14%)	6 (4%)	4	13
6	F	164/168 (98%)	146 (89%)	16 (10%)	2 (1%)	16	43
7	G	132/151 (87%)	109 (83%)	21 (16%)	2 (2%)	13	36
All	All	1663/1981 (84%)	1384 (83%)	230 (14%)	49 (3%)	6	17

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	83	VAL
2	B	171	GLU
2	B	278	VAL
4	D	203	ARG
4	D	215	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/363 (96%)	332 (95%)	17 (5%)	31	62
2	B	163/345 (47%)	150 (92%)	13 (8%)	15	37
3	C	286/313 (91%)	272 (95%)	14 (5%)	31	62
4	D	239/264 (90%)	227 (95%)	12 (5%)	30	61
5	E	154/159 (97%)	148 (96%)	6 (4%)	39	72
6	F	153/155 (99%)	145 (95%)	8 (5%)	29	60
7	G	88/123 (72%)	83 (94%)	5 (6%)	25	55
All	All	1432/1722 (83%)	1357 (95%)	75 (5%)	29	60

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	21	THR
3	C	321	LEU
6	F	165	LEU
3	C	30	HIS
3	C	131	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	4	HIS
3	C	107	ASN
6	F	28	GLN
3	C	33	HIS
3	C	46	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	C69	A	501	-	25,25,25	1.71	2 (8%)	34,35,35	2.45	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	C69	A	501	-	-	0/12/25/25	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	501	C69	CAE-NAF	-6.91	1.33	1.43
8	A	501	C69	CAQ-NAF	-3.40	1.33	1.37

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	501	C69	CAQ-CAR-SAS	-7.99	100.94	107.69
8	A	501	C69	CAB-OAA-CAD	-5.15	109.73	117.54
8	A	501	C69	CAL-OAM-CAN	-4.00	108.15	117.51
8	A	501	C69	OAW-CAQ-NAF	-3.19	122.36	125.12
8	A	501	C69	OAA-CAD-CAC	-3.17	118.80	124.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	501	C69	11	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/418 (95%)	0.23	14 (3%) 48 41	28, 60, 92, 107	0
2	B	195/394 (49%)	0.65	25 (12%) 5 3	35, 73, 109, 115	0
3	C	341/372 (91%)	0.05	1 (0%) 94 93	32, 52, 78, 95	0
4	D	277/300 (92%)	0.12	4 (1%) 78 73	30, 60, 96, 114	0
5	E	173/178 (97%)	0.32	9 (5%) 31 24	52, 79, 111, 119	0
6	F	166/168 (98%)	-0.00	1 (0%) 90 88	28, 56, 74, 87	0
7	G	138/151 (91%)	0.54	11 (7%) 15 10	49, 100, 118, 120	0
All	All	1689/1981 (85%)	0.23	65 (3%) 44 37	28, 63, 105, 120	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	ARG	5.4
2	B	178	LEU	5.4
5	E	154	ASP	4.8
7	G	151	VAL	4.7
2	B	174	SER	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
8	C69	A	501	23/23	0.92	0.24	0.95	71,75,83,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.