



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:40 PM GMT

PDB ID : 3UL1
Title : Mouse importin alpha: nucleoplasmin cNLS peptide complex
Authors : Marfori, M.; Forwood, J.K.; Lonhienne, T.G.; Kobe, B.
Deposited on : 2011-11-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	510	
2	A	22	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	426	3278	2090	553	625	10	5	8	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	20	MET	-	EXPRESSION TAG	UNP P52293
B	21	HIS	-	EXPRESSION TAG	UNP P52293
B	22	HIS	-	EXPRESSION TAG	UNP P52293
B	23	HIS	-	EXPRESSION TAG	UNP P52293
B	24	HIS	-	EXPRESSION TAG	UNP P52293
B	25	HIS	-	EXPRESSION TAG	UNP P52293
B	26	HIS	-	EXPRESSION TAG	UNP P52293
B	27	SER	-	EXPRESSION TAG	UNP P52293
B	28	SER	-	EXPRESSION TAG	UNP P52293
B	29	GLY	-	EXPRESSION TAG	UNP P52293
B	30	LEU	-	EXPRESSION TAG	UNP P52293
B	31	VAL	-	EXPRESSION TAG	UNP P52293
B	32	PRO	-	EXPRESSION TAG	UNP P52293
B	33	ARG	-	EXPRESSION TAG	UNP P52293
B	34	GLY	-	EXPRESSION TAG	UNP P52293
B	35	SER	-	EXPRESSION TAG	UNP P52293
B	36	GLY	-	EXPRESSION TAG	UNP P52293
B	37	MET	-	EXPRESSION TAG	UNP P52293
B	38	LEU	-	EXPRESSION TAG	UNP P52293
B	39	GLU	-	EXPRESSION TAG	UNP P52293
B	40	THR	-	EXPRESSION TAG	UNP P52293
B	41	ALA	-	EXPRESSION TAG	UNP P52293
B	42	ALA	-	EXPRESSION TAG	UNP P52293
B	43	ALA	-	EXPRESSION TAG	UNP P52293
B	44	LEU	-	EXPRESSION TAG	UNP P52293
B	45	PHE	-	EXPRESSION TAG	UNP P52293
B	46	GLU	-	EXPRESSION TAG	UNP P52293

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	47	ARG	-	EXPRESSION TAG	UNP P52293
B	48	ASN	-	EXPRESSION TAG	UNP P52293
B	49	HIS	-	EXPRESSION TAG	UNP P52293
B	50	MET	-	EXPRESSION TAG	UNP P52293
B	51	ASP	-	EXPRESSION TAG	UNP P52293
B	52	SER	-	EXPRESSION TAG	UNP P52293
B	53	PRO	-	EXPRESSION TAG	UNP P52293
B	54	ASP	-	EXPRESSION TAG	UNP P52293
B	55	LEU	-	EXPRESSION TAG	UNP P52293
B	56	GLY	-	EXPRESSION TAG	UNP P52293
B	57	THR	-	EXPRESSION TAG	UNP P52293
B	58	ASP	-	EXPRESSION TAG	UNP P52293
B	59	ASP	-	EXPRESSION TAG	UNP P52293
B	60	ASP	-	EXPRESSION TAG	UNP P52293
B	61	ASP	-	EXPRESSION TAG	UNP P52293
B	62	LEU	-	EXPRESSION TAG	UNP P52293
B	63	ALA	-	EXPRESSION TAG	UNP P52293
B	64	MET	-	EXPRESSION TAG	UNP P52293
B	65	ALA	-	EXPRESSION TAG	UNP P52293
B	66	ASP	-	EXPRESSION TAG	UNP P52293
B	67	ILE	-	EXPRESSION TAG	UNP P52293
B	68	GLY	-	EXPRESSION TAG	UNP P52293
B	69	SER	-	EXPRESSION TAG	UNP P52293

- Molecule 2 is a protein called Nucleoplasmin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	A	21	Total	C	N	O	0	0	0
			156	97	32	27			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	151	GLY	-	EXPRESSION TAG	UNP P05221
A	152	SER	-	EXPRESSION TAG	UNP P05221

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	279	Total	O	0	0
			279	279		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Importin subunit alpha-2



• Molecule 2: Nucleoplasmin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.16Å 90.09Å 99.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.07 – 1.90 42.07 – 1.90	Depositor EDS
% Data completeness (in resolution range)	92.5 (42.07-1.90) 98.2 (42.07-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.89Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.191 , 0.222 0.193 , 0.223	Depositor DCC
R_{free} test set	2821 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.467	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	0 of 55629 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3726	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.28	0/3363	0.44	0/4581
2	A	0.29	0/156	0.47	0/202
All	All	0.28	0/3519	0.44	0/4783

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3278	0	3379	23	0
2	A	156	0	182	0	0
3	A	13	0	0	0	0
3	B	279	0	0	3	0
All	All	3726	0	3561	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:VAL:HG12	1:B:75:ASN:HD22	1.62	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:425:LEU:HG	1:B:440:ILE:HG23	1.81	0.62
1:B:372:GLN:NE2	3:B:723:HOH:O	2.05	0.60
1:B:290:VAL:HG13	1:B:330:LYS:HE2	1.87	0.57
1:B:456:GLU:O	1:B:459:LYS:HG3	2.07	0.54
1:B:482:GLU:HA	1:B:485:TYR:CE2	2.43	0.53
1:B:74:VAL:HG12	1:B:75:ASN:ND2	2.26	0.50
1:B:330:LYS:HE3	3:B:702:HOH:O	2.12	0.49
1:B:482:GLU:HA	1:B:485:TYR:CD2	2.50	0.47
1:B:411:GLU:H	1:B:411:GLU:CD	2.17	0.47
1:B:284:GLU:O	1:B:288:MET:HG3	2.15	0.46
1:B:261:HIS:HE1	3:B:664:HOH:O	1.98	0.46
1:B:273:TRP:CD2	1:B:312:PRO:HB3	2.51	0.45
1:B:290:VAL:CG1	1:B:330:LYS:HD3	2.47	0.44
1:B:472:LYS:O	1:B:476:LEU:HG	2.17	0.44
1:B:238:ARG:O	1:B:239:ASN:HB2	2.18	0.44
1:B:386:LEU:HD21	1:B:425:LEU:HD13	2.00	0.43
1:B:73:THR:HA	1:B:76:TRP:NE1	2.32	0.43
1:B:223:CYS:SG	1:B:227:ARG:CZ	3.08	0.42
1:B:426:MET:HE1	1:B:444:ILE:HD11	2.01	0.42
1:B:290:VAL:HG11	1:B:330:LYS:HD3	2.02	0.42
1:B:262:HIS:O	1:B:268:LEU:HD21	2.20	0.42
1:B:386:LEU:HD21	1:B:425:LEU:CD1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	433/510 (85%)	427 (99%)	6 (1%)	0	100	100
2	A	19/22 (86%)	19 (100%)	0	0	100	100
All	All	452/532 (85%)	446 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	366/426 (86%)	361 (99%)	5 (1%)	74	71
2	A	15/15 (100%)	15 (100%)	0	100	100
All	All	381/441 (86%)	376 (99%)	5 (1%)	76	73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	109	GLN
1	B	240	LYS
1	B	247	ASP
1	B	352	GLN
1	B	459	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	75	ASN
1	B	228	ASN
1	B	350	ASN
1	B	372	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	426/510 (83%)	0.58	44 (10%) 9 9	19, 32, 81, 113	0
2	A	21/22 (95%)	0.34	1 (4%) 34 37	25, 40, 59, 69	0
All	All	447/532 (84%)	0.57	45 (10%) 9 10	19, 33, 80, 113	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	485	TYR	11.8
1	B	489	LEU	8.2
1	B	476	LEU	6.4
1	B	492	ILE	6.1
1	B	482	GLU	5.5
1	B	74	VAL	5.2
1	B	478	ARG	4.9
1	B	479	HIS	4.9
1	B	496	PHE	4.9
1	B	484	VAL	4.7
1	B	474	GLU	4.6
1	B	475	ALA	4.5
1	B	493	GLU	4.4
1	B	481	ASN	4.4
1	B	486	LYS	4.3
1	B	491	LEU	4.1
1	B	480	GLU	4.1
1	B	432	LYS	4.1
1	B	483	SER	3.9
1	B	490	ASN	3.9
1	B	477	GLN	3.5
1	B	88	ASN	3.4
1	B	497	SER	3.4
1	B	107	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	108	LYS	3.2
1	B	444	ILE	3.1
1	B	487	ALA	3.1
1	B	431	ALA	3.1
1	B	471	ASP	3.1
1	B	470	LEU	2.7
1	B	109	GLN	2.6
1	B	452	GLU	2.6
1	B	455	GLY	2.5
2	A	159	ALA	2.4
1	B	434	THR	2.4
1	B	494	LYS	2.4
1	B	189	ILE	2.3
1	B	437	ILE	2.3
1	B	159	VAL	2.2
1	B	73	THR	2.2
1	B	186	LEU	2.2
1	B	495	TYR	2.1
1	B	419	CYS	2.1
1	B	473	ILE	2.0
1	B	458	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.