



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:02 PM GMT

PDB ID : 4UPN  
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with N,N''-{[(2S)-3-aminopropane-1,2-diyl]bis(oxymethanediylbenzene-3,1-diyl)}di thiophene-2-carboximidamide  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2014-06-17  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	 5% 81% 14% ••
1	B	422	 2% 82% 14% ••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEM	B	750	-	-	-	X
4	7LN	A	800	-	-	-	X
4	7LN	B	800	-	-	-	X
5	ACT	A	860	-	-	-	X
5	ACT	B	860	-	-	-	X
6	ZN	A	1717	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEURONAL NITRIC OXIDE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	0	0	1
			3314	2121	567	605	21			
1	B	411	Total	C	N	O	S	0	0	0
			3345	2140	574	610	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 17	C 9	N 5	O 3	0	0
3	B	1	Total 17	C 9	N 5	O 3	0	0

- Molecule 4 is N,N''-{[(2S)-3-AMINOPROPANE-1,2-DIYL]BIS(OXYMETHANEDIYLBENZENE-3,1-DIYL)}DITHIOPHENE-2-CARBOXIMIDAMIDE (three-letter code: 7LN) (formula: C<sub>27</sub>H<sub>29</sub>N<sub>5</sub>O<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			36	27	5	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			36	27	5	2	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula:  $Zn$ ).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	96	Total	O	0	0
			96	96		
7	B	117	Total	O	0	0
			117	117		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: NEURONAL NITRIC OXIDE SYNTHASE



#### • Molecule 1: NEURONAL NITRIC OXIDE SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.07Å 111.27Å 165.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.38 – 2.09 49.33 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.6 (49.38-2.09) 97.7 (49.33-2.09)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.199 , 0.245 0.202 , 0.245	Depositor DCC
$R_{free}$ test set	2772 reflections (5.18%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.3	Xtriage
Anisotropy	0.689	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 56250 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7073	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, H4B, 7LN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.85	4/3407 (0.1%)	0.88	3/4623 (0.1%)
1	B	1.08	3/3438 (0.1%)	0.98	7/4661 (0.2%)
All	All	0.97	7/6845 (0.1%)	0.93	10/9284 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	441	TYR	CE2-CZ	9.43	1.50	1.38
1	A	676	TRP	CG-CD1	7.50	1.47	1.36
1	B	421	TRP	CB-CG	6.35	1.61	1.50
1	A	416	VAL	C-O	6.16	1.35	1.23
1	A	419	ILE	N-CA	5.97	1.58	1.46
1	A	532	GLU	C-O	5.63	1.34	1.23
1	B	471	ASP	C-O	5.30	1.33	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	430	ARG	NE-CZ-NH1	8.99	124.80	120.30
1	A	410	ARG	NE-CZ-NH1	8.93	124.77	120.30
1	B	430	ARG	NE-CZ-NH2	-6.95	116.83	120.30
1	B	471	ASP	CB-CG-OD1	6.23	123.90	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	489	ASP	CB-CG-OD2	-6.13	112.78	118.30
1	B	489	ASP	CB-CG-OD1	6.03	123.73	118.30
1	B	675	ASP	CB-CG-OD1	5.95	123.65	118.30
1	A	572	LEU	CB-CG-CD2	-5.72	101.28	111.00
1	B	371	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	A	586	GLY	N-CA-C	-5.05	100.47	113.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	588	TYR	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3314	0	3221	40	0
1	B	3345	0	3259	32	0
2	A	43	0	30	5	0
2	B	43	0	30	4	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	36	0	28	8	0
4	B	36	0	28	4	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0
7	A	96	0	0	6	0
7	B	117	0	0	2	0
All	All	7073	0	6632	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:800:7LN:H36	4:A:800:7LN:H24	1.63	0.80
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.64	0.79
4:B:800:7LN:H23	7:B:2096:HOH:O	1.82	0.78
2:A:750:HEM:HBB2	2:A:750:HEM:HHC	1.76	0.67
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	1.76	0.67
1:A:592:GLU:OE1	4:A:800:7LN:N07	2.31	0.64
4:A:800:7LN:C24	4:A:800:7LN:H36	2.29	0.61
1:B:325:GLY:O	1:B:332:MET:HG3	2.00	0.61
1:B:322:LEU:HD13	1:B:699:ARG:HH21	1.67	0.59
1:A:601:ASN:HB2	7:A:2072:HOH:O	2.02	0.59
1:A:337:LEU:HD21	4:A:800:7LN:H33	1.86	0.57
1:A:674:ALA:HB3	1:A:695:MET:HB3	1.87	0.55
1:A:380:ARG:HD3	1:A:400:GLU:OE2	2.07	0.54
1:A:439:PHE:CZ	1:A:443:CYS:SG	3.01	0.53
1:B:355:PHE:CE1	1:B:385:ASN:HB2	2.43	0.53
1:B:388:ILE:O	1:B:392:SER:N	2.39	0.53
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.89	0.53
4:A:800:7LN:C36	4:A:800:7LN:H24	2.36	0.52
2:A:750:HEM:CMC	2:A:750:HEM:HBC2	2.38	0.51
1:B:584:PHE:CD2	2:B:750:HEM:CAC	2.94	0.50
4:A:800:7LN:C05	4:A:800:7LN:C12	2.87	0.50
1:A:701:THR:OG1	1:A:702:PRO:HA	2.11	0.49
1:B:572:LEU:HD21	1:B:574:ILE:HD11	1.95	0.49
1:A:321:THR:HG22	7:A:2003:HOH:O	2.12	0.49
1:A:299:ARG:O	1:A:317:HIS:CE1	2.66	0.49
1:B:525:GLN:HG3	1:B:529:ASN:O	2.13	0.48
1:B:416:VAL:HG23	1:B:679:ILE:HG23	1.95	0.48
2:B:750:HEM:HBC2	2:B:750:HEM:CMC	2.44	0.48
2:B:750:HEM:C4B	4:B:800:7LN:S01	3.08	0.47
1:A:464:GLN:HB3	1:A:579:PHE:CE1	2.50	0.46
1:B:350:THR:OG1	1:B:353:GLN:NE2	2.48	0.46
1:A:684:SER:HB3	1:A:687:ILE:CG1	2.46	0.46
1:B:592:GLU:OE1	4:B:800:7LN:N07	2.49	0.46
1:B:492:THR:HG21	1:B:496:PRO:HG3	1.98	0.45
1:A:337:LEU:HD21	4:A:800:7LN:C33	2.45	0.45
1:A:473:ARG:HD3	7:A:2047:HOH:O	2.16	0.45
1:A:684:SER:O	1:A:685:GLY:C	2.54	0.45
1:B:593:ILE:HA	1:B:597:ASP:HB2	1.99	0.45
1:A:306:TRP:CD2	1:B:336:MET:HE3	2.52	0.45
1:A:664:ASN:OD1	1:A:667:ARG:NH2	2.50	0.45
1:A:523:LEU:C	1:A:524:LEU:HD23	2.37	0.45
1:A:330:ILE:HG23	1:A:330:ILE:O	2.16	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:TRP:CH2	2:A:750:HEM:HMC1	2.52	0.45
1:A:657:SER:HB2	7:A:2025:HOH:O	2.17	0.45
1:B:478:GLN:HB2	1:B:481:ARG:HG3	2.00	0.44
1:B:701:THR:HA	1:B:702:PRO:C	2.37	0.44
1:B:445:HIS:C	1:B:445:HIS:CD2	2.91	0.44
1:A:409:TRP:CE2	2:A:750:HEM:C2C	3.06	0.44
1:A:463:PRO:HG2	1:A:472:PHE:CZ	2.53	0.44
1:A:657:SER:CB	7:A:2025:HOH:O	2.66	0.44
1:A:306:TRP:CE2	1:B:336:MET:HE3	2.53	0.43
1:B:663:GLU:HB3	1:B:667:ARG:NH1	2.33	0.43
1:B:460:THR:O	1:B:582:CYS:HA	2.18	0.43
1:B:336:MET:HE2	1:B:336:MET:HB2	1.91	0.43
1:A:486:LYS:HB2	1:A:486:LYS:HE3	1.90	0.43
1:B:657:SER:HB2	7:B:2028:HOH:O	2.19	0.42
1:A:686:SER:HA	1:A:691:PHE:CG	2.54	0.42
1:B:496:PRO:HB2	1:B:602:SER:O	2.19	0.42
1:A:398:ASP:O	1:A:402:ILE:HD12	2.19	0.42
1:B:546:ILE:HG12	1:B:560:LYS:HA	2.02	0.42
1:A:476:ASN:HB3	1:A:478:GLN:O	2.20	0.42
1:A:508:GLN:HA	1:A:508:GLN:HE21	1.85	0.42
1:A:569:ASN:H	1:A:569:ASN:HD22	1.66	0.42
1:B:659:ILE:HD13	1:B:659:ILE:HA	1.94	0.42
1:A:686:SER:OG	1:B:682:PRO:HB2	2.20	0.41
1:A:332:MET:O	1:A:335:ILE:HG12	2.20	0.41
1:A:312:LEU:HB3	1:A:666:TYR:CD2	2.55	0.41
4:A:800:7LN:C23	7:A:2070:HOH:O	2.68	0.41
4:B:800:7LN:C12	4:B:800:7LN:C05	2.98	0.41
1:B:482:TYR:HA	1:B:518:ASP:O	2.21	0.41
1:A:306:TRP:CD1	1:B:336:MET:CE	3.04	0.41
1:B:303:VAL:HG23	1:B:312:LEU:HB2	2.03	0.41
1:A:445:HIS:C	1:A:445:HIS:CD2	2.93	0.41
1:A:684:SER:HB3	1:A:687:ILE:HD11	2.03	0.41
1:A:582:CYS:N	1:A:583:PRO:CD	2.83	0.41
1:A:306:TRP:CG	1:B:336:MET:HE3	2.57	0.40
1:A:686:SER:OG	1:B:595:VAL:CG1	2.70	0.40
1:B:302:LYS:HA	1:B:312:LEU:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/422 (96%)	388 (96%)	12 (3%)	4 (1%)	19	13
1	B	407/422 (96%)	393 (97%)	14 (3%)	0	100	100
All	All	811/844 (96%)	781 (96%)	26 (3%)	4 (0%)	34	30

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	ASP
1	A	685	GLY
1	A	309	ASP
1	A	465	ARG

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/377 (96%)	350 (96%)	13 (4%)	42	43
1	B	366/377 (97%)	353 (96%)	13 (4%)	42	43
All	All	729/754 (97%)	703 (96%)	26 (4%)	42	43

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	307	GLU
1	A	337	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	350	THR
1	A	370	LYS
1	A	454	ASN
1	A	486	LYS
1	A	508	GLN
1	A	523	LEU
1	A	547	ARG
1	A	569	ASN
1	A	645	LYS
1	A	660	LYS
1	A	715	VAL
1	B	337	LEU
1	B	353	GLN
1	B	390	SER
1	B	392	SER
1	B	423	LYS
1	B	454	ASN
1	B	481	ARG
1	B	486	LYS
1	B	503	GLU
1	B	507	GLN
1	B	547	ARG
1	B	627	ASP
1	B	717	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	454	ASN
1	A	508	GLN
1	A	569	ASN
1	A	697	ASN
1	B	353	GLN
1	B	364	GLN
1	B	454	ASN
1	B	535	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	750	1	30,50,50	2.73	12 (40%)	24,82,82	2.53	11 (45%)
3	H4B	A	760	-	13,18,18	1.73	4 (30%)	11,26,26	3.60	7 (63%)
4	7LN	A	800	-	38,39,39	1.48	4 (10%)	36,51,51	1.41	3 (8%)
5	ACT	A	860	-	1,3,3	1.51	0	0,3,3	0.00	-
2	HEM	B	750	1	30,50,50	2.44	8 (26%)	24,82,82	2.42	11 (45%)
3	H4B	B	760	-	13,18,18	1.47	2 (15%)	11,26,26	2.71	4 (36%)
4	7LN	B	800	-	38,39,39	1.23	6 (15%)	36,51,51	1.97	8 (22%)
5	ACT	B	860	-	1,3,3	1.01	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	7LN	A	800	-	-	1/21/29/29	0/4/4/4
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	7LN	B	800	-	-	1/21/29/29	0/4/4/4
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C2C-C1C	-8.43	1.36	1.52
2	B	750	HEM	C2D-C3D	-6.52	1.35	1.54
2	A	750	HEM	C2D-C3D	-6.02	1.36	1.54
2	A	750	HEM	C2C-C1C	-5.94	1.41	1.52
2	A	750	HEM	C3B-C4B	-5.77	1.46	1.51
2	A	750	HEM	C3D-C4D	-4.61	1.45	1.51
2	B	750	HEM	C3D-C4D	-3.97	1.46	1.51
3	A	760	H4B	C8A-N1	-3.50	1.28	1.34
4	A	800	7LN	C11-N07	-3.33	1.36	1.42
2	A	750	HEM	C3C-CAC	-3.29	1.45	1.51
4	B	800	7LN	C11-N07	-3.24	1.37	1.42
2	B	750	HEM	C2B-C1B	-2.77	1.42	1.51
2	B	750	HEM	C3C-CAC	-2.44	1.46	1.51
2	B	750	HEM	C3B-CAB	-2.26	1.47	1.51
4	A	800	7LN	C31-N27	-2.23	1.37	1.41
2	A	750	HEM	C2D-C1D	-2.20	1.44	1.51
2	A	750	HEM	C2A-C3A	-2.13	1.31	1.37
4	B	800	7LN	C26-N27	-2.01	1.35	1.39
2	A	750	HEM	C4A-CHB	2.05	1.45	1.39
2	A	750	HEM	CHD-C4C	2.16	1.41	1.36
2	A	750	HEM	FE-NB	2.23	2.09	1.97
4	B	800	7LN	C03-C02	2.33	1.51	1.35
3	A	760	H4B	C2-N2	2.34	1.38	1.34
4	B	800	7LN	C13-C14	2.39	1.43	1.38
2	B	750	HEM	C4C-NC	2.39	1.39	1.36
4	B	800	7LN	C05-S01	2.41	1.75	1.72
3	A	760	H4B	C11-C10	2.42	1.57	1.51
2	B	750	HEM	C1C-NC	2.47	1.39	1.36
3	A	760	H4B	C7-N8	2.64	1.49	1.46
3	B	760	H4B	O4-C4	2.76	1.31	1.24
3	B	760	H4B	C2-N3	2.93	1.40	1.35

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	800	7LN	C25-C26	3.03	1.49	1.45
2	A	750	HEM	CHC-C1C	3.42	1.44	1.36
4	A	800	7LN	C25-C26	4.40	1.51	1.45
4	A	800	7LN	C06-N07	5.29	1.37	1.30
2	A	750	HEM	C1C-NC	6.08	1.43	1.36

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	800	7LN	C23-C22-S21	-4.19	108.59	113.23
2	B	750	HEM	C1D-CHD-C4C	-3.92	119.27	125.82
4	B	800	7LN	C03-C02-S01	-3.87	108.95	113.23
2	B	750	HEM	CBA-CAA-C2A	-3.82	105.68	112.53
2	A	750	HEM	CBA-CAA-C2A	-3.43	106.37	112.53
3	A	760	H4B	N3-C2-N1	-3.34	120.05	125.53
4	A	800	7LN	C23-C22-S21	-3.33	109.54	113.23
2	B	750	HEM	C3B-C4B-NB	-2.83	106.21	111.63
4	A	800	7LN	C17-O18-C19	-2.76	106.21	112.81
2	A	750	HEM	C4B-CHC-C1C	-2.63	121.42	125.82
2	A	750	HEM	CAA-C2A-C3A	-2.62	121.52	129.00
4	B	800	7LN	C12-C13-C14	-2.48	116.70	120.24
3	A	760	H4B	C8A-C4A-N5	-2.42	115.80	118.85
3	B	760	H4B	N3-C2-N1	-2.29	121.77	125.53
2	B	750	HEM	C3C-CAC-CBC	-2.26	120.98	124.46
2	B	750	HEM	C4B-CHC-C1C	-2.21	122.12	125.82
4	B	800	7LN	C16-C11-N07	-2.00	115.97	120.53
4	B	800	7LN	C34-C35-C36	2.24	121.86	118.55
2	B	750	HEM	CMD-C2D-C3D	2.25	124.28	114.35
2	A	750	HEM	C2D-C3D-C4D	2.35	105.48	101.50
3	A	760	H4B	C4-N3-C2	2.58	119.52	115.94
2	B	750	HEM	CHC-C4B-NB	2.60	130.79	124.52
4	B	800	7LN	C14-C15-C16	2.65	122.47	118.55
2	A	750	HEM	C3C-CAC-CBC	2.71	128.61	124.46
2	A	750	HEM	CMD-C2D-C3D	2.77	126.61	114.35
3	B	760	H4B	N2-C2-N3	2.82	121.87	117.20
2	A	750	HEM	CAA-C2A-C1A	2.99	130.25	127.01
4	B	800	7LN	C13-C12-C11	3.10	124.21	119.79
2	B	750	HEM	CMC-C2C-C3C	3.20	124.53	116.53
2	A	750	HEM	CAA-CBA-CGA	3.40	118.97	112.75
2	A	750	HEM	CAD-C3D-C4D	3.65	125.36	112.47
2	B	750	HEM	CMB-C2B-C3B	3.82	126.06	116.53
3	A	760	H4B	N2-C2-N3	3.90	123.66	117.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	760	H4B	C4-N3-C2	4.12	121.65	115.94
2	B	750	HEM	CAD-C3D-C4D	4.26	127.50	112.47
3	A	760	H4B	C2-N1-C8A	5.00	125.77	114.54
2	B	750	HEM	CAD-C3D-C2D	5.07	127.79	113.22
4	A	800	7LN	C25-C26-N27	5.19	129.88	115.07
2	A	750	HEM	CMC-C2C-C3C	5.38	129.95	116.53
2	A	750	HEM	CAD-C3D-C2D	5.53	129.13	113.22
3	A	760	H4B	C4A-C8A-N8	5.65	125.08	118.43
3	B	760	H4B	C4-C4A-C8A	6.52	120.46	114.56
3	A	760	H4B	C4-C4A-C8A	6.57	120.51	114.56
4	B	800	7LN	C25-C26-N27	6.73	134.27	115.07

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	800	7LN	N06-C06-N07-C11
4	A	800	7LN	N06-C06-N07-C11

There are no ring outliers.

4 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	5	0
4	A	800	7LN	8	0
2	B	750	HEM	4	0
4	B	800	7LN	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	408/422 (96%)	0.53	23 (5%)	28 36	33, 59, 105, 132	0
1	B	411/422 (97%)	0.29	7 (1%)	73 78	31, 47, 76, 99	0
All	All	819/844 (97%)	0.41	30 (3%)	45 54	31, 52, 97, 132	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	5.3
1	A	355	PHE	5.1
1	A	551	PHE	4.7
1	B	718	GLY	4.2
1	A	558	GLY	3.7
1	A	299	ARG	3.6
1	B	350	THR	3.3
1	A	352	ASP	3.2
1	A	556	ASP	3.2
1	A	552	ASP	3.1
1	B	355	PHE	3.0
1	A	351	LYS	3.0
1	A	372	PHE	2.9
1	A	391	THR	2.9
1	B	321	THR	2.7
1	A	373	GLY	2.6
1	A	322	LEU	2.5
1	A	517	PHE	2.5
1	A	300	PHE	2.5
1	B	338	PRO	2.4
1	A	511	LYS	2.4
1	A	321	THR	2.3
1	A	366	TYR	2.3
1	A	393	THR	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	352	ASP	2.3
1	A	466	THR	2.3
1	A	386	LYS	2.2
1	A	488	PRO	2.1
1	A	392	SER	2.0
1	A	489	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	ZN	A	1717	1/1	0.99	0.17	4.17	55,55,55,55	0
4	7LN	A	800	36/36	0.93	0.20	3.81	37,65,129,132	0
4	7LN	B	800	36/36	0.91	0.19	2.84	36,58,109,110	0
5	ACT	B	860	4/4	0.94	0.19	2.58	55,58,59,59	0
2	HEM	B	750	43/43	0.97	0.17	2.56	32,33,41,45	0
5	ACT	A	860	4/4	0.94	0.16	2.47	60,63,64,64	0
2	HEM	A	750	43/43	0.98	0.13	-0.05	33,36,42,44	0
3	H4B	B	760	17/17	0.96	0.13	-0.24	36,39,41,42	0
3	H4B	A	760	17/17	0.96	0.11	-2.75	35,37,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.