



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:03 PM GMT

PDB ID : 4UPS
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with N-[3-({[(3S,5S)-5-{[(3-{[(Z)-imino(thiophen-2-yl)methyl]amino}benzyl)oxy]methyl}pyrrolidin-3-yl]oxy}methyl) phenyl]thiophene-2-carboximidamide
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-06-17
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	<div> <div>9%</div> <div>82%</div> <div>8%</div> <div>9%</div> </div>
1	B	443	<div> <div>9%</div> <div>84%</div> <div>6%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	6E5	B	800	-	-	-	X
5	ACT	A	861	-	-	-	X
5	ACT	B	860	-	-	-	X
6	GOL	A	880	-	-	-	X
6	GOL	B	880	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	404	Total	As	C	N	O	S	0	2	0
			3220	1	2048	564	591	16			
1	B	402	Total	As	C	N	O	S	0	0	0
			3201	1	2036	562	586	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	CONFLICT	UNP P29473
B	100	ARG	CYS	CONFLICT	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is N-[3-({[(3S,5S)-5-{{[(3-{{[(Z)-IMINO(THIOPHEN-2-YL)METHYL}AMINO}BENZYL)OXY]METHYL}PYRROLIDIN-3-YL]OXY}METHYL)PHENYL]THIOPHENE-2-CARBOXIMIDAMIDE (three-letter code: 6E5) (formula: $C_{29}H_{31}N_5O_2S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 38	C 29	N 5	O 2	S 2	0	0
4	B	1	Total 38	C 29	N 5	O 2	S 2	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	193	Total	O	0	0
			193	193		
8	B	153	Total	O	0	0
			153	153		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.78Å 106.47Å 157.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.33 – 1.95 39.30 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.6 (39.33-1.95) 98.7 (39.30-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.177 , 0.213 0.176 , 0.211	Depositor DCC
R_{free} test set	3516 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 70554 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6992	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, 6E5, H4B, CAS, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/3306	0.69	2/4502 (0.0%)
1	B	0.58	0/3280	0.70	2/4466 (0.0%)
All	All	0.58	0/6586	0.69	4/8968 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	B	240	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	294	LEU	CA-CB-CG	5.76	128.55	115.30
1	A	244	ARG	NE-CZ-NH1	5.34	122.97	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3220	0	3123	26	0
1	B	3201	0	3104	17	0
2	A	43	0	30	3	0
2	B	43	0	30	3	0
3	A	17	0	15	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	17	0	15	0	0
4	A	38	0	29	1	0
4	B	38	0	29	2	0
5	A	8	0	6	0	0
5	B	8	0	6	0	0
6	A	6	0	8	0	0
6	B	6	0	8	0	0
7	A	1	0	0	0	0
8	A	193	0	0	2	0
8	B	153	0	0	1	0
All	All	6992	0	6403	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:CAS:SG	1:A:384:CAS:AS	2.49	1.30
1:B:384:CAS:AS	1:B:384:CAS:SG	2.54	1.25
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.63	0.80
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	1.66	0.75
1:A:95:PRO:HB3	1:A:108:PRO:HB2	1.68	0.74
4:B:800:6E5:H32	4:B:800:6E5:N28	2.11	0.65
1:A:384:CAS:CB	1:A:384:CAS:AS	3.05	0.65
2:B:500:HEM:HHC	2:B:500:HEM:HBB2	1.79	0.65
1:B:71:PRO:HG2	1:B:84:ASP:HB3	1.81	0.63
1:A:240:ARG:HD3	1:A:298:PRO:CB	2.30	0.60
1:B:236:ARG:NH1	1:B:242:ASP:OD1	2.35	0.60
2:A:500:HEM:CBB	2:A:500:HEM:HHC	2.32	0.58
1:B:99:ARG:HG2	1:B:100:ARG:HD2	1.85	0.58
1:B:384:CAS:AS	1:B:384:CAS:CB	3.12	0.57
1:A:378:LEU:HB2	8:A:2140:HOH:O	2.05	0.56
1:A:95:PRO:CB	1:A:108:PRO:HB2	2.37	0.55
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.87	0.55
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.37	0.54
1:A:102:LEU:HD11	1:B:71:PRO:HB3	1.90	0.53
1:B:384:CAS:CE2	1:B:384:CAS:SG	2.96	0.52
1:A:281:TRP:HB2	1:A:304:LEU:HD21	1.90	0.52
1:A:449:TRP:HA	3:A:600:H4B:N1	2.25	0.51
1:B:236:ARG:HH11	1:B:242:ASP:CG	2.14	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:800:6E5:C32	4:B:800:6E5:N28	2.73	0.50
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.51	0.50
1:B:167:LEU:HG	1:B:348:LEU:HD12	1.93	0.49
1:A:384:CAS:AS	1:A:384:CAS:HB2	2.73	0.48
1:A:145:SER:O	1:A:149:GLU:HG2	2.13	0.47
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.96	0.47
1:A:240:ARG:HD2	1:A:241:GLY:O	2.13	0.47
2:B:500:HEM:CMC	2:B:500:HEM:HBC2	2.45	0.46
1:B:370:CYS:HB3	1:B:378:LEU:HD22	1.96	0.46
1:A:158:GLU:OE2	1:A:166:HIS:HD2	2.00	0.45
1:A:423:HIS:HB2	1:B:392:THR:HB	2.00	0.44
1:A:392:THR:HB	1:B:423:HIS:HB2	2.00	0.44
1:A:279:HIS:HD2	8:A:2113:HOH:O	2.00	0.44
1:A:472:SER:HA	1:A:473:PRO:C	2.38	0.44
2:B:500:HEM:HHC	2:B:500:HEM:CBB	2.48	0.43
1:B:390:ARG:NH1	1:B:390:ARG:HB2	2.34	0.43
1:B:330:LEU:HB2	8:B:2103:HOH:O	2.18	0.43
1:B:236:ARG:HG3	1:B:349:GLU:O	2.19	0.42
1:A:126:LEU:HA	1:A:126:LEU:HD12	1.80	0.41
1:A:340:ASN:H	1:A:340:ASN:HD22	1.69	0.41
2:A:500:HEM:HBB2	2:A:500:HEM:CHC	2.43	0.41
1:A:338:VAL:HG23	4:A:800:6E5:H04	2.03	0.40
1:A:384:CAS:SG	1:A:384:CAS:CE2	3.09	0.40
1:A:70:PHE:HB3	1:A:84:ASP:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	401/443 (90%)	389 (97%)	11 (3%)	1 (0%)	52 43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	397/443 (90%)	386 (97%)	11 (3%)	0	100	100
All	All	798/886 (90%)	775 (97%)	22 (3%)	1 (0%)	56	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/375 (92%)	330 (96%)	14 (4%)	37	22
1	B	341/375 (91%)	329 (96%)	12 (4%)	43	29
All	All	685/750 (91%)	659 (96%)	26 (4%)	40	25

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	72	ARG
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	107	LEU
1	A	125	LEU
1	A	154	GLU
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	340	ASN
1	A	458	LEU
1	A	468	ASN
1	B	69	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	100	ARG
1	B	105	LEU
1	B	124	GLN
1	B	141	LYS
1	B	149	GLU
1	B	330	LEU
1	B	340	ASN
1	B	378	LEU
1	B	379	GLU
1	B	389	THR
1	B	458	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	HIS
1	A	191	GLN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	178	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	384	1	5,8,9	1.04	0	2,9,11	1.26	0
1	CAS	B	384	1	5,8,9	1.28	0	2,9,11	1.53	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	0/0/0/0
1	CAS	B	384	1	-	0/0/7/9	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	384	CAS	O-C-CA	-2.11	119.99	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	384	CAS	4	0
1	B	384	CAS	3	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	500	1	30,50,50	2.05	9 (30%)	24,82,82	2.36	11 (45%)
3	H4B	A	600	-	13,18,18	0.81	0	11,26,26	2.65	5 (45%)
4	6E5	A	800	-	40,42,42	1.32	5 (12%)	35,56,56	1.83	6 (17%)
5	ACT	A	860	-	1,3,3	1.51	0	0,3,3	0.00	-
5	ACT	A	861	-	1,3,3	2.32	1 (100%)	0,3,3	0.00	-
6	GOL	A	880	-	5,5,5	0.39	0	5,5,5	0.41	0
2	HEM	B	500	1	30,50,50	2.07	6 (20%)	24,82,82	2.38	10 (41%)
3	H4B	B	600	-	13,18,18	0.99	0	11,26,26	2.15	6 (54%)
4	6E5	B	800	-	40,42,42	1.29	5 (12%)	35,56,56	1.85	4 (11%)
5	ACT	B	860	-	1,3,3	1.98	0	0,3,3	0.00	-
5	ACT	B	861	-	1,3,3	1.47	0	0,3,3	0.00	-
6	GOL	B	880	-	5,5,5	0.23	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	0/10/54/54	0/0/8/8
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
4	6E5	A	800	-	-	0/19/36/36	0/5/5/5
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
5	ACT	A	861	-	-	0/0/0/0	0/0/0/0
6	GOL	A	880	-	-	0/4/4/4	0/0/0/0
2	HEM	B	500	1	-	0/10/54/54	0/0/8/8
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
4	6E5	B	800	-	-	0/19/36/36	0/5/5/5
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0
5	ACT	B	861	-	-	0/0/0/0	0/0/0/0
6	GOL	B	880	-	-	0/4/4/4	0/0/0/0

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C2D-C3D	-6.35	1.35	1.54
2	B	500	HEM	C2D-C3D	-6.28	1.35	1.54
2	B	500	HEM	C2C-C1C	-5.82	1.41	1.52
2	A	500	HEM	C2C-C1C	-4.51	1.44	1.52
2	B	500	HEM	C3B-C4B	-3.73	1.48	1.51
4	A	800	6E5	C06-N07	-3.45	1.32	1.39
4	B	800	6E5	C06-N07	-3.19	1.32	1.39
2	A	500	HEM	C3D-C4D	-3.13	1.47	1.51
2	A	500	HEM	C3B-C4B	-3.06	1.49	1.51
4	B	800	6E5	C11-N07	-2.78	1.36	1.41
4	A	800	6E5	C31-N27	-2.65	1.36	1.41
4	A	800	6E5	C26-N27	-2.37	1.34	1.39
2	B	500	HEM	C2B-C1B	-2.24	1.44	1.51
4	B	800	6E5	C31-N27	-2.04	1.37	1.41
2	A	500	HEM	C2B-C1B	-2.02	1.45	1.51
2	A	500	HEM	FE-NB	2.28	2.09	1.97
5	A	861	ACT	CH3-C	2.32	1.52	1.48
2	A	500	HEM	C1C-NC	2.34	1.38	1.36
2	B	500	HEM	FE-NB	2.65	2.11	1.97
2	A	500	HEM	FE-NC	2.89	2.07	1.95
2	B	500	HEM	FE-NC	3.03	2.07	1.95
4	A	800	6E5	C05-C06	3.15	1.50	1.45
4	B	800	6E5	C05-C06	3.21	1.50	1.45
2	A	500	HEM	C4C-NC	3.50	1.40	1.36
4	A	800	6E5	C25-C26	3.73	1.50	1.45
4	B	800	6E5	C25-C26	4.16	1.51	1.45

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	6E5	C23-C22-S21	-6.32	106.23	113.23
4	B	800	6E5	C23-C22-S21	-6.28	106.28	113.23
4	B	800	6E5	C03-C02-S01	-6.07	106.51	113.23
4	A	800	6E5	C03-C02-S01	-5.45	107.19	113.23
2	B	500	HEM	CBA-CAA-C2A	-3.55	106.17	112.53
2	B	500	HEM	C1D-CHD-C4C	-3.32	120.27	125.82
3	A	600	H4B	N3-C2-N1	-3.12	120.42	125.53
3	B	600	H4B	N3-C2-N1	-3.06	120.52	125.53
2	A	500	HEM	CBA-CAA-C2A	-3.05	107.06	112.53
2	B	500	HEM	C3C-CAC-CBC	-3.00	119.85	124.46
2	A	500	HEM	C1D-CHD-C4C	-2.74	121.23	125.82
2	A	500	HEM	C3C-CAC-CBC	-2.59	120.48	124.46
4	A	800	6E5	C31-N27-C26	-2.24	122.82	128.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	HEM	CMA-C3A-C4A	-2.04	124.98	128.36
2	A	500	HEM	C2D-C3D-C4D	2.08	105.03	101.50
3	B	600	H4B	C4-N3-C2	2.09	118.84	115.94
2	B	500	HEM	C2D-C3D-C4D	2.11	105.08	101.50
2	A	500	HEM	CMD-C2D-C3D	2.12	123.71	114.35
3	B	600	H4B	C4A-C8A-N8	2.13	120.94	118.43
4	A	800	6E5	C25-C26-N27	2.26	121.51	115.07
3	A	600	H4B	C4A-C8A-N8	2.32	121.17	118.43
4	A	800	6E5	O38-C37-C35	2.34	115.57	109.99
2	A	500	HEM	CHC-C4B-NB	2.38	130.25	124.52
3	A	600	H4B	C2-N1-C8A	2.53	120.23	114.54
2	B	500	HEM	CAA-CBA-CGA	2.54	117.41	112.75
2	B	500	HEM	CMD-C2D-C3D	2.74	126.45	114.35
4	B	800	6E5	C25-C26-N27	2.84	123.16	115.07
4	B	800	6E5	C3'-C4'-C5'	2.89	106.29	103.59
3	B	600	H4B	N2-C2-N3	3.03	122.22	117.20
3	B	600	H4B	C2-N1-C8A	3.27	121.89	114.54
4	A	800	6E5	C3'-C4'-C5'	3.41	106.77	103.59
3	B	600	H4B	C4-C4A-C8A	3.42	117.66	114.56
2	A	500	HEM	CMB-C2B-C3B	3.49	125.23	116.53
2	B	500	HEM	CMB-C2B-C3B	3.51	125.28	116.53
2	A	500	HEM	CMC-C2C-C3C	3.84	126.12	116.53
2	A	500	HEM	CAD-C3D-C4D	3.92	126.29	112.47
2	B	500	HEM	CMC-C2C-C3C	4.02	126.56	116.53
3	A	600	H4B	C4-N3-C2	4.45	122.11	115.94
2	B	500	HEM	CAD-C3D-C4D	4.55	128.51	112.47
2	B	500	HEM	CAD-C3D-C2D	4.59	126.41	113.22
2	A	500	HEM	CAD-C3D-C2D	5.29	128.43	113.22
3	A	600	H4B	C4-C4A-C8A	5.32	119.38	114.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	3	0
3	A	600	H4B	1	0
4	A	800	6E5	1	0
2	B	500	HEM	3	0
4	B	800	6E5	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	403/443 (90%)	0.56	41 (10%) 9 14	27, 39, 67, 92	0
1	B	401/443 (90%)	0.54	40 (9%) 9 15	28, 42, 74, 108	0
All	All	804/886 (90%)	0.55	81 (10%) 9 14	27, 41, 72, 108	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	GLN	8.2
1	B	121	PRO	5.8
1	A	67	GLY	5.5
1	A	160	ALA	5.5
1	A	69	LYS	5.5
1	A	122	ALA	5.3
1	A	121	PRO	5.2
1	B	122	ALA	5.0
1	A	123	GLU	5.0
1	A	259	GLN	5.0
1	A	99	ARG	4.9
1	B	142	ARG	4.9
1	A	239	GLY	4.7
1	B	98	PRO	4.6
1	A	91	GLN	4.5
1	B	99	ARG	4.5
1	B	108	PRO	4.4
1	A	98	PRO	4.2
1	A	238	PRO	4.2
1	A	124	GLN	4.2
1	A	97	THR	4.2
1	B	143	SER	4.1
1	B	146	GLN	4.0
1	A	338	VAL	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	260	ASP	3.9
1	B	94	GLY	3.9
1	B	69	LYS	3.8
1	A	260	ASP	3.8
1	B	123	GLU	3.7
1	A	159	VAL	3.6
1	A	93	ASP	3.5
1	A	94	GLY	3.3
1	A	126	LEU	3.3
1	A	261	GLY	3.3
1	B	141	LYS	3.2
1	B	239	GLY	3.2
1	B	93	ASP	3.1
1	B	91	GLN	3.1
1	B	96	CYS	3.1
1	A	125	LEU	3.1
1	B	338	VAL	3.0
1	B	223	ARG	3.0
1	B	258	GLN	3.0
1	A	155	VAL	3.0
1	B	92	GLN	2.9
1	B	261	GLY	2.9
1	A	230	ILE	2.8
1	A	92	GLN	2.8
1	B	97	THR	2.7
1	A	127	SER	2.6
1	B	70	PHE	2.6
1	A	355	PHE	2.6
1	B	147	ALA	2.6
1	A	257	ARG	2.6
1	A	153	GLN	2.6
1	A	414	LEU	2.6
1	B	414	LEU	2.5
1	B	124	GLN	2.4
1	A	157	ALA	2.4
1	A	337	ALA	2.4
1	A	156	GLU	2.4
1	A	161	SER	2.2
1	A	130	ARG	2.2
1	A	95	PRO	2.2
1	A	232	VAL	2.2
1	A	152	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	232	VAL	2.2
1	A	450	ILE	2.2
1	B	140	ILE	2.2
1	B	323	GLU	2.1
1	B	230	ILE	2.1
1	B	337	ALA	2.1
1	B	390	ARG	2.1
1	B	245	ILE	2.1
1	B	130	ARG	2.1
1	B	144	GLY	2.1
1	B	451	VAL	2.1
1	B	454	ILE	2.0
1	B	251	VAL	2.0
1	A	449	TRP	2.0
1	A	142	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CAS	A	384	9/10	0.98	0.08	-	37,38,58,63	0
1	CAS	B	384	9/10	0.97	0.09	-	47,49,65,67	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ACT	B	860	4/4	0.92	0.24	19.15	46,47,49,50	0
6	GOL	A	880	6/6	0.92	0.27	3.99	58,62,64,67	0
6	GOL	B	880	6/6	0.86	0.29	3.85	65,69,70,72	0
4	6E5	B	800	38/38	0.89	0.27	2.64	29,80,119,120	0
5	ACT	A	861	4/4	0.96	0.21	2.14	28,32,33,35	0
4	6E5	A	800	38/38	0.92	0.24	1.51	27,76,117,118	0
3	H4B	A	600	17/17	0.98	0.18	0.99	32,34,37,38	0
2	HEM	A	500	43/43	0.97	0.18	0.71	25,28,35,38	0
2	HEM	B	500	43/43	0.98	0.15	0.53	27,32,38,41	0
5	ACT	B	861	4/4	0.98	0.13	-0.06	38,38,39,40	0
3	H4B	B	600	17/17	0.97	0.14	-0.22	30,33,35,35	0
5	ACT	A	860	4/4	0.97	0.08	-0.89	44,47,48,48	0
7	ZN	A	1483	1/1	0.99	0.04	-3.88	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.