



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:48 AM GMT

PDB ID : 2UWN  
Title : CRYSTAL STRUCTURE OF HUMAN COMPLEMENT FACTOR H, SCR DOMAINS 6-8 (H402 RISK VARIANT), IN COMPLEX WITH LIGAND.  
Authors : Prosser, B.E.; Johnson, S.; Roversi, P.; Herbert, A.P.; Blaum, B.S.; Tyrrell, J.; Jowitt, T.A.; Clark, S.J.; Terelli, E.; Uhrin, D.; Barlow, P.N.; Sim, R.B.; Day, A.J.; Lea, S.M.  
Deposited on : 2007-03-22  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	<div> <div>9%</div> <div>71%</div> <div>25%</div> <div>• •</div> </div>

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 1755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN COMPLEMENT FACTOR H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	2	0
			1489	944	250	281	14			

- Molecule 2 is SUGAR (SUCROSE OCTASULFATE) (three-letter code: SCR) (formula:  $C_{12}H_{22}O_{35}S_8$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	1
			110	24	70	16		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	135	Total	O	0	0
			135	135		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: HUMAN COMPLEMENT FACTOR H



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.76Å 92.49Å 57.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.35 19.66 – 2.35	Depositor EDS
% Data completeness (in resolution range)	(Not available) (15.00-2.35) 97.9 (19.66-2.35)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.44 (at 2.35Å)	Xtriage
Refinement program	TNT 5.6.1	Depositor
R, $R_{free}$	0.214 , 0.246 0.227 , 0.274	Depositor DCC
$R_{free}$ test set	834 reflections (11.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.1	Xtriage
Anisotropy	0.513	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 60.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 8386 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	1755	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ACT, SO4, SCR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.64	3/1546 (0.2%)	0.64	4/2098 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	486	GLY	C-O	11.69	1.42	1.23
1	A	324	PRO	C-N	7.00	1.50	1.34
1	A	343	PRO	C-N	-6.44	1.19	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	343	PRO	O-C-N	-8.59	108.95	122.70
1	A	343	PRO	CA-C-N	7.20	133.04	117.20
1	A	486	GLY	C-N-CA	6.13	137.01	121.70
1	A	324	PRO	N-CA-C	-5.03	99.03	112.10

There are no chirality outliers.

All (3) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	323	LYS	Mainchain,Peptide
1	A	486	GLY	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1489	0	1379	47	3
2	A	110	0	42	17	3
3	A	4	0	3	1	0
4	A	2	0	0	1	0
5	A	15	0	0	0	0
6	A	135	0	0	15	2
All	All	1755	0	1424	61	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:462:GLU:HB3	1:A:463:SER:HA	1.30	1.09
1:A:390[A]:TYR:CD1	6:A:2035:HOH:O	2.29	0.84
1:A:453:ILE:HG21	1:A:460:ILE:HD11	1.60	0.83
1:A:485:ASP:H	1:A:486:GLY:HA2	1.45	0.79
1:A:344:TYR:HB2	6:A:2014:HOH:O	1.83	0.77
2:A:1507[A]:SCR:O94	2:A:1507[A]:SCR:H14	1.88	0.74
2:A:1507[A]:SCR:O53	2:A:1507[A]:SCR:O10	2.06	0.74
1:A:402:HIS:NE2	2:A:1507[A]:SCR:O64	2.19	0.73
1:A:335:LEU:HD21	1:A:372:ILE:HD11	1.72	0.71
1:A:453:ILE:CG2	1:A:460:ILE:HD11	2.21	0.70
1:A:323:LYS:HB3	1:A:324:PRO:CD	2.20	0.70
2:A:1507[A]:SCR:S12	2:A:1507[A]:SCR:H1	2.32	0.69
1:A:392:PRO:HB3	2:A:1507[A]:SCR:O72	1.94	0.68
1:A:390[A]:TYR:CE1	6:A:2035:HOH:O	2.46	0.67
1:A:413:ASP:HA	1:A:428:THR:HG22	1.76	0.67
1:A:462:GLU:CB	1:A:463:SER:HA	2.12	0.66

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:323:LYS:HB3	1:A:324:PRO:HD3	1.80	0.63
1:A:327:TYR:N	6:A:2005:HOH:O	1.95	0.62
1:A:462:GLU:HB3	1:A:463:SER:CA	2.18	0.61
2:A:1507[A]:SCR:O91	2:A:1507[A]:SCR:H1	2.00	0.61
1:A:347:VAL:HG13	1:A:351:LYS:HD2	1.80	0.61
1:A:485:ASP:N	1:A:486:GLY:HA2	2.15	0.61
2:A:1507[A]:SCR:O23	2:A:1507[A]:SCR:O93	2.21	0.59
1:A:322:LEU:C	1:A:323:LYS:HD3	2.24	0.58
1:A:342:ARG:NE	6:A:2015:HOH:O	2.28	0.58
1:A:323:LYS:N	1:A:323:LYS:HD3	2.19	0.57
2:A:1507[A]:SCR:O33	2:A:1507[A]:SCR:O2	2.24	0.56
2:A:1507[A]:SCR:O91	2:A:1507[A]:SCR:C1	2.52	0.56
2:A:1507[A]:SCR:O23	2:A:1507[A]:SCR:H1	2.04	0.56
1:A:411[B]:SER:OG	6:A:2052:HOH:O	2.18	0.55
1:A:345:PHE:HA	1:A:346:PRO:C	2.28	0.55
1:A:355:TYR:CE2	1:A:372:ILE:HG23	2.42	0.54
2:A:1507[A]:SCR:H74	2:A:1507[A]:SCR:H94	1.56	0.54
2:A:1507[B]:SCR:O23	2:A:1507[B]:SCR:H1	2.08	0.54
2:A:1507[A]:SCR:O91	2:A:1507[A]:SCR:O5	2.25	0.53
1:A:441:ARG:NH1	3:A:1509:ACT:OXT	2.40	0.53
1:A:455:ILE:HG23	4:A:1511:CL:CL	2.45	0.53
1:A:322:LEU:HD11	1:A:376:GLN:HB2	1.92	0.52
1:A:423:PRO:HA	1:A:424:LYS:CG	2.40	0.51
1:A:392:PRO:HB3	2:A:1507[B]:SCR:O53	2.10	0.51
1:A:423:PRO:HA	1:A:424:LYS:HG3	1.94	0.50
1:A:478:LYS:NZ	6:A:2112:HOH:O	2.13	0.50
1:A:389:CYS:HB3	1:A:436:TRP:CE2	2.47	0.49
1:A:375:THR:HG22	1:A:376:GLN:N	2.27	0.49
1:A:483:THR:HA	1:A:504:THR:O	2.13	0.49
1:A:424:LYS:HE2	6:A:2135:HOH:O	2.14	0.48
1:A:427:THR:OG1	6:A:2061:HOH:O	2.20	0.48
1:A:464:GLN:HB2	6:A:2093:HOH:O	2.14	0.47
1:A:342:ARG:CG	6:A:2015:HOH:O	2.62	0.47
1:A:396:ASN:ND2	6:A:2043:HOH:O	2.30	0.46
2:A:1507[A]:SCR:S12	2:A:1507[A]:SCR:O23	2.73	0.46
1:A:502:GLN:HB2	6:A:2086:HOH:O	2.14	0.46
1:A:327:TYR:HB2	6:A:2005:HOH:O	2.16	0.45
1:A:473:ALA:HB2	1:A:499:TRP:CH2	2.51	0.45
1:A:327:TYR:OH	1:A:338:GLU:HG2	2.18	0.44
1:A:439:THR:HA	1:A:440:PRO:HD3	1.91	0.43
1:A:365:SER:N	6:A:2029:HOH:O	2.39	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1507[A]:SCR:H15	2:A:1507[A]:SCR:O54	2.20	0.42
1:A:451:SER:HB3	1:A:465:TYR:HE1	1.84	0.41
2:A:1507[A]:SCR:C13	2:A:1507[A]:SCR:O5	2.68	0.41
1:A:493:THR:H	1:A:500:SER:HB2	1.86	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2017:HOH:O	6:A:2116:HOH:O[4_466]	0.35	1.85
6:A:2095:HOH:O	6:A:2105:HOH:O[8_556]	1.83	0.37
1:A:354:SER:OG	2:A:1507[A]:SCR:O24[6_455]	2.03	0.17
1:A:444:ARG:NH1	2:A:1507[A]:SCR:O94[3_555]	2.08	0.12
1:A:354:SER:OG	2:A:1507[B]:SCR:O24[6_455]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	187/187 (100%)	173 (92%)	13 (7%)	1 (0%)	34 39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	324	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/161 (101%)	152 (94%)	10 (6%)	23	26

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	323	LYS
1	A	338	GLU
1	A	347	VAL
1	A	390[A]	TYR
1	A	390[B]	TYR
1	A	411[A]	SER
1	A	411[B]	SER
1	A	476	GLN
1	A	485	ASP
1	A	488	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SCR	A	1507[A]	1	56,56,56	1.20	5 (8%)	79,92,92	2.31	26 (32%)
2	SCR	A	1507[B]	-	56,56,56	0.87	1 (1%)	79,92,92	1.91	17 (21%)
3	ACT	A	1509	-	1,3,3	1.72	0	0,3,3	0.00	-
5	SO4	A	1512	-	4,4,4	1.49	0	6,6,6	1.71	1 (16%)
5	SO4	A	1513	-	4,4,4	1.49	0	6,6,6	1.71	1 (16%)
5	SO4	A	1514	-	4,4,4	1.48	0	6,6,6	1.71	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SCR	A	1507[A]	1	-	0/48/88/88	0/2/2/2
2	SCR	A	1507[B]	-	-	0/48/88/88	0/2/2/2
3	ACT	A	1509	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1512	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1513	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1514	-	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1507[A]	SCR	O3-C3	-2.23	1.41	1.46
2	A	1507[B]	SCR	C14-C13	2.04	1.57	1.53
2	A	1507[A]	SCR	O33-S3	2.16	1.53	1.45
2	A	1507[A]	SCR	C4-C3	2.31	1.57	1.52
2	A	1507[A]	SCR	C11-C12	2.37	1.54	1.52
2	A	1507[A]	SCR	C12-C13	4.80	1.62	1.54

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1507[A]	SCR	C3-O3-S3	-8.24	103.07	118.77
2	A	1507[B]	SCR	C2-O2-S2	-5.93	107.48	118.77
2	A	1507[B]	SCR	C13-O91-S12	-5.72	108.48	118.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1507[A]	SCR	C14-O71-S13	-5.61	108.09	118.77
2	A	1507[B]	SCR	C3-O3-S3	-5.59	108.12	118.77
2	A	1507[B]	SCR	C14-O71-S13	-5.51	108.28	118.77
2	A	1507[B]	SCR	C4-O4-S4	-5.29	108.69	118.77
2	A	1507[A]	SCR	C2-O2-S2	-4.94	109.36	118.77
2	A	1507[A]	SCR	O5-C5-C6	-4.53	97.35	106.61
2	A	1507[B]	SCR	C12-O1-C1	-4.06	106.84	117.53
2	A	1507[A]	SCR	C4-O4-S4	-4.00	111.16	118.77
2	A	1507[A]	SCR	C11-C12-C13	-3.43	107.55	115.31
2	A	1507[A]	SCR	C3-C4-C5	-2.98	104.07	110.55
2	A	1507[A]	SCR	C13-O91-S12	-2.97	113.23	118.36
2	A	1507[B]	SCR	C6-C5-C4	-2.26	107.62	113.35
2	A	1507[A]	SCR	C12-O1-C1	-2.25	111.59	117.53
2	A	1507[B]	SCR	C1-O5-C5	-2.23	109.41	113.75
2	A	1507[A]	SCR	C4-C3-C2	-2.15	105.69	110.43
2	A	1507[B]	SCR	C1-C2-C3	-2.04	106.46	110.75
2	A	1507[A]	SCR	O91-C13-C12	2.46	117.68	110.98
2	A	1507[A]	SCR	C6-C5-C4	2.53	119.76	113.35
2	A	1507[A]	SCR	O73-S13-O72	2.64	123.97	112.46
2	A	1507[A]	SCR	O10-C15-C14	2.70	108.09	103.52
2	A	1507[B]	SCR	O73-S13-O72	2.76	124.49	112.46
2	A	1507[A]	SCR	O23-S2-O22	2.78	124.57	112.46
2	A	1507[A]	SCR	O2-C2-C3	2.79	114.66	108.48
2	A	1507[B]	SCR	O23-S2-O22	2.88	125.00	112.46
2	A	1507[B]	SCR	O93-S12-O92	2.88	125.02	112.46
2	A	1507[B]	SCR	O33-S3-O32	2.92	125.17	112.46
2	A	1507[B]	SCR	O53-S14-O52	2.96	125.37	112.46
2	A	1507[A]	SCR	O83-S11-O82	2.97	125.38	112.46
2	A	1507[B]	SCR	O63-S6-O62	2.97	125.41	112.46
2	A	1507[B]	SCR	O43-S4-O42	2.98	125.42	112.46
2	A	1507[A]	SCR	O93-S12-O92	2.99	125.47	112.46
2	A	1507[B]	SCR	O83-S11-O82	2.99	125.48	112.46
2	A	1507[A]	SCR	O43-S4-O42	3.10	125.94	112.46
2	A	1507[A]	SCR	O3-C3-C4	3.15	115.46	108.48
2	A	1507[A]	SCR	O63-S6-O62	3.30	126.85	112.46
2	A	1507[A]	SCR	O6-C6-C5	3.60	114.83	107.90
2	A	1507[A]	SCR	O5-C1-C2	3.67	116.88	109.47
5	A	1514	SO4	O4-S-O3	4.02	125.34	108.98
5	A	1512	SO4	O4-S-O3	4.03	125.35	108.98
5	A	1513	SO4	O4-S-O3	4.03	125.37	108.98
2	A	1507[A]	SCR	O1-C1-C2	4.14	115.28	107.57
2	A	1507[A]	SCR	O34-S3-O32	4.62	125.95	108.56

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1507[A]	SCR	O51-C16-C15	4.68	116.91	107.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1507[A]	SCR	15	2
2	A	1507[B]	SCR	2	1
3	A	1509	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	187/187 (100%)	0.49	16 (8%)	13 21	16, 36, 60, 70	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	485	ASP	4.9
1	A	344	TYR	4.7
1	A	484	ALA	4.6
1	A	479	LEU	4.0
1	A	390[A]	TYR	3.5
1	A	320	MET	3.3
1	A	465	TYR	3.1
1	A	496	LYS	2.9
1	A	322	LEU	2.8
1	A	323	LYS	2.7
1	A	339	ASN	2.5
1	A	460	ILE	2.4
1	A	343	PRO	2.2
1	A	462	GLU	2.2
1	A	463	SER	2.1
1	A	425	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SCR	A	1507[A]	55/55	0.75	0.32	1.99	70,99,110,110	55
2	SCR	A	1507[B]	55/55	0.75	0.32	1.98	51,73,81,81	55
4	CL	A	1511	1/1	0.95	0.20	1.00	58,58,58,58	0
4	CL	A	1510	1/1	0.98	0.13	-0.24	40,40,40,40	0
5	SO4	A	1514	5/5	0.80	0.25	-	102,103,103,103	0
5	SO4	A	1512	5/5	0.94	0.18	-	72,72,73,73	0
5	SO4	A	1513	5/5	0.95	0.17	-	91,91,91,91	0
3	ACT	A	1509	4/4	0.85	0.25	-	69,69,69,69	4

## 6.5 Other polymers

There are no such residues in this entry.