



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:44 PM GMT

PDB ID : 1V26
Title : Crystal structure of tt0168 from *Thermus thermophilus* HB8
Authors : Hisanaga, Y.; Ago, H.; Nakatsu, T.; Hamada, K.; Ida, K.; Kanda, H.; Yamamoto, M.; Hori, T.; Arii, Y.; Sugahara, M.; Kuramitsu, S.; Yokoyama, S.; Miyano, M.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2003-10-07
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	541	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>10%</div> </div> </div>
1	B	541	<div> <div>7%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MYR	B	2001	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called long-chain-fatty-acid-CoA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	489	Total	C	N	O	S	0	0	0
			3802	2436	661	695	10			
1	B	510	Total	C	N	O	S	0	0	0
			3964	2536	688	730	10			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is MYRISTIC ACID (three-letter code: MYR) (formula: C₁₄H₂₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			15	14	1		
3	B	1	Total	C	O	0	0
			15	14	1		

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	251	Total	O	0	0
			251	251		
5	B	221	Total	O	0	0
			221	221		

- Molecule 1: long-chain-fatty-acid-CoA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.51Å 101.38Å 176.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.10 – 2.50 46.35 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.10-2.50) 99.7 (46.35-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.66 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.204 , 0.240 0.203 , 0.237	Depositor DCC
R_{free} test set	4116 reflections (11.17%)	DCC
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40998 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8316	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, MG, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/3890	0.63	8/5285 (0.2%)
1	B	0.29	0/4056	0.62	6/5511 (0.1%)
All	All	0.29	0/7946	0.62	14/10796 (0.1%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	473	TRP	C-N-CA	-6.90	104.45	121.70
1	A	473	TRP	C-N-CA	-6.55	105.31	121.70
1	B	432	ASP	CB-CG-OD2	5.94	123.65	118.30
1	A	432	ASP	CB-CG-OD2	5.92	123.62	118.30
1	A	126	ASP	CB-CG-OD2	5.50	123.25	118.30
1	B	15	ASP	CB-CG-OD2	5.42	123.17	118.30
1	B	126	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	212	ASP	CB-CG-OD2	5.33	123.10	118.30
1	A	120	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	449	ASP	CB-CG-OD2	5.23	123.01	118.30
1	B	120	ASP	CB-CG-OD2	5.19	122.97	118.30
1	A	15	ASP	CB-CG-OD2	5.06	122.86	118.30
1	A	255	ASP	CB-CG-OD2	5.05	122.84	118.30
1	B	509	ASP	CB-CG-OD2	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3802	0	3814	28	0
1	B	3964	0	3963	32	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	15	0	27	3	0
3	B	15	0	27	2	0
4	A	23	0	12	0	0
4	B	23	0	12	0	0
5	A	251	0	0	1	0
5	B	221	0	0	0	0
All	All	8316	0	7855	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:471:PRO:HD3	1:B:540:GLY:HA2	1.24	1.14
1:B:438:ILE:HG21	1:B:477:PRO:HD3	1.48	0.96
1:B:330:SER:N	1:B:331:PRO:HA	2.01	0.76
1:B:225:VAL:HG11	1:B:277:VAL:HG11	1.67	0.75
1:B:471:PRO:HD3	1:B:540:GLY:CA	2.12	0.74
1:A:330:SER:N	1:A:331:PRO:HA	2.04	0.72
1:A:310:ILE:O	1:A:314:GLU:HG2	1.92	0.70
1:B:372:GLU:HB2	1:B:374:ARG:HD3	1.79	0.63
1:A:308:SER:O	1:A:312:ARG:HB2	2.00	0.61
1:B:19:ASN:HD22	1:B:21:TRP:H	1.47	0.61
1:B:275:PRO:HB2	1:B:474:GLN:NE2	2.16	0.61
1:A:474:GLN:HG3	5:A:3091:HOH:O	2.01	0.59
1:A:446:SER:HB3	1:A:449:ASP:HB2	1.85	0.58
1:B:305:ALA:H	1:B:322:GLN:NE2	2.01	0.58
1:B:92:VAL:HB	1:B:93:PRO:HD3	1.85	0.57
1:B:329:THR:C	1:B:331:PRO:HA	2.26	0.56
1:B:335:GLN:HE21	1:B:361:ILE:HG21	1.70	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:HIS:HE1	1:B:126:ASP:OD1	1.89	0.56
1:B:471:PRO:CD	1:B:540:GLY:HA2	2.17	0.54
1:A:19:ASN:HD22	1:A:21:TRP:H	1.56	0.54
1:B:128:ASN:HD22	1:B:128:ASN:H	1.55	0.52
1:A:264:ASP:HB3	1:A:293:LYS:HE3	1.91	0.52
1:B:293:LYS:HG3	1:B:294:THR:HG23	1.92	0.52
1:B:438:ILE:HD13	1:B:477:PRO:HG3	1.91	0.51
1:A:92:VAL:HB	1:A:93:PRO:HD3	1.91	0.51
1:A:329:THR:C	1:A:331:PRO:HA	2.32	0.50
1:A:419:ILE:HD11	1:A:434:LEU:HG	1.94	0.49
1:A:438:ILE:HD13	1:A:477:PRO:HB3	1.94	0.49
1:B:305:ALA:H	1:B:322:GLN:HE22	1.61	0.48
1:B:275:PRO:HB2	1:B:474:GLN:HE22	1.79	0.48
1:B:125:PHE:CZ	1:B:148:VAL:HG22	2.48	0.47
1:B:214:THR:HG21	3:B:2001:MYR:H111	1.96	0.47
1:A:228:MET:HA	1:A:233:ALA:HB2	1.97	0.46
1:A:82:HIS:HE1	1:A:126:ASP:OD1	1.98	0.46
1:A:128:ASN:H	1:A:128:ASN:HD22	1.63	0.46
1:B:477:PRO:HB2	1:B:508:PRO:HA	1.98	0.46
1:A:23:PHE:CE1	1:A:242:LEU:HD13	2.52	0.45
1:A:477:PRO:HG2	1:A:508:PRO:HA	1.99	0.45
1:B:234:TRP:CD2	3:B:2001:MYR:H62	2.51	0.45
1:A:305:ALA:H	1:A:322:GLN:NE2	2.14	0.45
1:A:127:PRO:HG3	1:A:150:ASP:HB2	1.99	0.45
1:A:323:GLY:HA3	3:A:1001:MYR:H72	1.99	0.44
1:B:419:ILE:HB	1:B:432:ASP:HB3	1.99	0.44
1:B:372:GLU:HG3	1:B:374:ARG:NH1	2.33	0.44
1:B:128:ASN:ND2	1:B:128:ASN:H	2.16	0.43
1:A:234:TRP:CE2	3:A:1001:MYR:H51	2.53	0.43
1:A:225:VAL:HG11	1:A:277:VAL:HG11	1.99	0.43
1:B:498:LEU:HD21	1:B:507:LEU:HD11	2.00	0.43
1:A:77:THR:HA	1:A:124:LEU:O	2.18	0.43
1:A:370:ASP:OD2	1:A:374:ARG:HB2	2.19	0.42
1:B:77:THR:HA	1:B:124:LEU:O	2.20	0.42
1:A:128:ASN:H	1:A:128:ASN:ND2	2.18	0.41
1:A:153:ALA:HA	1:A:154:PRO:HD3	1.91	0.41
1:A:305:ALA:H	1:A:322:GLN:HE22	1.69	0.41
1:B:225:VAL:HG13	1:B:254:LEU:HD22	2.03	0.40
1:A:299:VAL:HG11	3:A:1001:MYR:H101	2.03	0.40
1:B:419:ILE:HD11	1:B:434:LEU:HG	2.04	0.40
1:B:185:THR:HG22	1:B:191:PRO:HG3	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:103:ASN:HA	1:B:104:PRO:HD3	1.89	0.40
1:A:225:VAL:HG13	1:A:254:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/541 (89%)	469 (98%)	11 (2%)	1 (0%)	52	75
1	B	502/541 (93%)	489 (97%)	11 (2%)	2 (0%)	39	61
All	All	983/1082 (91%)	958 (98%)	22 (2%)	3 (0%)	46	68

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	330	SER
1	B	330	SER
1	B	518	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	395/437 (90%)	380 (96%)	15 (4%)	40	67

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	411/437 (94%)	396 (96%)	15 (4%)	42	69
All	All	806/874 (92%)	776 (96%)	30 (4%)	41	68

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	109	LYS
1	A	172	ARG
1	A	185	THR
1	A	187	THR
1	A	231	VAL
1	A	235	CYS
1	A	236	LEU
1	A	283	ASP
1	A	291	ARG
1	A	336	ASN
1	A	422	TRP
1	A	435	LYS
1	A	449	ASP
1	A	526	LEU
1	B	55	GLN
1	B	74	ARG
1	B	109	LYS
1	B	128	ASN
1	B	139	GLU
1	B	155	GLU
1	B	172	ARG
1	B	223	LEU
1	B	231	VAL
1	B	235	CYS
1	B	236	LEU
1	B	303	SER
1	B	422	TRP
1	B	435	LYS
1	B	448	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	55	GLN
1	A	82	HIS
1	A	128	ASN
1	A	144	GLN
1	A	322	GLN
1	A	335	GLN
1	A	336	ASN
1	B	19	ASN
1	B	55	GLN
1	B	82	HIS
1	B	128	ASN
1	B	145	HIS
1	B	232	ASN
1	B	322	GLN
1	B	335	GLN
1	B	336	ASN
1	B	474	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MYR	A	1001	4	14,14,15	0.30	0	12,13,15	0.76	0
4	AMP	A	1002	3,2	20,25,25	1.23	2 (10%)	22,38,38	2.28	3 (13%)
3	MYR	B	2001	4	14,14,15	0.30	0	12,13,15	0.70	0
4	AMP	B	2002	3,2	20,25,25	1.24	2 (10%)	22,38,38	2.41	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MYR	A	1001	4	-	0/11/12/13	0/0/0/0
4	AMP	A	1002	3,2	-	0/6/26/26	0/3/3/3
3	MYR	B	2001	4	-	0/11/12/13	0/0/0/0
4	AMP	B	2002	3,2	-	0/6/26/26	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1002	AMP	P-O1P	2.43	1.59	1.51
4	B	2002	AMP	P-O1P	2.47	1.59	1.51
4	A	1002	AMP	O4'-C1'	4.12	1.46	1.41
4	B	2002	AMP	O4'-C1'	4.20	1.46	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2002	AMP	N3-C2-N1	-10.00	121.23	128.89
4	A	1002	AMP	N3-C2-N1	-9.45	121.66	128.89
4	A	1002	AMP	C2'-C1'-N9	-2.72	110.14	114.29
4	B	2002	AMP	C2'-C1'-N9	-2.61	110.30	114.29
4	B	2002	AMP	C4-C5-N7	-2.39	107.28	109.48
4	A	1002	AMP	C4-C5-N7	-2.37	107.30	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	MYR	3	0
3	B	2001	MYR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	489/541 (90%)	0.14	32 (6%)	22 25	10, 21, 70, 100	0
1	B	510/541 (94%)	0.11	39 (7%)	17 18	11, 21, 71, 102	0
All	All	999/1082 (92%)	0.13	71 (7%)	19 21	10, 21, 70, 102	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	493	LEU	8.8
1	A	481	VAL	8.5
1	B	514	ALA	6.2
1	B	541	ALA	5.9
1	B	490	PRO	5.5
1	A	526	LEU	5.3
1	A	478	LEU	5.2
1	B	515	GLU	5.1
1	A	502	PHE	4.9
1	B	463	ALA	4.9
1	B	497	LEU	4.8
1	B	516	GLU	4.7
1	B	517	ILE	4.4
1	B	528	ARG	4.3
1	A	531	ARG	4.2
1	A	468	ILE	4.1
1	A	501	GLY	4.1
1	B	533	GLN	4.0
1	B	8	ALA	3.9
1	B	480	VAL	3.9
1	B	464	ALA	3.9
1	A	528	ARG	3.7
1	A	464	ALA	3.6
1	B	489	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	512	VAL	3.5
1	A	510	ALA	3.5
1	A	511	TYR	3.4
1	B	452	ASN	3.4
1	A	480	VAL	3.4
1	A	507	LEU	3.4
1	B	513	PHE	3.3
1	A	8	ALA	3.2
1	B	492	GLU	3.2
1	B	534	TYR	3.2
1	B	495	GLU	3.1
1	A	529	ALA	3.0
1	A	479	ALA	3.0
1	A	539	GLY	2.9
1	B	519	ARG	2.8
1	A	533	GLN	2.8
1	B	371	GLU	2.7
1	A	530	LEU	2.7
1	B	498	LEU	2.7
1	A	532	GLU	2.7
1	B	518	PRO	2.7
1	B	504	LYS	2.6
1	A	534	TYR	2.5
1	B	510	ALA	2.5
1	A	527	LYS	2.5
1	A	155	GLU	2.5
1	A	509	ASP	2.4
1	B	502	PHE	2.4
1	B	372	GLU	2.4
1	B	479	ALA	2.4
1	B	348	GLU	2.3
1	B	494	ASN	2.3
1	B	450	LEU	2.3
1	A	449	ASP	2.3
1	A	448	VAL	2.3
1	A	504	LYS	2.2
1	A	452	ASN	2.2
1	A	465	VAL	2.2
1	A	402	GLU	2.2
1	B	540	GLY	2.1
1	B	500	ALA	2.1
1	A	405	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	530	LEU	2.1
1	B	152	LYS	2.1
1	B	155	GLU	2.0
1	A	469	PRO	2.0
1	B	478	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MYR	B	2001	15/16	0.88	0.26	5.00	17,18,20,20	0
3	MYR	A	1001	15/16	0.87	0.19	1.95	6,7,9,9	0
2	MG	A	3001	1/1	0.96	0.12	-0.84	35,35,35,35	0
4	AMP	A	1002	23/23	0.95	0.11	-0.89	7,13,13,14	0
4	AMP	B	2002	23/23	0.96	0.12	-1.23	16,17,19,20	0
2	MG	B	3002	1/1	0.97	0.09	-3.48	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.