



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:55 AM GMT

PDB ID : 2VA0  
Title : DIFFERENTIAL REGULATION OF THE XYLAN DEGRADING APPARATUS OF CELLVIBRIO JAPONICUS BY A NOVEL TWO COMPONENT SYSTEM  
Authors : Murray, J.W.; Emami, K.; Topakas, E.; Nagy, T.; Henshaw, J.; Jackson, K.A.; Nelson, K.E.; Mongodin, E.F.; Lewis, R.J.; Gilbert, H.J.  
Deposited on : 2007-08-28  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	131	<div> <div>7%</div> <div>57%</div> <div>16%</div> <div>•</div> <div>24%</div> </div>
1	B	131	<div> <div>2%</div> <div>73%</div> <div>13%</div> <div>• •</div> <div>9%</div> </div>
1	C	131	<div> <div>•</div> <div>64%</div> <div>10%</div> <div>•</div> <div>24%</div> </div>
1	D	131	<div> <div>5%</div> <div>56%</div> <div>15%</div> <div>• •</div> <div>25%</div> </div>
1	E	131	<div> <div>5%</div> <div>73%</div> <div>14%</div> <div>• •</div> <div>8%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	131	<div><div></div><div>5%</div><div>57%</div><div>16%</div><div>•</div><div>24%</div></div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	S	0	0	0
			800	504	155	138	3			
1	B	119	Total	C	N	O	S	0	0	0
			973	612	182	175	4			
1	C	99	Total	C	N	O	S	0	0	0
			803	505	155	140	3			
1	D	98	Total	C	N	O	S	0	0	0
			795	501	154	137	3			
1	E	120	Total	C	N	O	S	0	0	0
			980	616	186	174	4			
1	F	99	Total	C	N	O	S	0	0	0
			800	504	155	138	3			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	MET	-	EXPRESSION TAG	UNP B3PFT7
A	166	LEU	-	EXPRESSION TAG	UNP B3PFT7
A	167	GLU	-	EXPRESSION TAG	UNP B3PFT7
A	168	HIS	-	EXPRESSION TAG	UNP B3PFT7
A	169	HIS	-	EXPRESSION TAG	UNP B3PFT7
A	170	HIS	-	EXPRESSION TAG	UNP B3PFT7
A	171	HIS	-	EXPRESSION TAG	UNP B3PFT7
A	172	HIS	-	EXPRESSION TAG	UNP B3PFT7
A	173	HIS	-	EXPRESSION TAG	UNP B3PFT7
B	43	MET	-	EXPRESSION TAG	UNP B3PFT7
B	166	LEU	-	EXPRESSION TAG	UNP B3PFT7
B	167	GLU	-	EXPRESSION TAG	UNP B3PFT7
B	168	HIS	-	EXPRESSION TAG	UNP B3PFT7
B	169	HIS	-	EXPRESSION TAG	UNP B3PFT7
B	170	HIS	-	EXPRESSION TAG	UNP B3PFT7
B	171	HIS	-	EXPRESSION TAG	UNP B3PFT7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	172	HIS	-	EXPRESSION TAG	UNP B3PFT7
B	173	HIS	-	EXPRESSION TAG	UNP B3PFT7
C	43	MET	-	EXPRESSION TAG	UNP B3PFT7
C	166	LEU	-	EXPRESSION TAG	UNP B3PFT7
C	167	GLU	-	EXPRESSION TAG	UNP B3PFT7
C	168	HIS	-	EXPRESSION TAG	UNP B3PFT7
C	169	HIS	-	EXPRESSION TAG	UNP B3PFT7
C	170	HIS	-	EXPRESSION TAG	UNP B3PFT7
C	171	HIS	-	EXPRESSION TAG	UNP B3PFT7
C	172	HIS	-	EXPRESSION TAG	UNP B3PFT7
C	173	HIS	-	EXPRESSION TAG	UNP B3PFT7
D	43	MET	-	EXPRESSION TAG	UNP B3PFT7
D	166	LEU	-	EXPRESSION TAG	UNP B3PFT7
D	167	GLU	-	EXPRESSION TAG	UNP B3PFT7
D	168	HIS	-	EXPRESSION TAG	UNP B3PFT7
D	169	HIS	-	EXPRESSION TAG	UNP B3PFT7
D	170	HIS	-	EXPRESSION TAG	UNP B3PFT7
D	171	HIS	-	EXPRESSION TAG	UNP B3PFT7
D	172	HIS	-	EXPRESSION TAG	UNP B3PFT7
D	173	HIS	-	EXPRESSION TAG	UNP B3PFT7
E	43	MET	-	EXPRESSION TAG	UNP B3PFT7
E	166	LEU	-	EXPRESSION TAG	UNP B3PFT7
E	167	GLU	-	EXPRESSION TAG	UNP B3PFT7
E	168	HIS	-	EXPRESSION TAG	UNP B3PFT7
E	169	HIS	-	EXPRESSION TAG	UNP B3PFT7
E	170	HIS	-	EXPRESSION TAG	UNP B3PFT7
E	171	HIS	-	EXPRESSION TAG	UNP B3PFT7
E	172	HIS	-	EXPRESSION TAG	UNP B3PFT7
E	173	HIS	-	EXPRESSION TAG	UNP B3PFT7
F	43	MET	-	EXPRESSION TAG	UNP B3PFT7
F	166	LEU	-	EXPRESSION TAG	UNP B3PFT7
F	167	GLU	-	EXPRESSION TAG	UNP B3PFT7
F	168	HIS	-	EXPRESSION TAG	UNP B3PFT7
F	169	HIS	-	EXPRESSION TAG	UNP B3PFT7
F	170	HIS	-	EXPRESSION TAG	UNP B3PFT7
F	171	HIS	-	EXPRESSION TAG	UNP B3PFT7
F	172	HIS	-	EXPRESSION TAG	UNP B3PFT7
F	173	HIS	-	EXPRESSION TAG	UNP B3PFT7

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		
3	F	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	10	Total 10	O 10	0	0
4	B	28	Total 28	O 28	0	0
4	C	23	Total 23	O 23	0	0
4	D	23	Total 23	O 23	0	0
4	E	11	Total 11	O 11	0	0
4	F	7	Total 7	O 7	0	0
4	D	1	Total 1	O 1	0	0
4	B	1	Total 1	O 1	0	0
4	F	1	Total 1	O 1	0	0
4	A	1	Total 1	O 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN




- Molecule 1: ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN



- Molecule 1: ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN



- Molecule 1: ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN





GLY  
PRO  
ASP  
ARG  
ASP  
ASP  
ILE  
ILE  
TRP  
GLU  
MET  
PHE  
LEU  
LEU  
ASN  
LEU  
GLU  
HIS  
HIS  
HIS  
HIS  
HIS

- Molecule 1: ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN



MET  
TYR  
LEU  
PRO  
ASP  
ASP  
S49  
R65  
L70  
T71  
Q72  
D73  
V85  
R90  
D94  
L100  
T112  
H115  
D118  
Q121  
K126  
D136  
G137  
N138  
T144  
G148  
R149  
D150  
E151  
G152  
P153  
D154  
E160  
M161  
F162  
L163  
E164  
N165  
H168  
HIS  
HIS  
HIS  
HIS

- Molecule 1: ABFS ARABINOFURANOSIDASE TWO COMPONENT SYSTEM SENSOR PROTEIN



MET  
TYR  
LEU  
PRO  
ASP  
ASP  
S49  
P50  
R53  
L54  
M61  
R65  
Q72  
D73  
V80  
R83  
F84  
V85  
D89  
R90  
N91  
M105  
V106  
L107  
A108  
P109  
T112  
A113  
K114  
H115  
V120  
R123  
M124  
R125  
K126  
L127  
D136  
G137  
N138  
T144  
I145  
S146  
A147  
GLY  
ARG  
ASP  
GLU

GLY  
PRO  
ASP  
ARG  
ASP  
ASP  
ILE  
ILE  
TRP  
GLU  
MET  
PHE  
LEU  
LEU  
ASN  
LEU  
GLU  
HIS  
HIS  
HIS  
HIS  
HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.77Å 80.01Å 101.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.04 – 2.60 37.04 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.04-2.60) 99.8 (37.04-2.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.13 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.199 , 0.259 0.200 , 0.255	Depositor DCC
$R_{free}$ test set	1430 reflections (5.45%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.9	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 43.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 27692 reflections (0.007%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5290	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	2/815 (0.2%)	0.96	2/1104 (0.2%)
1	B	0.91	1/992 (0.1%)	1.06	7/1343 (0.5%)
1	C	0.89	1/818 (0.1%)	1.07	5/1108 (0.5%)
1	D	0.89	1/810 (0.1%)	1.01	6/1097 (0.5%)
1	E	0.85	0/1000	1.01	4/1354 (0.3%)
1	F	0.78	0/815	0.94	3/1104 (0.3%)
All	All	0.86	5/5250 (0.1%)	1.01	27/7110 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	2
All	All	0	3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	136	ASP	CB-CG	-6.55	1.38	1.51
1	B	85	VAL	CB-CG2	-6.32	1.39	1.52
1	C	73	ASP	CB-CG	-6.22	1.38	1.51
1	A	136	ASP	CB-CG	-5.49	1.40	1.51
1	A	73	ASP	CB-CG	-5.43	1.40	1.51

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	73	ASP	CB-CG-OD1	-8.75	110.42	118.30
1	D	136	ASP	CB-CG-OD1	-8.59	110.57	118.30
1	C	65	ARG	NE-CZ-NH2	-7.88	116.36	120.30
1	C	65	ARG	NE-CZ-NH1	7.50	124.05	120.30
1	B	58	MET	CG-SD-CE	-7.31	88.51	100.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	47	ASP	Peptide
1	E	149	ARG	Peptide
1	E	150	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	800	0	834	18	0
1	B	973	0	981	23	0
1	C	803	0	833	12	0
1	D	795	0	829	18	0
1	E	980	0	994	13	0
1	F	800	0	834	18	0
2	A	5	0	0	1	0
2	B	5	0	0	1	0
2	C	5	0	0	1	0
2	D	5	0	0	1	0
2	E	5	0	0	1	0
2	F	5	0	0	1	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	F	1	0	0	0	0
4	A	11	0	0	1	0
4	B	29	0	0	5	0
4	C	23	0	0	0	0
4	D	24	0	0	1	0
4	E	11	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	8	0	0	1	0
All	All	5290	0	5305	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100:LEU:CD1	1:B:105:MET:CE	2.28	1.10
1:F:112:THR:HG22	1:F:115:HIS:H	1.03	1.08
1:B:100:LEU:HD13	1:B:105:MET:CE	1.85	1.07
1:D:112:THR:HG22	1:D:115:HIS:H	1.26	1.00
1:D:112:THR:HG21	2:D:1147:PO4:O1	1.61	1.00

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/131 (74%)	95 (98%)	2 (2%)	0	100	100
1	B	115/131 (88%)	111 (96%)	4 (4%)	0	100	100
1	C	97/131 (74%)	95 (98%)	2 (2%)	0	100	100
1	D	96/131 (73%)	95 (99%)	1 (1%)	0	100	100
1	E	118/131 (90%)	114 (97%)	2 (2%)	2 (2%)	11	22
1	F	97/131 (74%)	95 (98%)	1 (1%)	1 (1%)	19	39
All	All	620/786 (79%)	605 (98%)	12 (2%)	3 (0%)	34	60

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	150	ASP
1	E	154	ASP
1	F	83	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/119 (75%)	85 (96%)	4 (4%)	34	62
1	B	108/119 (91%)	101 (94%)	7 (6%)	21	42
1	C	90/119 (76%)	88 (98%)	2 (2%)	60	83
1	D	89/119 (75%)	86 (97%)	3 (3%)	44	72
1	E	108/119 (91%)	100 (93%)	8 (7%)	17	34
1	F	89/119 (75%)	83 (93%)	6 (7%)	20	40
All	All	573/714 (80%)	543 (95%)	30 (5%)	29	54

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	85	VAL
1	E	72	GLN
1	F	112	THR
1	D	136	ASP
1	E	73	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN
1	A	121	GLN
1	B	124	ASN
1	E	72	GLN
1	E	121	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	1148	-	4,4,4	0.52	0	6,6,6	0.26	0
2	PO4	B	1169	-	4,4,4	0.39	0	6,6,6	0.29	0
2	PO4	C	1147	-	4,4,4	0.57	0	6,6,6	0.29	0
2	PO4	D	1147	-	4,4,4	0.30	0	6,6,6	0.34	0
2	PO4	E	1169	-	4,4,4	0.58	0	6,6,6	0.29	0
2	PO4	F	1148	-	4,4,4	0.40	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1148	-	-	0/0/0/0	0/0/0/0
2	PO4	B	1169	-	-	0/0/0/0	0/0/0/0
2	PO4	C	1147	-	-	0/0/0/0	0/0/0/0
2	PO4	D	1147	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	E	1169	-	-	0/0/0/0	0/0/0/0
2	PO4	F	1148	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1148	PO4	1	0
2	B	1169	PO4	1	0
2	C	1147	PO4	1	0
2	D	1147	PO4	1	0
2	E	1169	PO4	1	0
2	F	1148	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	99/131 (75%)	0.49	9 (9%) 11 7	3, 9, 17, 22	0
1	B	119/131 (90%)	0.19	2 (1%) 73 68	3, 9, 24, 35	0
1	C	99/131 (75%)	0.01	1 (1%) 84 81	3, 9, 19, 22	0
1	D	98/131 (74%)	0.34	6 (6%) 25 18	3, 9, 17, 22	0
1	E	120/131 (91%)	0.29	7 (5%) 26 20	3, 11, 36, 49	0
1	F	99/131 (75%)	0.30	6 (6%) 25 18	3, 9, 18, 21	0
All	All	634/786 (80%)	0.27	31 (4%) 33 26	3, 9, 22, 49	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	80	TRP	6.5
1	E	150	ASP	5.0
1	A	80	TRP	4.7
1	F	80	TRP	4.5
1	E	149	ARG	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	A	1149	1/1	0.98	0.20	1.67	7,7,7,7	0
3	CL	C	1148	1/1	0.99	0.16	-0.61	3,3,3,3	0
2	PO4	D	1147	5/5	0.97	0.17	-1.73	7,9,13,15	0
3	CL	F	1149	1/1	0.99	0.13	-2.99	8,8,8,8	0
2	PO4	B	1169	5/5	0.98	0.11	-3.33	14,18,21,23	0
2	PO4	A	1148	5/5	0.98	0.17	-	29,29,30,32	0
2	PO4	E	1169	5/5	0.98	0.19	-	16,17,20,20	0
2	PO4	F	1148	5/5	0.96	0.17	-	24,27,28,28	0
2	PO4	C	1147	5/5	0.96	0.15	-	12,12,17,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.