



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:51 PM GMT

PDB ID : 1VIQ
Title : Crystal structure of putative ADP ribose pyrophosphatase
Authors : Structural GenomiX
Deposited on : 2003-12-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	<div> <div>71% 20% 7%</div> </div>
1	B	220	<div> <div>63% 23% 5% 9%</div> </div>
1	C	220	<div> <div>65% 18% 5% 12%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribose pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	0	0
			1597	1010	281	302	4			
1	B	201	Total	C	N	O	S	0	1	0
			1595	1006	285	300	4			
1	C	194	Total	C	N	O	S	0	0	0
			1533	969	270	290	4			

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	UNP Q93K97
A	0	SER	-	cloning artifact	UNP Q93K97
A	28	SER	PHE	variant	UNP Q93K97
A	209	GLU	-	cloning artifact	UNP Q93K97
A	210	GLY	-	cloning artifact	UNP Q93K97
A	211	GLY	-	cloning artifact	UNP Q93K97
A	212	SER	-	cloning artifact	UNP Q93K97
A	213	HIS	-	cloning artifact	UNP Q93K97
A	214	HIS	-	cloning artifact	UNP Q93K97
A	215	HIS	-	cloning artifact	UNP Q93K97
A	216	HIS	-	cloning artifact	UNP Q93K97
A	217	HIS	-	cloning artifact	UNP Q93K97
A	218	HIS	-	cloning artifact	UNP Q93K97
B	-1	MET	-	cloning artifact	UNP Q93K97
B	0	SER	-	cloning artifact	UNP Q93K97
B	28	SER	PHE	variant	UNP Q93K97
B	209	GLU	-	cloning artifact	UNP Q93K97
B	210	GLY	-	cloning artifact	UNP Q93K97
B	211	GLY	-	cloning artifact	UNP Q93K97
B	212	SER	-	cloning artifact	UNP Q93K97
B	213	HIS	-	cloning artifact	UNP Q93K97
B	214	HIS	-	cloning artifact	UNP Q93K97
B	215	HIS	-	cloning artifact	UNP Q93K97

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	216	HIS	-	cloning artifact	UNP Q93K97
B	217	HIS	-	cloning artifact	UNP Q93K97
B	218	HIS	-	cloning artifact	UNP Q93K97
C	-1	MET	-	cloning artifact	UNP Q93K97
C	0	SER	-	cloning artifact	UNP Q93K97
C	28	SER	PHE	variant	UNP Q93K97
C	209	GLU	-	cloning artifact	UNP Q93K97
C	210	GLY	-	cloning artifact	UNP Q93K97
C	211	GLY	-	cloning artifact	UNP Q93K97
C	212	SER	-	cloning artifact	UNP Q93K97
C	213	HIS	-	cloning artifact	UNP Q93K97
C	214	HIS	-	cloning artifact	UNP Q93K97
C	215	HIS	-	cloning artifact	UNP Q93K97
C	216	HIS	-	cloning artifact	UNP Q93K97
C	217	HIS	-	cloning artifact	UNP Q93K97
C	218	HIS	-	cloning artifact	UNP Q93K97


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	80	Total O 80 80	0	0
2	B	91	Total O 91 91	0	0
2	C	80	Total O 80 80	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-ribose pyrophosphatase



- Molecule 1: ADP-ribose pyrophosphatase



- Molecule 1: ADP-ribose pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	98.18Å 98.18Å 155.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	51.98 – 2.40 51.98 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (51.98-2.40) 99.9 (51.98-2.40)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.39Å)	Xtriage
Refinement program	REFMAC 4	Depositor
R, R_{free}	0.244 , 0.314 0.222 , 0.285	Depositor DCC
R_{free} test set	1764 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.830	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.9	EDS
Estimated twinning fraction	0.047 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 34608 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4976	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1628	0.88	4/2209 (0.2%)
1	B	0.68	0/1629	0.92	5/2207 (0.2%)
1	C	0.64	0/1562	0.85	8/2117 (0.4%)
All	All	0.65	0/4819	0.88	17/6533 (0.3%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	14	ASP	CB-CG-OD2	7.74	125.27	118.30
1	B	83	ASP	CB-CG-OD2	7.50	125.06	118.30
1	C	164	ASP	CB-CG-OD2	6.87	124.48	118.30
1	C	14	ASP	CB-CG-OD2	6.78	124.40	118.30
1	A	83	ASP	CB-CG-OD2	6.72	124.35	118.30
1	B	65	ASP	CB-CG-OD2	6.49	124.14	118.30
1	C	31	ASP	CB-CG-OD2	6.48	124.14	118.30
1	A	69	ASP	CB-CG-OD2	6.06	123.75	118.30
1	B	166	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	B	148	ASP	CB-CG-OD2	5.90	123.61	118.30
1	C	65	ASP	CB-CG-OD2	5.88	123.59	118.30
1	C	185	ASP	CB-CG-OD1	5.66	123.40	118.30
1	B	185	ASP	CB-CG-OD2	5.31	123.08	118.30
1	C	83	ASP	CB-CG-OD2	5.30	123.07	118.30
1	C	106	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	91	LEU	CA-CB-CG	5.05	126.91	115.30
1	C	139	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1597	0	1552	20	1
1	B	1595	0	1565	42	0
1	C	1533	0	1496	33	0
2	A	80	0	0	4	0
2	B	91	0	0	6	0
2	C	80	0	0	1	0
All	All	4976	0	4613	82	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:GLN:HG2	2:B:229:HOH:O	1.66	0.96
1:B:177:GLN:O	1:B:181:GLU:HG3	1.78	0.84
1:A:114:GLU:HG2	1:B:68:ARG:NH2	1.96	0.80
1:A:72:VAL:HG21	1:A:118:LEU:HD11	1.72	0.71
1:B:139:ARG:NH2	1:C:185:ASP:OD2	2.17	0.71
1:A:61:LEU:HD12	1:A:108:ALA:HB1	1.74	0.69
1:B:84:THR:CG2	1:C:139:ARG:HH11	2.07	0.68
1:B:185:ASP:OD2	1:C:139:ARG:NH2	2.20	0.67
1:B:176:TYR:HB2	2:B:240:HOH:O	1.95	0.67
1:B:131:ALA:HB2	2:B:247:HOH:O	1.95	0.65
1:B:22:THR:HG21	1:B:25[B]:ARG:CZ	2.27	0.65
1:B:54:GLU:OE2	1:B:55:ARG:N	2.31	0.64
1:B:195:TRP:CD1	1:B:199:HIS:CE1	2.87	0.63
1:B:84:THR:HG23	1:C:139:ARG:HD2	1.84	0.60
1:B:84:THR:HG23	1:C:139:ARG:HH11	1.67	0.59
1:C:48:VAL:HG11	1:C:50:ARG:HH11	1.67	0.59
1:C:152:ALA:HA	2:C:230:HOH:O	2.04	0.58
1:B:164:ASP:N	2:B:285:HOH:O	2.39	0.56
1:C:68:ARG:HD3	1:C:70:GLU:OE1	2.07	0.55
1:C:68:ARG:NH1	1:C:70:GLU:OE2	2.41	0.53
1:A:68:ARG:HD3	1:A:70:GLU:OE1	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:LEU:HB3	1:C:24:TYR:CE1	2.43	0.53
1:A:99:GLU:O	1:A:102:GLU:HB2	2.09	0.53
1:C:121:LYS:HB2	1:C:146:GLU:HB3	1.91	0.53
1:B:69:ASP:OD2	1:B:172:ARG:NH2	2.40	0.53
1:B:177:GLN:OE1	1:B:181:GLU:OE2	2.28	0.52
1:B:6:LEU:HG	1:C:91:LEU:HD22	1.92	0.52
1:B:111:GLU:O	1:B:115:GLU:HB2	2.10	0.51
1:B:200:HIS:HD2	1:B:204:LYS:NZ	2.09	0.51
1:C:163:GLU:HG2	1:C:165:ILE:HD11	1.94	0.50
1:C:22:THR:HA	1:C:31:ASP:OD1	2.12	0.50
1:B:51:GLU:O	1:B:52:ILE:HD13	2.11	0.50
1:C:104:VAL:HG11	1:C:141:SER:HB2	1.94	0.50
1:B:68:ARG:HD3	1:B:70:GLU:OE1	2.11	0.49
1:B:22:THR:HG22	1:B:31:ASP:OD1	2.12	0.49
1:B:172:ARG:HG2	1:B:172:ARG:HH11	1.78	0.49
1:B:102:GLU:OE2	1:B:110:ARG:NH1	2.35	0.49
1:B:54:GLU:OE2	1:B:56:GLY:N	2.25	0.48
1:A:107:VAL:HG22	1:A:110:ARG:HH12	1.79	0.48
1:B:51:GLU:C	1:B:52:ILE:HD13	2.33	0.48
1:B:80:ALA:HB3	1:C:133:PRO:O	2.14	0.48
1:B:195:TRP:NE1	1:B:199:HIS:CE1	2.83	0.47
1:C:95:ALA:C	1:C:111:GLU:HG3	2.34	0.47
1:B:109:ARG:NH1	1:B:120:VAL:O	2.48	0.47
1:B:75:GLU:HA	1:B:90:LEU:O	2.14	0.47
1:B:80:ALA:HB2	1:C:52:ILE:HD12	1.97	0.47
1:A:86:GLU:HB3	2:A:295:HOH:O	2.15	0.46
1:A:61:LEU:CD1	1:A:108:ALA:HB1	2.42	0.46
1:B:36:ARG:HD3	2:B:290:HOH:O	2.15	0.45
1:C:65:ASP:HA	1:C:66:PRO:HD2	1.79	0.45
1:C:48:VAL:HG11	1:C:50:ARG:NH1	2.31	0.45
1:C:12:LYS:HA	1:C:15:VAL:HG23	1.98	0.45
1:C:136:THR:HG23	1:C:138:GLU:H	1.82	0.45
1:B:84:THR:CG2	1:C:139:ARG:HD2	2.45	0.44
1:A:102:GLU:OE1	1:A:110:ARG:NH1	2.51	0.44
1:B:53:PHE:CE1	1:C:30:LEU:HD11	2.53	0.44
1:A:126:VAL:HG13	1:A:195:TRP:CG	2.53	0.44
1:C:6:LEU:CB	1:C:7:PRO:CD	2.96	0.43
1:B:121:LYS:NZ	2:B:235:HOH:O	2.51	0.43
1:B:33:TYR:O	1:B:49:ARG:HA	2.18	0.43
1:C:74:ILE:O	1:C:74:ILE:HG13	2.18	0.43
1:A:130:LEU:HD21	1:A:139:ARG:NH1	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:GLN:HA	1:A:177:GLN:HE21	1.84	0.43
1:A:18:ILE:N	1:A:34:ARG:O	2.48	0.43
1:C:176:TYR:O	1:C:179:VAL:HB	2.19	0.43
1:B:121:LYS:HB2	1:B:146:GLU:HB3	2.01	0.43
1:B:57:HIS:CD2	1:B:139:ARG:HB3	2.54	0.43
1:C:126:VAL:HG12	1:C:127:LEU:N	2.33	0.43
1:B:165:ILE:CG2	1:B:166:ARG:N	2.82	0.42
1:A:39:LEU:HA	2:A:220:HOH:O	2.18	0.42
1:C:48:VAL:CG1	1:C:50:ARG:HH11	2.32	0.42
1:C:176:TYR:CZ	1:C:180:GLU:HG2	2.55	0.42
1:A:120:VAL:HG23	2:A:222:HOH:O	2.19	0.42
1:C:105:GLU:OE2	1:C:109:ARG:NE	2.50	0.42
1:C:38:ARG:HD3	1:C:38:ARG:HA	1.79	0.42
1:A:55:ARG:O	1:A:55:ARG:HG3	2.19	0.41
1:A:170:VAL:HG11	1:A:178:TRP:CZ3	2.55	0.41
1:A:190:VAL:O	1:A:194:GLN:HG3	2.21	0.41
1:A:121:LYS:HB3	1:A:122:ARG:H	1.68	0.41
1:B:23:LEU:HB3	1:C:24:TYR:CD1	2.56	0.41
1:A:13:ASN:HB2	2:A:229:HOH:O	2.21	0.41
1:B:24:TYR:CZ	1:B:26:GLY:HA3	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:ARG:NH2	1:A:185:ASP:OD2[6_556]	2.12	0.08

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	200/220 (91%)	191 (96%)	9 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	198/220 (90%)	181 (91%)	17 (9%)	0	100	100
1	C	190/220 (86%)	182 (96%)	6 (3%)	2 (1%)	17	25
All	All	588/660 (89%)	554 (94%)	32 (5%)	2 (0%)	46	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	121	LYS
1	C	100	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/186 (89%)	149 (90%)	17 (10%)	9	13
1	B	168/186 (90%)	143 (85%)	25 (15%)	4	4
1	C	160/186 (86%)	144 (90%)	16 (10%)	9	14
All	All	494/558 (88%)	436 (88%)	58 (12%)	7	8

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	15	VAL
1	A	16	GLU
1	A	17	ILE
1	A	21	GLU
1	A	28	SER
1	A	29	SER
1	A	61	LEU
1	A	68	ARG
1	A	85	SER
1	A	91	LEU
1	A	104	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	121	LYS
1	A	147	VAL
1	A	183	LYS
1	A	201	GLN
1	A	206	GLU
1	B	6	LEU
1	B	25[A]	ARG
1	B	25[B]	ARG
1	B	49	ARG
1	B	50	ARG
1	B	54	GLU
1	B	74	ILE
1	B	77	ILE
1	B	84	THR
1	B	85	SER
1	B	91	LEU
1	B	97	MET
1	B	103	SER
1	B	104	VAL
1	B	113	ILE
1	B	115	GLU
1	B	121	LYS
1	B	127	LEU
1	B	139	ARG
1	B	147	VAL
1	B	153	SER
1	B	166	ARG
1	B	177	GLN
1	B	181	GLU
1	B	201	GLN
1	C	15	VAL
1	C	18	ILE
1	C	22	THR
1	C	29	SER
1	C	38	ARG
1	C	50	ARG
1	C	68	ARG
1	C	86	GLU
1	C	91	LEU
1	C	99	GLU
1	C	105	GLU
1	C	119	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	139	ARG
1	C	147	VAL
1	C	174	GLN
1	C	204	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	GLN
1	A	177	GLN
1	A	200	HIS
1	B	57	HIS
1	B	177	GLN
1	B	200	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/220 (92%)	-0.44	2 (0%) 84 83	21, 36, 53, 59	0
1	B	201/220 (91%)	-0.58	0 100 100	19, 33, 46, 55	0
1	C	194/220 (88%)	-0.34	3 (1%) 76 75	22, 39, 54, 59	0
All	All	599/660 (90%)	-0.46	5 (0%) 87 87	19, 35, 53, 59	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	19	ALA	2.9
1	A	27	PHE	2.7
1	C	21	GLU	2.2
1	C	48	VAL	2.1
1	A	48	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.