



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:57 PM GMT

PDB ID : 3VOL
Title : X-ray Crystal Structure of PAS-HAMP Aer2 in the CN-bound Form
Authors : Sawai, H.; Sugimoto, H.; Shiro, Y.; Aono, S.
Deposited on : 2012-01-27
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div></div> <div> <div></div> <div>39%</div> <div>18%</div> <div>•</div> <div>41%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1123 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aerotaxis transducer Aer2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	0	0
			1067	654	206	202	5			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	152	MET	-	EXPRESSION TAG	UNP Q9I6V6
A	153	GLY	-	EXPRESSION TAG	UNP Q9I6V6
A	154	SER	-	EXPRESSION TAG	UNP Q9I6V6
A	155	SER	-	EXPRESSION TAG	UNP Q9I6V6
A	156	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	157	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	158	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	159	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	160	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	161	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	162	SER	-	EXPRESSION TAG	UNP Q9I6V6
A	163	SER	-	EXPRESSION TAG	UNP Q9I6V6
A	164	GLY	-	EXPRESSION TAG	UNP Q9I6V6
A	165	LEU	-	EXPRESSION TAG	UNP Q9I6V6
A	166	VAL	-	EXPRESSION TAG	UNP Q9I6V6
A	167	PRO	-	EXPRESSION TAG	UNP Q9I6V6
A	168	ARG	-	EXPRESSION TAG	UNP Q9I6V6
A	169	GLY	-	EXPRESSION TAG	UNP Q9I6V6
A	170	SER	-	EXPRESSION TAG	UNP Q9I6V6
A	171	HIS	-	EXPRESSION TAG	UNP Q9I6V6
A	172	MET	-	EXPRESSION TAG	UNP Q9I6V6

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is CYANIDE ION (three-letter code: CYN) (formula: CN).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			2	1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total	O	0	0
			11	11		

- Molecule 1: Aerotaxis transducer Aer2



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	83.02Å 83.02Å 107.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.90 – 2.40 43.12 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.90-2.40) 99.5 (43.12-2.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.55 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.223 , 0.268 0.212 , 0.254	Depositor DCC
R_{free} test set	432 reflections (4.76%)	DCC
Wilson B-factor (Å ²)	58.8	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 9083 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1123	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CYN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/1082 (0.1%)	0.98	1/1461 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	GLU	CG-CD	5.21	1.59	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	276	ARG	NE-CZ-NH1	5.30	122.95	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1067	0	1049	30	0
2	A	43	0	30	7	0
3	A	2	0	0	0	0
4	A	11	0	0	0	0
All	All	1123	0	1079	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:HEM:HMB1	2:A:401:HEM:HBB2	1.49	0.94
2:A:401:HEM:HMC1	2:A:401:HEM:HBC2	1.51	0.92
2:A:401:HEM:CMC	2:A:401:HEM:HBC2	2.09	0.83
2:A:401:HEM:HBB2	2:A:401:HEM:CMB	2.12	0.78
1:A:288:GLU:HG2	1:A:289:GLU:OE1	1.86	0.75
1:A:288:GLU:HG2	1:A:289:GLU:H	1.64	0.63
2:A:401:HEM:HBA2	2:A:401:HEM:HHA	1.81	0.61
1:A:243:LEU:C	1:A:243:LEU:HD23	2.24	0.58
1:A:229:ASN:O	1:A:232:VAL:HG23	2.04	0.57
1:A:293:GLU:O	1:A:297:SER:HB2	2.07	0.54
2:A:401:HEM:CBB	2:A:401:HEM:HMB1	2.32	0.53
1:A:289:GLU:O	1:A:293:GLU:HG3	2.10	0.52
1:A:234:HIS:O	1:A:237:PRO:HD3	2.10	0.51
2:A:401:HEM:CBA	2:A:401:HEM:HHA	2.41	0.51
1:A:241:ARG:HH11	1:A:241:ARG:CG	2.25	0.49
1:A:247:LEU:HG	1:A:268:PRO:HD3	1.94	0.49
1:A:241:ARG:HH11	1:A:241:ARG:HG2	1.77	0.48
1:A:302:ALA:O	1:A:305:ALA:O	2.31	0.48
1:A:260:ARG:NH1	1:A:285:ASP:OD2	2.47	0.47
1:A:198:MET:HE3	1:A:225:LEU:HB3	1.95	0.47
1:A:171:HIS:O	1:A:175:ILE:HG13	2.14	0.46
1:A:214:ARG:NH2	1:A:221:ASP:OD1	2.48	0.46
1:A:305:ALA:O	1:A:306:GLY:C	2.53	0.46
1:A:208:ARG:HG3	1:A:208:ARG:H	1.57	0.44
1:A:198:MET:HB3	1:A:198:MET:HE2	1.78	0.44
1:A:171:HIS:HA	1:A:174:ARG:CZ	2.49	0.43
1:A:231:ASP:O	1:A:233:PHE:N	2.50	0.43
1:A:243:LEU:O	1:A:243:LEU:HD23	2.19	0.43
1:A:264:LEU:HD12	1:A:266:VAL:HG23	2.01	0.43
1:A:172:MET:O	1:A:176:LYS:HG3	2.20	0.42
1:A:231:ASP:C	1:A:233:PHE:H	2.22	0.42
1:A:295:GLU:HG2	1:A:296:VAL:N	2.35	0.41
1:A:204:GLU:O	1:A:208:ARG:HG3	2.20	0.41
1:A:231:ASP:C	1:A:233:PHE:N	2.74	0.41
1:A:197:TYR:CG	1:A:198:MET:N	2.88	0.41
1:A:187:MET:HE1	1:A:206:LEU:HD11	2.02	0.41
1:A:225:LEU:O	1:A:228:ALA:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	136/233 (58%)	125 (92%)	10 (7%)	1 (1%)	26 38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	274	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	111/190 (58%)	99 (89%)	12 (11%)	8 11

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	MET
1	A	212	ASP
1	A	217	LEU
1	A	232	VAL
1	A	241	ARG
1	A	260	ARG
1	A	264	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	272	ASP
1	A	282	GLN
1	A	286	ARG
1	A	294	GLN
1	A	297	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	401	1	30,50,50	2.00	5 (16%)	24,82,82	3.11	12 (50%)
3	CYN	A	402	-	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	401	1	-	2/10/54/54	0/0/8/8
3	CYN	A	402	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	HEM	C3B-C4B	-6.06	1.46	1.51
2	A	401	HEM	C2C-C1C	-5.14	1.42	1.52
2	A	401	HEM	C3D-C4D	-4.01	1.46	1.51
2	A	401	HEM	CAA-C2A	2.02	1.55	1.52
2	A	401	HEM	C1C-NC	3.91	1.40	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	C3C-CAC-CBC	-6.37	114.68	124.46
2	A	401	HEM	C4B-CHC-C1C	-6.16	115.52	125.82
2	A	401	HEM	CAA-CBA-CGA	-4.50	104.49	112.75
2	A	401	HEM	C3B-CAB-CBB	-2.77	120.21	124.46
2	A	401	HEM	CAA-C2A-C3A	-2.42	122.10	129.00
2	A	401	HEM	CMD-C2D-C3D	2.28	124.43	114.35
2	A	401	HEM	CAA-C2A-C1A	2.56	129.79	127.01
2	A	401	HEM	CBA-CAA-C2A	2.76	117.48	112.53
2	A	401	HEM	CMB-C2B-C3B	3.98	126.46	116.53
2	A	401	HEM	CAD-C3D-C4D	4.47	128.22	112.47
2	A	401	HEM	CAD-C3D-C2D	4.79	126.98	113.22
2	A	401	HEM	CMC-C2C-C3C	5.57	130.44	116.53

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	HEM	C3A-C2A-CAA-CBA
2	A	401	HEM	C1A-C2A-CAA-CBA

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	HEM	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/233 (59%)	0.32	3 (2%) 65 64	39, 64, 92, 110	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	298	GLN	2.4
1	A	291	ARG	2.4
1	A	216	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	HEM	A	401	43/43	0.99	0.15	-0.21	39,50,62,70	0
3	CYN	A	402	2/2	0.99	0.16	-	37,37,37,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.