



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:59 PM GMT

PDB ID : 3VOT
Title : Crystal structure of L-amino acid ligase from *Bacillus licheniformis*
Authors : Suzuki, M.; Takahashi, Y.; Noguchi, A.; Arai, T.; Yagasaki, M.; Kino, K.; Saito, J.
Deposited on : 2012-02-08
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	<div> <div>5%</div> <div>88%</div> <div>8%</div> <div>••</div> </div>
1	B	425	<div> <div>11%</div> <div>85%</div> <div>7%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PG4	B	507	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7123 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-amino acid ligase, BL00235.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	4	0
			3288	2105	565	609	9			
1	B	394	Total	C	N	O	S	0	3	0
			3123	2000	529	585	9			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	4	Total Ca 4 4	0	0
3	A	5	Total Ca 5 5	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Cl 1 1	0	0
4	A	1	Total Cl 1 1	0	0

- Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 13 8 5	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	341	Total O 342 342	0	1
6	B	291	Total O 292 292	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-amino acid ligase, BL00235



- Molecule 1: L-amino acid ligase, BL00235



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.09 Å 90.03 Å 154.91 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.46 – 1.80 29.46 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.1 (29.46-1.80) 96.2 (29.46-1.80)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.80 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.196 , 0.241 0.195 , 0.241	Depositor DCC
R_{free} test set	3625 reflections (5.21%)	DCC
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.210	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 73160 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7123	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.96 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.2829e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PG4, ADP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/3374	0.55	0/4565
1	B	0.42	0/3198	0.53	0/4329
All	All	0.42	0/6572	0.54	0/8894

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3288	0	3268	16	0
1	B	3123	0	3084	16	0
2	A	27	0	11	0	0
2	B	27	0	11	0	0
3	A	5	0	0	0	0
3	B	4	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	13	0	18	0	0
6	A	342	0	0	0	0
6	B	292	0	0	4	0
All	All	7123	0	6392	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:ARG:HG2	1:B:111:ARG:HH11	1.51	0.75
1:B:153:VAL:HG22	1:B:196:VAL:HG23	1.76	0.68
1:B:54:LEU:HD23	1:B:56:LEU:HD22	1.78	0.66
1:B:95:GLU:HG2	6:B:749:HOH:O	1.98	0.64
1:A:315[A]:GLN:HA	1:A:315[A]:GLN:HE21	1.69	0.57
1:A:203:GLY:HA3	1:A:276:LEU:O	2.08	0.54
1:B:353:ASP:HB2	1:B:414:ILE:HG13	1.90	0.54
1:B:117:ARG:HG3	1:B:130:PHE:CD2	2.44	0.53
1:B:170:LEU:O	1:B:174:VAL:HG23	2.08	0.52
1:B:111:ARG:HG2	1:B:111:ARG:NH1	2.23	0.52
1:B:68:ARG:HD3	6:B:834:HOH:O	2.09	0.52
1:B:209:GLU:OE1	1:B:238:TYR:OH	2.25	0.50
1:A:117:ARG:HG3	1:A:130[B]:PHE:CD2	2.48	0.49
1:A:163[A]:ARG:NH2	1:A:229:LYS:O	2.45	0.49
1:A:232:PHE:CE2	1:A:381:PRO:HB3	2.48	0.48
1:A:295:GLY:O	1:A:298:HIS:HB3	2.15	0.47
1:A:27:LEU:HD11	1:A:311:GLN:HG3	1.97	0.46
1:A:315[A]:GLN:HA	1:A:315[A]:GLN:NE2	2.31	0.45
1:A:133:PHE:CZ	1:A:139:LEU:HD11	2.52	0.45
1:A:113:LYS:HG3	1:A:153:VAL:O	2.17	0.44
1:A:245:LYS:HE2	1:A:245:LYS:HB3	1.76	0.44
1:B:136:LEU:HB3	1:B:174:VAL:HG11	2.00	0.43
1:B:230:GLY:HA3	1:B:231:PRO:HA	1.80	0.43
1:B:116:THR:O	1:B:120:LEU:HG	2.19	0.43
1:A:141:ASN:HD22	1:A:141:ASN:C	2.22	0.42
1:B:315:GLN:HG2	6:B:637:HOH:O	2.18	0.42
1:B:299:TYR:OH	1:B:365:LYS:HG3	2.20	0.41
1:B:326:PHE:HB3	6:B:868:HOH:O	2.20	0.41
1:A:331:ARG:O	1:A:331:ARG:HG3	2.20	0.41
1:A:136:LEU:HD11	1:A:171:GLU:HG2	2.03	0.41
1:A:116:THR:O	1:A:120:LEU:HG	2.21	0.40
1:A:260:VAL:HG13	1:A:265:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	411/425 (97%)	399 (97%)	11 (3%)	1 (0%)	52	35
1	B	389/425 (92%)	376 (97%)	13 (3%)	0	100	100
All	All	800/850 (94%)	775 (97%)	24 (3%)	1 (0%)	56	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	12	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/363 (98%)	341 (96%)	15 (4%)	36	18
1	B	338/363 (93%)	329 (97%)	9 (3%)	52	36
All	All	694/726 (96%)	670 (96%)	24 (4%)	45	25

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LYS
1	A	44	ASN
1	A	132	GLU
1	A	136	LEU
1	A	141	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	176	LYS
1	A	192	LYS
1	A	234	GLU
1	A	238	TYR
1	A	250	LEU
1	A	278	LYS
1	A	331	ARG
1	A	376[A]	LYS
1	A	376[B]	LYS
1	A	378	LEU
1	B	30	LYS
1	B	61	GLU
1	B	133	PHE
1	B	195	ILE
1	B	238	TYR
1	B	278	LYS
1	B	329	GLU
1	B	344	GLN
1	B	365	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	114	ASN
1	A	141	ASN
1	B	5	ASN
1	B	109	ASN
1	B	416	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 11 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADP	A	501	3	22,29,29	1.01	2 (9%)	27,45,45	1.83	3 (11%)
2	ADP	B	501	3	22,29,29	1.04	2 (9%)	27,45,45	1.88	4 (14%)
5	PG4	B	507	-	12,12,12	0.50	0	11,11,11	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	501	3	-	0/12/32/32	0/3/3/3
2	ADP	B	501	3	-	0/12/32/32	0/3/3/3
5	PG4	B	507	-	-	0/10/10/10	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ADP	O4'-C1'	2.03	1.43	1.41
2	B	501	ADP	O4'-C1'	2.25	1.44	1.41
2	A	501	ADP	C5-C4	3.01	1.47	1.40
2	B	501	ADP	C5-C4	3.05	1.47	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ADP	N3-C2-N1	-7.18	123.39	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	ADP	N3-C2-N1	-6.89	123.62	128.89
2	B	501	ADP	C2'-C1'-N9	-3.90	108.34	114.29
2	A	501	ADP	C2'-C1'-N9	-3.59	108.80	114.29
2	B	501	ADP	C4-C5-N7	-2.85	106.86	109.48
2	A	501	ADP	C4-C5-N7	-2.29	107.37	109.48
2	B	501	ADP	O4'-C1'-N9	2.28	112.87	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	411/425 (96%)	0.12	21 (5%) 32 26	11, 25, 53, 71	0
1	B	394/425 (92%)	0.41	47 (11%) 6 4	12, 26, 68, 97	0
All	All	805/850 (94%)	0.27	68 (8%) 14 11	11, 26, 59, 97	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	136	LEU	6.4
1	B	2	THR	6.0
1	B	194	GLY	5.9
1	B	175	ARG	5.7
1	B	154	ASN	5.3
1	B	57	PHE	5.2
1	A	155	GLY	5.1
1	B	174	VAL	5.0
1	B	328	GLY	4.9
1	B	144	LEU	4.8
1	B	153	VAL	4.7
1	B	193	THR	4.3
1	B	177	VAL	4.3
1	B	155	GLY	4.3
1	B	56	LEU	4.2
1	B	135	THR	4.1
1	B	133	PHE	4.1
1	B	176	LYS	4.1
1	B	230	GLY	3.8
1	B	382	HIS	3.7
1	A	183	ARG	3.6
1	B	134	HIS	3.6
1	B	140	GLU	3.5
1	A	328	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	171	GLU	3.3
1	B	138	ASP	3.2
1	A	378	LEU	3.2
1	B	146	TYR	3.2
1	B	329	GLU	3.2
1	B	417	ASN	3.2
1	B	166	ASP	3.1
1	A	382	HIS	3.1
1	A	372	ARG	3.0
1	A	172	GLU	3.0
1	B	232	PHE	3.0
1	B	139	LEU	2.9
1	B	379	PRO	2.9
1	A	376[A]	LYS	2.9
1	A	141	ASN	2.7
1	B	247	GLU	2.7
1	B	58	GLU	2.7
1	A	185	LEU	2.6
1	B	229	LYS	2.6
1	B	130	PHE	2.6
1	B	170	LEU	2.6
1	B	372	ARG	2.5
1	A	381	PRO	2.5
1	A	2	THR	2.5
1	B	279	ASP	2.5
1	A	142	ARG	2.5
1	B	61	GLU	2.4
1	A	327	GLU	2.4
1	B	173	ALA	2.4
1	B	59	ASP	2.4
1	B	383	PHE	2.4
1	A	380	TYR	2.3
1	A	383	PHE	2.3
1	B	145	SER	2.3
1	B	167	ARG	2.3
1	B	137	ALA	2.3
1	A	187	ARG	2.2
1	B	152	PRO	2.2
1	A	143	LYS	2.2
1	A	186	ASN	2.1
1	B	234	GLU	2.1
1	A	154	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	168	LYS	2.1
1	A	278	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	PG4	B	507	13/13	0.73	0.27	2.03	43,46,55,59	0
2	ADP	A	501	27/27	0.91	0.12	-0.26	23,25,42,42	0
2	ADP	B	501	27/27	0.90	0.11	-0.41	26,30,39,40	0
4	CL	A	507	1/1	0.99	0.03	-2.87	23,23,23,23	0
4	CL	B	506	1/1	0.99	0.05	-5.00	26,26,26,26	0
3	CA	B	503	1/1	0.98	0.03	-	43,43,43,43	0
3	CA	A	503	1/1	0.94	0.08	-	49,49,49,49	0
3	CA	B	504	1/1	0.94	0.08	-	46,46,46,46	0
3	CA	A	502	1/1	0.97	0.21	-	43,43,43,43	0
3	CA	B	505	1/1	0.99	0.04	-	33,33,33,33	0
3	CA	B	502	1/1	0.98	0.03	-	39,39,39,39	0
3	CA	A	504	1/1	0.96	0.07	-	42,42,42,42	0
3	CA	A	506	1/1	0.92	0.06	-	59,59,59,59	0
3	CA	A	505	1/1	0.99	0.03	-	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.