



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:01 AM GMT

PDB ID : 2VPS  
Title : STRUCTURE OF THE BIFUNCTIONAL LEISHMANIA MAJOR TRY-  
PANOTHIONE SYNTHETASE-AMIDASE  
Authors : Fyfe, P.K.; Oza, S.L.; Fairlamb, A.H.; Hunter, W.N.  
Deposited on : 2008-03-04  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	652	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4939 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPANOTHIONE SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	603	Total	C	N	O	S	0	0	0
			4887	3126	831	909	21			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Cl	0	0
			5	5		

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Br	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	45	Total	O	0	0
			45	45		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: TRYPANOTHIONE SYNTHETASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.05Å 85.60Å 168.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.33 – 2.75 20.00 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.5 (28.33-2.75) 99.5 (20.00-2.75)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 2.75Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.204 , 0.250 0.208 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	53.3	Xtriage
Anisotropy	0.374	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 46.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 27197 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4939	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/5024	0.73	3/6814 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	57	TYR	N-CA-C	5.46	125.76	111.00
1	A	3	SER	N-CA-C	5.19	125.01	111.00
1	A	503	GLU	N-CA-C	-5.15	97.10	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	281	THR	CB

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	502	PHE	Peptide
1	A	56	LYS	Peptide
1	A	621	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4887	0	4674	52	0
2	A	5	0	0	0	0
3	A	2	0	0	0	0
4	A	45	0	0	4	0
All	All	4939	0	4674	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:SER:O	1:A:352:THR:HB	1.77	0.84
1:A:79:ASN:H	1:A:79:ASN:ND2	1.91	0.68
1:A:514:ALA:HB2	1:A:550:ILE:HD13	1.76	0.67
1:A:579:ASN:C	1:A:579:ASN:HD22	2.01	0.64
1:A:590:LYS:HE3	1:A:595:TYR:CZ	2.34	0.63
1:A:265:TYR:CE1	1:A:369:LEU:HD11	2.34	0.62
1:A:121:PRO:HB3	1:A:186:LEU:HD21	1.84	0.59
1:A:620:THR:O	1:A:624:SER:HB2	2.06	0.55
1:A:623:ASP:OD1	1:A:623:ASP:N	2.39	0.55
1:A:178:ASP:N	1:A:178:ASP:OD1	2.39	0.54
1:A:308:GLU:HA	1:A:311:TRP:CE2	2.43	0.53
1:A:400:PHE:CE2	1:A:416:MET:HG3	2.45	0.52
1:A:374:THR:HG22	1:A:632:LYS:HA	1.93	0.51
1:A:178:ASP:HB3	4:A:2022:HOH:O	2.10	0.51
1:A:281:THR:CG2	4:A:2008:HOH:O	2.58	0.50
1:A:284:HIS:O	1:A:288:MET:HG2	2.12	0.50
1:A:472:GLU:O	1:A:474:ARG:N	2.46	0.49
1:A:150:ARG:NH2	1:A:177:ILE:HD12	2.28	0.48
1:A:281:THR:HG21	4:A:2008:HOH:O	2.14	0.48
1:A:220:LEU:HD21	1:A:612:ILE:HD11	1.96	0.47
1:A:79:ASN:N	1:A:79:ASN:ND2	2.60	0.47
1:A:550:ILE:HG22	1:A:551:VAL:H	1.80	0.47
1:A:512:ASN:OD1	1:A:550:ILE:HG21	2.15	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LEU:HD22	1:A:175:ASP:HA	1.97	0.45
1:A:219:TYR:CD2	1:A:219:TYR:C	2.89	0.45
1:A:612:ILE:CG2	1:A:613:ARG:N	2.80	0.44
1:A:218:PRO:HA	1:A:269:ASN:HD21	1.82	0.44
1:A:10:SER:HB2	1:A:322:GLN:HB2	1.99	0.44
1:A:123:THR:OG1	1:A:126:ASN:N	2.51	0.43
1:A:645:ILE:O	1:A:645:ILE:HG22	2.19	0.43
1:A:420:GLU:HA	1:A:424:LEU:O	2.19	0.43
1:A:372:GLN:HE21	1:A:372:GLN:HB2	1.64	0.43
1:A:79:ASN:HD22	1:A:79:ASN:H	1.63	0.43
1:A:614:GLU:O	1:A:625:PRO:HD2	2.18	0.43
1:A:531:LYS:HG3	1:A:584:GLN:NE2	2.33	0.43
1:A:66:TRP:CZ2	1:A:70:ARG:HG2	2.55	0.42
1:A:588:LEU:N	1:A:588:LEU:CD1	2.82	0.42
1:A:321:GLN:HE22	1:A:501:TYR:H	1.68	0.42
1:A:592:ASP:O	1:A:593:ASP:HB2	2.19	0.42
1:A:488:LEU:HD12	1:A:492:LEU:HD13	2.02	0.42
1:A:472:GLU:N	1:A:472:GLU:OE1	2.53	0.41
1:A:79:ASN:HD22	1:A:79:ASN:N	2.18	0.41
1:A:620:THR:O	1:A:624:SER:CB	2.68	0.41
1:A:178:ASP:CB	4:A:2022:HOH:O	2.68	0.41
1:A:286:ILE:HG12	1:A:523:HIS:CD2	2.56	0.41
1:A:509:ILE:HB	1:A:510:PRO:HD3	2.01	0.41
1:A:294:VAL:HG21	1:A:304:PHE:CZ	2.56	0.41
1:A:375:ARG:NH2	1:A:379:PHE:CD2	2.89	0.41
1:A:84:ILE:HA	1:A:87:LEU:HD22	2.02	0.40
1:A:88:LYS:O	1:A:101:VAL:HG22	2.22	0.40
1:A:622:VAL:HB	1:A:623:ASP:OD1	2.21	0.40
1:A:33:TYR:HB2	1:A:57:TYR:CG	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	595/652 (91%)	559 (94%)	32 (5%)	4 (1%)	26	59

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	352	THR
1	A	473	GLU
1	A	439	ASP
1	A	622	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	515/556 (93%)	480 (93%)	35 (7%)	20	46

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	SER
1	A	16	HIS
1	A	48	GLU
1	A	78	VAL
1	A	79	ASN
1	A	87	LEU
1	A	149	TYR
1	A	163	LYS
1	A	178	ASP
1	A	181	GLU
1	A	199	GLU
1	A	219	TYR
1	A	221	LEU
1	A	237	MET
1	A	296	GLU
1	A	352	THR
1	A	358	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	362	LYS
1	A	369	LEU
1	A	372	GLN
1	A	383	ARG
1	A	387	MET
1	A	433	ASP
1	A	456	LYS
1	A	474	ARG
1	A	477	ASN
1	A	525	GLU
1	A	550	ILE
1	A	579	ASN
1	A	581	ILE
1	A	587	GLU
1	A	620	THR
1	A	623	ASP
1	A	624	SER
1	A	646	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	79	ASN
1	A	86	GLN
1	A	152	HIS
1	A	233	ASN
1	A	270	HIS
1	A	293	GLN
1	A	321	GLN
1	A	346	ASN
1	A	372	GLN
1	A	384	ASN
1	A	579	ASN
1	A	583	GLN
1	A	584	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	603/652 (92%)	0.21	35 (5%)	26 20	47, 61, 75, 88	12 (1%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	372	GLN	5.5
1	A	228	THR	5.4
1	A	476	GLU	5.3
1	A	233	ASN	4.9
1	A	551	VAL	4.8
1	A	483	LYS	4.8
1	A	477	ASN	4.1
1	A	232	ALA	3.8
1	A	14	PRO	3.7
1	A	646	ASP	3.5
1	A	644	ASP	3.4
1	A	645	ILE	3.3
1	A	178	ASP	3.3
1	A	238	ASN	3.2
1	A	647	LYS	3.0
1	A	124	ASP	3.0
1	A	621	GLY	2.8
1	A	213	LYS	2.7
1	A	180	ASP	2.6
1	A	229	GLU	2.5
1	A	230	SER	2.5
1	A	446	SER	2.5
1	A	181	GLU	2.4
1	A	153	LYS	2.4
1	A	15	GLY	2.4
1	A	200	GLY	2.4
1	A	262	VAL	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	415	CYS	2.3
1	A	166	HIS	2.3
1	A	125	ALA	2.2
1	A	485	LYS	2.2
1	A	474	ARG	2.1
1	A	622	VAL	2.0
1	A	214	GLU	2.0
1	A	224	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	1659	1/1	0.96	0.11	-1.22	37,37,37,37	0
2	CL	A	1653	1/1	0.98	0.10	-1.80	52,52,52,52	0
3	BR	A	1657	1/1	0.98	0.06	-5.62	71,71,71,71	0
2	CL	A	1655	1/1	0.98	0.07	-	47,47,47,47	0
2	CL	A	1654	1/1	0.91	0.08	-	80,80,80,80	0
2	CL	A	1656	1/1	0.93	0.08	-	52,52,52,52	0
3	BR	A	1658	1/1	0.96	0.07	-	101,101,101,101	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.