



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:05 AM GMT

PDB ID : 2VVS  
Title : BTGH84 STRUCTURE IN COMPLEX WITH PUGNAC  
Authors : Macauley, M.S.; Bubb, A.; Martinez-Fleites, C.; Davies, G.J.; Vocadlo, D.J.  
Deposited on : 2008-06-11  
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1611 (2.26-2.22)
Clashscore	102246	1764 (2.26-2.22)
Ramachandran outliers	100387	1724 (2.26-2.22)
Sidechain outliers	100360	1724 (2.26-2.22)
RSRZ outliers	91569	1616 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	737	<div> <div>5%</div> <div>63%</div> <div>14%</div> <div>•</div> <div>21%</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-GLCNACASE BT\_4395.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	580	Total	C	N	O	S	0	2	0
			4727	3027	795	888	17			

- Molecule 2 is O-(2-ACETAMIDO-2-DEOXY D-GLUCOPYRANOSYLIDENE) AMINO-N-PHENYLCARBAMATE (three-letter code: OAN) (formula: C<sub>15</sub>H<sub>19</sub>N<sub>3</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			25	15	3	7		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	111	Total	O	0	0
			111	111		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	185.64Å 51.34Å 81.46Å 90.00° 96.97° 90.00°	Depositor
Resolution (Å)	64.82 – 2.24 64.80 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.8 (64.82-2.24) 99.8 (64.80-2.24)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 2.25Å)	Xtriage
Refinement program	REFMAC 5.4.0065	Depositor
R, $R_{free}$	0.203 , 0.243 0.202 , 0.243	Depositor DCC
$R_{free}$ test set	1845 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.4	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 54.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 36976 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4863	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.78	2/4855 (0.0%)	0.79	7/6587 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	580	PHE	CE1-CZ	10.90	1.58	1.37
1	A	576	PHE	CG-CD1	5.98	1.47	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	580	PHE	CB-CG-CD1	-11.78	112.55	120.80
1	A	520	ASN	CB-CA-C	7.10	124.60	110.40
1	A	104	LEU	CA-CB-CG	-7.05	99.07	115.30
1	A	45	LEU	CA-CB-CG	7.03	131.46	115.30
1	A	580	PHE	CG-CD1-CE1	-6.75	113.37	120.80
1	A	580	PHE	CD1-CG-CD2	6.33	126.52	118.30
1	A	576	PHE	CB-CG-CD1	-5.51	116.94	120.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	460	TYR	Sidechain
1	A	576	PHE	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4727	0	4610	64	0
2	A	25	0	19	3	0
3	A	111	0	0	2	0
All	All	4863	0	4629	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LYS:HD3	1:A:216:LYS:H	1.44	0.81
1:A:408:ARG:NH2	1:A:419:GLU:OE2	2.20	0.75
1:A:536:GLN:HG2	1:A:590:TYR:OH	1.87	0.75
1:A:438:GLU:OE1	1:A:441:MET:HG3	1.87	0.74
1:A:539:PHE:O	1:A:543:GLN:HG3	1.92	0.70
1:A:318:ILE:HG21	1:A:366:MET:HE1	1.78	0.64
1:A:297:LEU:O	1:A:301:LEU:HB2	1.99	0.62
1:A:438:GLU:OE1	1:A:441:MET:CG	2.49	0.61
1:A:173:ALA:HB1	1:A:174:PRO:HA	1.84	0.59
1:A:210:GLN:H	1:A:210:GLN:CD	2.06	0.59
1:A:301:LEU:HD12	1:A:307:ILE:HD11	1.86	0.58
1:A:130:ARG:HG2	1:A:391:ALA:HB1	1.86	0.58
1:A:78:LYS:HE2	1:A:128:ARG:HG2	1.86	0.57
1:A:535:GLN:HE22	1:A:562:LYS:NZ	2.04	0.56
2:A:900:OAN:OAR	2:A:900:OAN:HAT	2.04	0.55
1:A:81:TYR:CE2	1:A:123:ASP:HB3	2.41	0.55
1:A:354:PRO:HB2	1:A:399:THR:HG22	1.90	0.54
1:A:341:PRO:HD2	1:A:373:PRO:HA	1.89	0.53
1:A:236:SER:HB3	1:A:274:GLN:HB2	1.91	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LYS:CD	1:A:216:LYS:H	2.10	0.52
1:A:59:ILE:HA	1:A:92:LEU:O	2.09	0.52
1:A:308:MET:HA	1:A:335:TYR:O	2.09	0.52
1:A:310:THR:HA	1:A:337:TRP:O	2.09	0.52
1:A:519:ARG:HB3	1:A:520:ASN:HD22	1.76	0.51
1:A:155:LYS:NZ	3:A:2030:HOH:O	2.42	0.51
1:A:128:ARG:HD3	1:A:129:TYR:CZ	2.45	0.51
1:A:41:LEU:HB2	1:A:104:LEU:HD21	1.93	0.49
1:A:186:ALA:HA	1:A:189:GLN:HE21	1.76	0.49
1:A:406:ALA:O	1:A:410:ILE:HG12	2.13	0.49
1:A:355:VAL:O	1:A:399:THR:HG23	2.12	0.49
1:A:517:GLU:O	1:A:517:GLU:HG3	2.12	0.49
1:A:163:TYR:CD2	1:A:163:TYR:C	2.87	0.48
1:A:445:PRO:O	1:A:449:ARG:HG3	2.14	0.48
1:A:462:LYS:O	1:A:466:GLU:HG3	2.14	0.47
1:A:516:VAL:C	1:A:518:GLY:H	2.18	0.47
1:A:283:ASN:O	1:A:284:LYS:C	2.52	0.47
1:A:126:SER:HB2	1:A:394:PRO:HD2	1.95	0.47
1:A:354:PRO:HB2	1:A:399:THR:CG2	2.45	0.46
1:A:261:ILE:O	1:A:265:PHE:HB3	2.15	0.46
1:A:562:LYS:HB3	1:A:563:PRO:HD3	1.97	0.46
1:A:408:ARG:HE	1:A:408:ARG:HB2	1.48	0.46
1:A:38:VAL:O	1:A:42:LYS:HG3	2.16	0.46
2:A:900:OAN:CAT	2:A:900:OAN:OAR	2.64	0.45
1:A:132:VAL:HG22	1:A:371:THR:HG23	1.99	0.44
1:A:129:TYR:O	1:A:368:GLY:HA2	2.17	0.44
1:A:574:LYS:NZ	1:A:578:GLN:HE22	2.15	0.44
1:A:138:GLY:O	1:A:140:PRO:HD3	2.18	0.44
1:A:266:ALA:HA	1:A:272:ILE:HD12	1.99	0.44
1:A:85:VAL:O	1:A:118:GLU:HG3	2.17	0.44
1:A:517:GLU:CG	1:A:517:GLU:O	2.66	0.44
1:A:483:LEU:HG	1:A:499:VAL:HG11	2.00	0.44
1:A:288:ASN:HA	1:A:289:PRO:HD2	1.77	0.44
1:A:163:TYR:O	1:A:163:TYR:CD2	2.72	0.43
1:A:424:HIS:HB2	1:A:502:PHE:HE1	1.83	0.43
1:A:301:LEU:HD22	1:A:305:ILE:HD12	2.00	0.43
1:A:203:VAL:HG22	1:A:236:SER:HB2	2.01	0.42
1:A:428:LEU:HB3	1:A:431:ASN:HD22	1.84	0.42
1:A:78:LYS:HE2	1:A:128:ARG:CG	2.49	0.42
1:A:444:GLN:N	1:A:445:PRO:HD2	2.35	0.42
1:A:41:LEU:O	1:A:45:LEU:HD22	2.20	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:ASN:O	1:A:289:PRO:C	2.59	0.41
1:A:242:ASP:O	1:A:244:ILE:HG23	2.20	0.41
1:A:430:PRO:HD3	3:A:2088:HOH:O	2.20	0.41
1:A:325:TRP:O	1:A:329:ARG:NH2	2.53	0.41
1:A:337:TRP:CE2	2:A:900:OAN:HAH3	2.57	0.40
1:A:443:ILE:HD12	1:A:443:ILE:C	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	577/737 (78%)	546 (95%)	31 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	510/647 (79%)	485 (95%)	25 (5%)	31	33

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	27	LEU
1	A	45	LEU
1	A	55	MET
1	A	78	LYS
1	A	89	GLU
1	A	104	LEU
1	A	128	ARG
1	A	216	LYS
1	A	219	ARG
1	A	221	LEU
1	A	287	SER
1	A	292	ASN
1	A	300	LYS
1	A	301	LEU
1	A	316	SER
1	A	337	TRP
1	A	408	ARG
1	A	423	MET
1	A	455	LYS
1	A	458	LYS
1	A	519	ARG
1	A	537	GLN
1	A	588	THR
1	A	589	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	26	GLN
1	A	156	ASN
1	A	189	GLN
1	A	520	ASN
1	A	529	ASN
1	A	535	GLN
1	A	543	GLN
1	A	578	GLN
1	A	581	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	OAN	A	900	-	22,26,26	1.34	2 (9%)	22,35,35	0.88	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OAN	A	900	-	-	0/12/35/35	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	900	OAN	CAS-NAO	-3.94	1.34	1.41
2	A	900	OAN	CAA-NAY	2.95	1.35	1.27

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	900	OAN	CAB-NAI-CAG	2.11	124.93	122.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	900	OAN	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	580/737 (78%)	0.57	38 (6%) 22 23	33, 45, 57, 91	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	591	MET	8.1
1	A	592	PRO	8.1
1	A	590	TYR	7.8
1	A	588	THR	6.6
1	A	463	ALA	4.7
1	A	454	PHE	4.3
1	A	456	GLU	4.3
1	A	576	PHE	4.0
1	A	462	LYS	3.7
1	A	44	LEU	3.6
1	A	524	PHE	3.4
1	A	25	TYR	3.4
1	A	460	TYR	3.2
1	A	26	GLN	3.2
1	A	19	ILE	3.1
1	A	574	LYS	3.1
1	A	23	ALA	3.0
1	A	522	SER	3.0
1	A	580	PHE	3.0
1	A	575	PHE	2.9
1	A	469	GLN	2.9
1	A	113	ASP	2.8
1	A	525	LEU	2.7
1	A	27	LEU	2.6
1	A	21	LEU	2.6
1	A	519	ARG	2.5
1	A	111	LEU	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	117	PRO	2.5
1	A	244	ILE	2.5
1	A	571	THR	2.5
1	A	589	ASP	2.4
1	A	245[A]	SER	2.2
1	A	87	GLU	2.2
1	A	17	LYS	2.1
1	A	459	ASN	2.0
1	A	31	GLU	2.0
1	A	467	THR	2.0
1	A	465	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	OAN	A	900	25/25	0.95	0.11	-2.98	34,41,52,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.