



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:04 PM GMT

PDB ID : 3W15  
Title : Structure of peroxisomal targeting signal 2 (PTS2) of *Saccharomyces cerevisiae* 3-ketoacyl-CoA thiolase in complex with Pex7p and Pex21p  
Authors : Pan, D.; Nakatsu, T.; Kato, H.  
Deposited on : 2012-11-06  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	<div> <div>11%</div> <div>83%</div> <div>8%</div> <div>9%</div> </div>
2	B	101	<div> <div>12%</div> <div>69%</div> <div>29%</div> </div>
3	C	389	<div> <div>%</div> <div>98%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NO3	A	401	-	X	-	-
4	NO3	A	402	-	X	-	-
4	NO3	B	301	-	X	-	-
4	NO3	C	401	-	X	-	-
4	NO3	C	402	-	X	-	X
4	NO3	C	403	-	X	-	-
4	NO3	C	404	-	X	-	X
4	NO3	C	405	-	X	-	X
4	NO3	C	406	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisomal targeting signal 2 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	7	0
			2650	1688	455	492	15			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP P39108
A	0	SER	-	EXPRESSION TAG	UNP P39108
A	?	-	GLU	DELETION	UNP P39108
A	?	-	SER	DELETION	UNP P39108
A	?	-	ALA	DELETION	UNP P39108
A	?	-	THR	DELETION	UNP P39108
A	?	-	ILE	DELETION	UNP P39108
A	?	-	LYS	DELETION	UNP P39108
A	?	-	ARG	DELETION	UNP P39108
A	?	-	THR	DELETION	UNP P39108
A	?	-	VAL	DELETION	UNP P39108

- Molecule 2 is a protein called Peroxisomal membrane protein PEX21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	72	Total	C	N	O	S	0	0	0
			548	338	93	114	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	188	GLY	-	EXPRESSION TAG	UNP P50091
B	189	SER	-	EXPRESSION TAG	UNP P50091


- Molecule 3 is a protein called 3-ketoacyl-CoA thiolase, peroxisomal, Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	389	Total	C	N	O	S	0	4	0
			3021	1939	491	584	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	EXPRESSION TAG	UNP P27796
C	0	SER	-	EXPRESSION TAG	UNP P27796
C	16	ARG	-	LINKER	UNP P0AEX9
C	17	SER	-	LINKER	UNP P0AEX9

- Molecule 4 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	C	5	Total	Mg	0	0
			5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	136	Total	O	0	0
			136	136		
6	B	28	Total	O	0	0
			28	28		
6	C	399	Total	O	0	0
			399	399		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peroxisomal targeting signal 2 receptor



- Molecule 2: Peroxisomal membrane protein PEX21



- Molecule 3: 3-ketoacyl-CoA thiolase, peroxisomal, Maltose-binding periplasmic protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.89Å 52.75Å 97.57Å 90.00° 106.59° 90.00°	Depositor
Resolution (Å)	32.20 – 1.80 32.01 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (32.20-1.80) 99.7 (32.01-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.94 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.189 , 0.224 0.188 , 0.225	Depositor DCC
$R_{free}$ test set	3618 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 46.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 71446 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6824	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/2738	0.58	1/3725 (0.0%)
2	B	0.38	0/550	0.49	0/734
3	C	0.40	0/3103	0.52	0/4214
All	All	0.40	0/6391	0.54	1/8673 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121	LEU	CA-CB-CG	5.33	127.56	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2507	17	0
2	B	548	0	536	1	0
3	C	3021	0	2975	5	0
4	A	8	0	0	0	0
4	B	4	0	0	0	0
4	C	24	0	0	0	0
5	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	5	0	0	0	0
6	A	136	0	0	0	0
6	B	28	0	0	0	0
6	C	399	0	0	0	0
All	All	6824	0	6018	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:ASP:OD2	1:A:251:ARG:HG2	1.83	0.79
1:A:116:ASN:HD22	1:A:187:GLN:HE21	1.46	0.64
1:A:149:GLU:O	1:A:153:MET:HG2	2.07	0.55
1:A:113:ASN:O	1:A:117:ARG:HA	2.09	0.52
2:B:267:ASN:HB3	2:B:269:GLU:HG2	1.94	0.50
1:A:43:LEU:HD12	1:A:368:LEU:HD22	1.94	0.49
1:A:185:HIS:CD2	1:A:233:PRO:HA	2.47	0.49
1:A:219:SER:HB3	3:C:67:VAL:HB	1.95	0.48
1:A:308[A]:CYS:SG	1:A:345:VAL:HG11	2.54	0.48
1:A:121:LEU:CD1	1:A:131:ILE:HG12	2.44	0.48
1:A:155:ASP:O	1:A:158:ASN:HB2	2.13	0.48
3:C:95:GLU:HG3	3:C:119:LYS:HB3	1.99	0.45
1:A:234:TYR:O	1:A:250:ILE:HG12	2.18	0.43
3:C:308[B]:GLU:OE2	3:C:312:LYS:HD3	2.19	0.43
3:C:246:PRO:HA	3:C:249:TRP:CE2	2.55	0.41
1:A:121:LEU:HD21	1:A:190:VAL:HG13	2.02	0.41
1:A:193[A]:CYS:SG	1:A:226:CYS:HB3	2.60	0.41
1:A:312:ARG:HB2	1:A:335:ILE:HD11	2.02	0.41
1:A:2:LEU:HB2	1:A:370:VAL:HB	2.03	0.41
1:A:127:GLY:HA2	1:A:175:ASN:O	2.22	0.40
3:C:188:TYR:OH	3:C:191:GLY:HA2	2.21	0.40
1:A:121:LEU:HD13	1:A:131:ILE:HG12	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/368 (91%)	327 (98%)	8 (2%)	0	100	100
2	B	66/101 (65%)	63 (96%)	3 (4%)	0	100	100
3	C	391/389 (100%)	386 (99%)	5 (1%)	0	100	100
All	All	792/858 (92%)	776 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/320 (90%)	285 (99%)	2 (1%)	88	86
2	B	65/92 (71%)	65 (100%)	0	100	100
3	C	313/315 (99%)	313 (100%)	0	100	100
All	All	665/727 (92%)	663 (100%)	2 (0%)	94	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	LEU
1	A	197	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NO3	A	401	-	3,3,3	3.27	3 (100%)	3,3,3	0.05	0
4	NO3	A	402	-	3,3,3	3.27	3 (100%)	3,3,3	0.15	0
4	NO3	B	301	-	3,3,3	3.20	3 (100%)	3,3,3	0.12	0
4	NO3	C	401	-	3,3,3	3.14	3 (100%)	3,3,3	0.25	0
4	NO3	C	402	-	3,3,3	3.26	3 (100%)	3,3,3	0.27	0
4	NO3	C	403	-	3,3,3	3.25	3 (100%)	3,3,3	0.13	0
4	NO3	C	404	-	3,3,3	3.23	3 (100%)	3,3,3	0.11	0
4	NO3	C	405	-	3,3,3	3.23	3 (100%)	3,3,3	0.39	0
4	NO3	C	406	-	3,3,3	3.26	3 (100%)	3,3,3	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NO3	A	401	-	-	0/0/0/0	0/0/0/0
4	NO3	A	402	-	-	0/0/0/0	0/0/0/0
4	NO3	B	301	-	-	0/0/0/0	0/0/0/0
4	NO3	C	401	-	-	0/0/0/0	0/0/0/0
4	NO3	C	402	-	-	0/0/0/0	0/0/0/0
4	NO3	C	403	-	-	0/0/0/0	0/0/0/0
4	NO3	C	404	-	-	0/0/0/0	0/0/0/0
4	NO3	C	405	-	-	0/0/0/0	0/0/0/0
4	NO3	C	406	-	-	0/0/0/0	0/0/0/0

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	401	NO3	O2-N	2.83	1.39	1.25
4	C	403	NO3	O3-N	2.83	1.39	1.25
4	C	404	NO3	O3-N	2.83	1.40	1.25
4	B	301	NO3	O3-N	2.85	1.40	1.25
4	C	403	NO3	O2-N	2.87	1.40	1.25
4	C	401	NO3	O3-N	2.87	1.40	1.25
4	C	402	NO3	O2-N	2.88	1.40	1.25
4	A	401	NO3	O3-N	2.88	1.40	1.25
4	A	401	NO3	O2-N	2.89	1.40	1.25
4	B	301	NO3	O2-N	2.91	1.40	1.25
4	C	406	NO3	O3-N	2.92	1.40	1.25
4	A	402	NO3	O2-N	2.92	1.40	1.25
4	A	402	NO3	O3-N	2.93	1.40	1.25
4	C	406	NO3	O2-N	2.94	1.40	1.25
4	C	404	NO3	O2-N	2.94	1.40	1.25
4	C	405	NO3	O2-N	2.97	1.40	1.25
4	C	402	NO3	O3-N	2.98	1.40	1.25
4	C	405	NO3	O3-N	3.00	1.40	1.25
4	C	401	NO3	O1-N	3.64	1.39	1.24
4	C	405	NO3	O1-N	3.67	1.39	1.24
4	B	301	NO3	O1-N	3.76	1.39	1.24
4	C	404	NO3	O1-N	3.83	1.40	1.24
4	C	406	NO3	O1-N	3.84	1.40	1.24
4	C	402	NO3	O1-N	3.84	1.40	1.24
4	A	402	NO3	O1-N	3.87	1.40	1.24
4	A	401	NO3	O1-N	3.93	1.40	1.24
4	C	403	NO3	O1-N	3.94	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/368 (91%)	0.59	42 (12%) 5 4	10, 20, 45, 49	0
2	B	72/101 (71%)	0.82	12 (16%) 2 1	15, 36, 47, 49	0
3	C	389/389 (100%)	-0.33	4 (1%) 84 82	9, 16, 27, 33	0
All	All	797/858 (92%)	0.16	58 (7%) 18 14	9, 18, 42, 49	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	151	THR	8.6
1	A	154	VAL	6.9
1	A	354	GLY	6.4
1	A	49	GLY	5.8
1	A	153	MET	5.7
1	A	273	SER	5.7
1	A	328	THR	5.6
1	A	1	MET	5.5
1	A	48	SER	5.4
1	A	155	ASP	5.1
1	A	47	ARG	5.0
2	B	197	SER	4.8
2	B	266	ASN	4.2
1	A	148	LEU	4.2
1	A	330	ALA	4.2
2	B	263	PHE	4.1
1	A	252	MET	4.1
2	B	224	SER	4.1
1	A	353	TRP	4.0
1	A	150	ILE	3.9
1	A	161	ILE	3.9
1	A	159	ALA	3.8
1	A	274	CYS	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	327	SER	3.8
1	A	158	ASN	3.6
1	A	156	PRO	3.6
1	A	331	THR	3.6
2	B	249	THR	3.5
1	A	356	PRO	3.5
1	A	325	THR	3.4
1	A	275	ILE	3.3
1	A	374	LEU	3.2
1	A	149	GLU	3.1
1	A	355	LYS	3.0
3	C	-1	GLY	3.0
1	A	162	LEU	3.0
2	B	250	LEU	2.9
2	B	267	ASN	2.9
2	B	269	GLU	2.8
1	A	324	LYS	2.8
2	B	210	CYS	2.8
1	A	152	LYS	2.8
1	A	352	LEU	2.7
1	A	21	ASP	2.7
2	B	268	GLY	2.7
1	A	251	ARG	2.5
1	A	163	LYS	2.5
3	C	117	ASN	2.5
1	A	147	PRO	2.3
1	A	213	ASN	2.3
3	C	190	ASN	2.2
1	A	181[A]	GLN	2.2
1	A	45	ILE	2.2
1	A	135	LEU	2.2
2	B	225	SER	2.1
2	B	198	GLU	2.1
1	A	46	ASP	2.1
3	C	191	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NO3	C	402	4/4	0.88	0.22	13.47	31,32,33,33	0
4	NO3	C	405	4/4	0.84	0.20	11.52	30,30,30,30	0
4	NO3	C	404	4/4	0.84	0.24	6.12	36,36,36,37	0
4	NO3	B	301	4/4	0.94	0.09	0.88	28,28,28,28	0
4	NO3	A	401	4/4	0.85	0.18	0.28	40,40,40,41	0
4	NO3	C	403	4/4	0.94	0.10	0.13	26,27,27,27	0
4	NO3	C	401	4/4	0.95	0.09	-0.33	25,26,26,27	0
5	MG	A	403	1/1	0.99	0.07	-2.47	22,22,22,22	0
4	NO3	C	406	4/4	0.84	0.17	-	47,47,47,47	0
5	MG	C	409	1/1	0.97	0.14	-	24,24,24,24	0
4	NO3	A	402	4/4	0.95	0.24	-	48,48,48,48	0
5	MG	C	410	1/1	0.98	0.06	-	18,18,18,18	0
5	MG	C	408	1/1	0.93	0.05	-	31,31,31,31	0
5	MG	C	407	1/1	0.99	0.12	-	19,19,19,19	0
5	MG	C	411	1/1	0.94	0.09	-	39,39,39,39	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.