



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:08 AM GMT

PDB ID : 2W2X  
Title : COMPLEX OF RAC2 AND PLCG2 SPPH DOMAIN  
Authors : Opaleye, O.; Bunney, T.D.; Roe, S.M.; Pearl, L.H.  
Deposited on : 2008-11-04  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	<div> <div>3%</div> <div> <div></div> <div>53%</div> <div>31%</div> <div>10%</div> <div>• •</div> </div> </div>
1	B	185	<div> <div>3%</div> <div> <div></div> <div>59%</div> <div>25%</div> <div>8%</div> <div>8%</div> </div> </div>
2	C	124	<div> <div>2%</div> <div> <div></div> <div>40%</div> <div>36%</div> <div>10%</div> <div>•</div> <div>12%</div> </div> </div>
3	D	124	<div> <div>3%</div> <div> <div></div> <div>55%</div> <div>21%</div> <div>6%</div> <div>•</div> <div>15%</div> </div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAS-RELATED C3 BOTULINUM TOXIN SUBSTRATE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	177	Total	C	N	O	S	0	0	0
			1333	859	216	252	6			
1	B	170	Total	C	N	O	S	0	0	0
			1250	806	200	239	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	VAL	GLY	ENGINEERED MUTATION	UNP P15153
B	12	VAL	GLY	ENGINEERED MUTATION	UNP P15153

- Molecule 2 is a protein called 1-PHOSPHATIDYLINOSITOL-4,5-BISPHOSPHATE PHOSPHODIESTERASE GAMMA-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	109	Total	C	N	O	S	0	0	0
			860	553	143	160	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	26	PHE	TYR	CONFLICT	UNP P16885
C	88	ASP	TYR	CONFLICT	UNP P16885


- Molecule 3 is a protein called 1-PHOSPHATIDYLINOSITOL-4,5-BISPHOSPHATE PHOSPHODIESTERASE GAMMA-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	105	Total	C	N	O	S	0	0	0
			834	536	136	158	4			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	26	PHE	TYR	CONFLICT	UNP P16885
D	88	ASP	TYR	CONFLICT	UNP P16885
D	97	LYS	ARG	CONFLICT	UNP P16885

- Molecule 4 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		
4	B	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	12	Total	O	0	0
			12	12		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	11	Total 11	O 11	0	0
6	C	6	Total 6	O 6	0	0
6	D	8	Total 8	O 8	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.57Å 84.46Å 74.20Å 90.00° 112.21° 90.00°	Depositor
Resolution (Å)	69.41 – 2.30 68.69 – 2.30	Depositor EDS
% Data completeness (in resolution range)	83.5 (69.41-2.30) 80.0 (68.69-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.24 (at 2.29Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.234 , 0.294 0.221 , 0.280	Depositor DCC
$R_{free}$ test set	2653 reflections (5.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.3	Xtriage
Anisotropy	0.818	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 66.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 26827 reflections (0.007%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4380	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	9.42	18/1358 (1.3%)	1.85	29/1856 (1.6%)
1	B	7.19	7/1276 (0.5%)	1.58	13/1753 (0.7%)
2	C	8.69	13/875 (1.5%)	1.64	19/1178 (1.6%)
3	D	6.68	8/851 (0.9%)	1.33	11/1150 (1.0%)
All	All	8.16	46/4360 (1.1%)	1.64	72/5937 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

The worst 5 of 46 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	83	SER	CA-CB	159.79	3.92	1.52
1	B	44	VAL	C-O	159.16	4.25	1.23
1	B	51	VAL	C-O	134.19	3.78	1.23
1	A	48	SER	C-O	129.12	3.68	1.23
1	B	35	THR	C-O	113.69	3.39	1.23

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	SER	CA-CB-OG	-34.16	18.98	111.20
1	B	51	VAL	CA-C-O	-31.50	53.94	120.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	44	VAL	CA-C-O	-31.48	53.99	120.10
1	A	86	SER	N-CA-CB	29.93	155.40	110.50
1	A	86	SER	CA-CB-OG	-20.70	55.30	111.20

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	30	GLY	Peptide
1	A	31	GLU	Peptide
1	B	30	GLY	Peptide
1	B	31	GLU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1333	0	1282	91	0
1	B	1250	0	1164	54	0
2	C	860	0	783	60	0
3	D	834	0	760	41	0
4	A	32	0	12	3	0
4	B	32	0	12	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	12	0	0	0	0
6	B	11	0	0	0	0
6	C	6	0	0	0	0
6	D	8	0	0	0	0
All	All	4380	0	4013	243	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 243 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:52:SER:C	2:C:53:LEU:N	1.84	1.30
2:C:33:LYS:C	2:C:34:LEU:N	1.86	1.28
1:A:83:SER:C	1:A:84:LEU:N	1.87	1.27
2:C:103:GLU:C	2:C:104:TRP:N	1.87	1.27
1:A:130:LYS:C	1:A:131:GLU:N	1.91	1.23

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/185 (89%)	148 (90%)	14 (8%)	3 (2%)	11	9
1	B	163/185 (88%)	151 (93%)	11 (7%)	1 (1%)	30	36
2	C	93/124 (75%)	86 (92%)	4 (4%)	3 (3%)	5	3
3	D	93/124 (75%)	84 (90%)	3 (3%)	6 (6%)	1	0
All	All	514/618 (83%)	469 (91%)	32 (6%)	13 (2%)	7	5

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	31	GLU
2	C	47	ASP
2	C	114	LYS
3	D	50	LEU
1	B	26	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/159 (89%)	129 (92%)	12 (8%)	13	16
1	B	127/159 (80%)	119 (94%)	8 (6%)	22	29
2	C	87/109 (80%)	77 (88%)	10 (12%)	7	7
3	D	86/109 (79%)	78 (91%)	8 (9%)	11	13
All	All	441/536 (82%)	403 (91%)	38 (9%)	13	15

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	120	ARG
2	C	31	ASP
3	D	70	GLN
1	B	134	LEU
2	C	33	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	162	GLN
2	C	8	GLN
3	D	8	GLN
1	B	26	ASN
2	C	70	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GSP	A	1178	5	25,34,34	2.23	2 (8%)	31,54,54	2.12	7 (22%)
4	GSP	B	1178	5	25,34,34	2.31	3 (12%)	31,54,54	1.74	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GSP	A	1178	5	-	0/15/38/38	0/3/3/3
4	GSP	B	1178	5	-	0/15/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1178	GSP	PG-S1G	-10.36	1.70	1.90
4	A	1178	GSP	PG-S1G	-9.91	1.71	1.90
4	B	1178	GSP	C2-N1	2.00	1.38	1.35
4	A	1178	GSP	C6-N1	3.05	1.38	1.33
4	B	1178	GSP	C6-N1	3.08	1.38	1.33

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1178	GSP	N3-C2-N1	-5.49	119.08	127.44
4	A	1178	GSP	PA-O3A-PB	-4.99	118.73	132.73
4	A	1178	GSP	C2'-C1'-N9	-4.89	106.83	114.29
4	B	1178	GSP	N3-C2-N1	-4.84	120.07	127.44
4	B	1178	GSP	C5-C6-N1	-3.04	119.43	123.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1178	GSP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	177/185 (95%)	0.35	5 (2%) 56 66	34, 58, 102, 131	8 (4%)
1	B	170/185 (91%)	0.26	5 (2%) 55 64	41, 63, 105, 163	2 (1%)
2	C	109/124 (87%)	0.29	3 (2%) 56 66	44, 63, 96, 109	0
3	D	105/124 (84%)	0.29	4 (3%) 44 53	34, 60, 96, 131	0
All	All	561/618 (90%)	0.30	17 (3%) 54 63	34, 62, 104, 163	10 (1%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	51	VAL	8.5
1	A	46	VAL	5.3
1	A	1	SER	4.7
2	C	43	THR	4.5
2	C	113	TRP	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GSP	B	1178	32/32	0.98	0.14	-0.64	32,60,83,120	4
4	GSP	A	1178	32/32	0.98	0.14	-0.64	33,58,91,107	0
5	MG	B	1179	1/1	0.89	0.19	-	46,46,46,46	0
5	MG	A	1179	1/1	0.92	0.13	-	47,47,47,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.