



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:59 PM GMT

PDB ID : 1W4L  
Title : COMPLEX OF TCACHE WITH BIS-ACTING GALANTHAMINE  
DERIVATIVE  
Authors : Greenblatt, H.M.; Guillou, C.; Guenard, D.; Badet, B.; Thal, C.; Silman, I.;  
Sussman, J.L.  
Deposited on : 2004-07-25  
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	<div> <div>4%</div> <div>81%</div> <div>13%</div> <div>...</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4608 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLCHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	529	Total	C	N	O	S	0	0	0
			4204	2699	713	770	22			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is GALANTHAMINE DERIVATIVE (three-letter code: GL8) (formula: C<sub>32</sub>H<sub>37</sub>N<sub>2</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			39	32	2	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	309	Total	O	0	0
			309	309		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ACETYLCHOLINESTERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.33Å 111.33Å 136.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.41 – 2.16 39.41 – 2.16	Depositor EDS
% Data completeness (in resolution range)	98.6 (39.41-2.16) 98.7 (39.41-2.16)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.00 (at 2.16Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.184 , 0.218 0.188 , 0.223	Depositor DCC
$R_{free}$ test set	2621 reflections (4.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtriage
Anisotropy	0.669	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 50.3	EDS
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 52748 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4608	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GL8, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	3/4325 (0.1%)	0.94	10/5873 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	175	MET	SD-CE	-8.15	1.32	1.77
1	A	82	GLU	CB-CG	6.45	1.64	1.52
1	A	82	GLU	CG-CD	-5.90	1.43	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	ARG	NE-CZ-NH2	-11.89	114.35	120.30
1	A	243	ARG	NE-CZ-NH1	8.19	124.40	120.30
1	A	518	VAL	CB-CA-C	-7.66	96.86	111.40
1	A	22	VAL	CB-CA-C	-6.86	98.36	111.40
1	A	129	VAL	CB-CA-C	-6.62	98.83	111.40
1	A	143	LEU	CA-CB-CG	-6.11	101.24	115.30
1	A	349	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	A	243	ARG	CG-CD-NE	-5.57	100.10	111.80
1	A	221	ARG	N-CA-CB	-5.54	100.63	110.60
1	A	518	VAL	CG1-CB-CG2	5.14	119.12	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	442	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4204	0	4053	84	0
2	A	28	0	26	0	0
3	A	28	0	25	0	0
4	A	39	0	37	0	0
5	A	309	0	0	11	1
All	All	4608	0	4141	84	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:LEU:HG	5:A:2272:HOH:O	1.24	1.29
1:A:468:ARG:HB2	5:A:2277:HOH:O	1.50	1.11
1:A:194:VAL:HG12	5:A:2127:HOH:O	1.55	1.05
1:A:468:ARG:HD3	5:A:2274:HOH:O	1.73	0.88
1:A:22:VAL:HG13	1:A:133:LYS:HG3	1.62	0.82
1:A:366:LEU:HD23	1:A:531:LEU:HD11	1.60	0.81
1:A:194:VAL:CG1	1:A:219:PHE:HA	2.14	0.77
1:A:468:ARG:NE	5:A:2277:HOH:O	2.17	0.74
1:A:453:VAL:HG22	1:A:456:LEU:HG	1.68	0.74
1:A:346:LYS:HG2	1:A:385:ILE:HD13	1.71	0.73
1:A:366:LEU:CD2	1:A:531:LEU:HD11	2.20	0.71
1:A:36:ALA:HB2	1:A:175:MET:HE2	1.72	0.71
1:A:194:VAL:HG13	1:A:219:PHE:HA	1.75	0.69
1:A:373:LEU:HD12	1:A:378:TRP:HE1	1.59	0.68
1:A:453:VAL:CG2	1:A:456:LEU:HG	2.24	0.67

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:MET:HE3	5:A:2018:HOH:O	1.94	0.66
1:A:36:ALA:HB2	1:A:175:MET:CE	2.26	0.65
1:A:452:LEU:HD21	1:A:467:ARG:NE	2.11	0.65
1:A:452:LEU:HD23	1:A:463:GLU:HG3	1.80	0.64
1:A:452:LEU:HD22	1:A:467:ARG:HH21	1.63	0.63
1:A:452:LEU:HD22	1:A:467:ARG:NH2	2.14	0.62
1:A:383:ASN:HD22	1:A:383:ASN:C	2.02	0.62
1:A:243:ARG:NH2	5:A:2155:HOH:O	2.32	0.61
1:A:489:GLU:O	1:A:491:LYS:HE2	2.02	0.59
1:A:366:LEU:HD23	1:A:535:THR:HG21	1.85	0.58
1:A:19:ARG:NH2	1:A:26:HIS:HB2	2.20	0.57
1:A:316:LYS:NZ	5:A:2201:HOH:O	2.36	0.57
1:A:221:ARG:HG2	1:A:476:PHE:CE2	2.41	0.56
1:A:373:LEU:CD1	1:A:378:TRP:HE1	2.19	0.55
1:A:499:GLU:HG2	1:A:501:LYS:HE3	1.89	0.55
1:A:36:ALA:CB	1:A:175:MET:CE	2.86	0.54
1:A:50:PRO:HA	1:A:175:MET:HE3	1.88	0.54
1:A:46:ARG:NE	1:A:163:GLU:OE2	2.41	0.53
1:A:9:ASN:ND2	1:A:14:LYS:HE2	2.23	0.53
1:A:452:LEU:HD21	1:A:467:ARG:CZ	2.38	0.53
1:A:82:GLU:HG3	1:A:85:ASN:HD22	1.73	0.52
1:A:346:LYS:HG2	1:A:385:ILE:CD1	2.39	0.52
1:A:452:LEU:CD2	1:A:467:ARG:NH2	2.72	0.52
1:A:449:GLY:HA2	1:A:466:SER:OG	2.11	0.51
1:A:197:PHE:CB	1:A:223:ILE:HB	2.41	0.50
1:A:464:ALA:O	1:A:468:ARG:HG2	2.11	0.50
1:A:9:ASN:ND2	1:A:14:LYS:HG2	2.26	0.50
1:A:47:ARG:HG3	5:A:2112:HOH:O	2.11	0.49
1:A:366:LEU:CD2	1:A:535:THR:HG21	2.42	0.49
1:A:320:LEU:HD23	1:A:320:LEU:C	2.33	0.49
1:A:135:LEU:HD23	1:A:143:LEU:HD13	1.95	0.49
1:A:461:GLU:CD	1:A:461:GLU:H	2.17	0.48
1:A:197:PHE:HB2	1:A:223:ILE:HB	1.95	0.48
1:A:50:PRO:O	1:A:52:LYS:HE3	2.13	0.48
1:A:146:LEU:C	1:A:146:LEU:HD12	2.35	0.47
1:A:221:ARG:HD3	1:A:476:PHE:O	2.14	0.47
1:A:450:LEU:N	1:A:451:PRO:CD	2.78	0.46
1:A:132:GLY:HA3	1:A:143:LEU:HD22	1.98	0.46
1:A:450:LEU:HD12	1:A:456:LEU:HD11	1.98	0.46
1:A:405:MET:HA	1:A:408:VAL:HG12	1.98	0.45
1:A:19:ARG:NE	1:A:26:HIS:HB3	2.31	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:420:LEU:HD23	1:A:502:PHE:HB3	1.99	0.45
1:A:506:ASN:OD1	1:A:508:GLU:HG2	2.17	0.45
1:A:228:SER:HB2	1:A:229:PRO:HD2	1.98	0.45
1:A:19:ARG:CZ	1:A:26:HIS:HB2	2.47	0.44
1:A:451:PRO:HA	1:A:458:TYR:CD1	2.52	0.44
1:A:499:GLU:CG	1:A:501:LYS:HE3	2.47	0.44
1:A:223:ILE:HA	1:A:320:LEU:O	2.18	0.44
1:A:349:ARG:NH2	1:A:376:THR:OG1	2.38	0.44
1:A:129:VAL:HG22	5:A:2053:HOH:O	2.18	0.44
1:A:499:GLU:HB3	1:A:501:LYS:HG3	2.00	0.43
1:A:349:ARG:HA	1:A:349:ARG:HD2	1.78	0.43
1:A:426:ARG:CZ	1:A:430:LEU:HD12	2.49	0.42
1:A:173:GLN:OE1	1:A:205:SER:HB3	2.19	0.42
1:A:264:HIS:CD2	5:A:2164:HOH:O	2.71	0.42
1:A:531:LEU:O	1:A:531:LEU:HD12	2.20	0.41
1:A:194:VAL:HG13	1:A:219:PHE:CA	2.48	0.41
1:A:450:LEU:HD12	1:A:456:LEU:CD1	2.49	0.41
1:A:491:LYS:HA	1:A:491:LYS:HD3	1.92	0.41
1:A:19:ARG:HE	1:A:26:HIS:HB3	1.84	0.41
1:A:439:ILE:HB	1:A:442:TYR:CD1	2.56	0.41
1:A:455:GLU:CD	1:A:455:GLU:H	2.24	0.41
1:A:452:LEU:HA	1:A:452:LEU:HD23	1.81	0.41
1:A:383:ASN:C	1:A:383:ASN:ND2	2.74	0.41
1:A:185:GLN:HA	1:A:189:GLY:O	2.20	0.41
1:A:22:VAL:CG1	1:A:133:LYS:HG3	2.42	0.41
1:A:211:LEU:HD23	1:A:314:PHE:HB3	2.04	0.40
1:A:408:VAL:CG2	1:A:418:THR:HG21	2.51	0.40
1:A:19:ARG:NE	1:A:26:HIS:CB	2.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2002:HOH:O	5:A:2182:HOH:O[3_564]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/543 (97%)	503 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	458/474 (97%)	437 (95%)	21 (5%)	33	29

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	15	VAL
1	A	22	VAL
1	A	52	LYS
1	A	82	GLU
1	A	129	VAL
1	A	143	LEU
1	A	243	ARG
1	A	274	LEU
1	A	287	ILE
1	A	288	PHE
1	A	294	PRO
1	A	330	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	340	SER
1	A	358	LEU
1	A	383	ASN
1	A	430	LEU
1	A	468	ARG
1	A	473	TRP
1	A	499	GLU
1	A	518	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	68	GLN
1	A	257	ASN
1	A	383	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	1537	1,3	14,14,15	0.78	0	15,19,21	0.80	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	1538	3	14,14,15	0.66	0	15,19,21	0.80	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1537	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1538	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1538	NAG	C2-N2-C7	-2.45	119.89	123.04
3	A	1537	NAG	C2-N2-C7	-2.25	120.15	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	1536	1	14,14,15	0.81	0	15,19,21	1.18	2 (13%)
2	NAG	A	1539	1	14,14,15	1.17	1 (7%)	15,19,21	1.16	1 (6%)
4	GL8	A	1540	-	43,44,44	3.42	21 (48%)	56,64,64	2.76	16 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1536	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1539	1	-	0/6/23/26	0/1/1/1
4	GL8	A	1540	-	-	0/12/65/65	0/5/6/6

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1540	GL8	C26-C25	-6.05	1.39	1.48
4	A	1540	GL8	C31-C32	-5.96	1.39	1.48
4	A	1540	GL8	C25-N2	-5.31	1.32	1.39
4	A	1540	GL8	C32-N2	-5.07	1.33	1.39
4	A	1540	GL8	C6-C14	-4.82	1.47	1.52
4	A	1540	GL8	O1-C5	-3.67	1.42	1.47
4	A	1540	GL8	C8-C7	2.02	1.43	1.39
4	A	1540	GL8	C9-C15	2.06	1.44	1.41
4	A	1540	GL8	C29-C30	2.14	1.43	1.38
4	A	1540	GL8	C13-C14	2.53	1.42	1.38
4	A	1540	GL8	C4-C5	2.58	1.57	1.52
4	A	1540	GL8	C28-C27	2.68	1.44	1.38
4	A	1540	GL8	C12-C11	2.74	1.56	1.52
4	A	1540	GL8	O2-C7	2.76	1.41	1.37
4	A	1540	GL8	O3-C3	2.77	1.49	1.43
4	A	1540	GL8	O32-C32	2.88	1.28	1.22
4	A	1540	GL8	O25-C25	3.16	1.29	1.22
2	A	1539	NAG	C1-C2	3.32	1.57	1.52
4	A	1540	GL8	C12-C6	3.65	1.58	1.54
4	A	1540	GL8	C3-C2	4.20	1.55	1.49
4	A	1540	GL8	C6-C1	5.80	1.58	1.51
4	A	1540	GL8	C4-C3	13.10	1.66	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1540	GL8	C14-C6-C5	-9.09	94.89	99.97
4	A	1540	GL8	O3-C3-C2	-7.37	97.91	109.94
4	A	1540	GL8	C4-C5-C6	-7.30	110.50	115.22
4	A	1540	GL8	C13-O1-C5	-4.48	98.97	105.34
4	A	1540	GL8	C5-C4-C3	-4.25	107.17	113.16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1540	GL8	C12-C6-C1	-3.44	107.89	112.21
4	A	1540	GL8	C3-C2-C1	-3.22	121.12	124.14
2	A	1539	NAG	C3-C4-C5	-2.84	105.25	110.20
4	A	1540	GL8	C22-C23-C24	-2.82	100.01	113.27
2	A	1536	NAG	C4-C3-C2	-2.72	107.00	111.23
2	A	1536	NAG	C2-N2-C7	-2.43	119.92	123.04
4	A	1540	GL8	C12-C6-C14	2.44	118.31	114.51
4	A	1540	GL8	O32-C32-N2	2.47	127.44	124.78
4	A	1540	GL8	C6-C14-C13	2.49	109.05	107.36
4	A	1540	GL8	O1-C5-C4	2.66	112.42	109.72
4	A	1540	GL8	C12-C11-N1	2.74	118.65	113.06
4	A	1540	GL8	C12-C6-C5	3.92	116.84	110.68
4	A	1540	GL8	C23-C24-N2	3.95	121.00	112.41
4	A	1540	GL8	O1-C5-C6	7.65	110.96	105.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	529/543 (97%)	-0.11	21 (3%)	42 52	21, 31, 45, 63	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	535	THR	5.4
1	A	485	PRO	4.5
1	A	55	SER	3.0
1	A	489	GLU	2.9
1	A	201	ALA	2.8
1	A	146	LEU	2.7
1	A	534	ALA	2.7
1	A	400	VAL	2.6
1	A	120	PHE	2.5
1	A	115	ILE	2.5
1	A	148	TYR	2.4
1	A	118	GLY	2.4
1	A	533	ASN	2.3
1	A	206	VAL	2.2
1	A	200	SER	2.2
1	A	114	TRP	2.2
1	A	116	TYR	2.1
1	A	224	LEU	2.1
1	A	203	GLY	2.0
1	A	508	GLU	2.0
1	A	196	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	1538	14/15	0.82	0.38	-	59,61,64,64	0
3	NAG	A	1537	14/15	0.91	0.17	-	40,43,50,56	0

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GL8	A	1540	39/39	0.94	0.18	0.39	22,25,39,43	0
2	NAG	A	1539	14/15	0.84	0.40	-	61,65,67,70	0
2	NAG	A	1536	14/15	0.79	0.21	-	57,60,63,66	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.