



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:13 AM GMT

PDB ID : 2WBG  
Title : Structure of family 1 beta-glucosidase from *Thermotoga maritima* in complex with 3-imino-2-oxa-(+)-castanospermine  
Authors : Aguilar, M.; Gloster, T.M.; Turkenburg, J.P.; Garcia-Moreno, M.I.; Ortiz Mellet, C.; Davies, G.J.; Garcia Fernandez, J.M.  
Deposited on : 2009-02-27  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 0%, yellow 0%, green 89%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>89%</span> <span>5% • 5%</span> </div> </div>
1	B	468	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 0%, yellow 0%, green 87%, grey 7%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>87%</span> <span>7% 5%</span> </div> </div>
1	C	468	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 0%, yellow 0%, green 90%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>90%</span> <span>5% 5%</span> </div> </div>
1	D	468	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 0%, yellow 0%, green 87%, grey 7%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>87%</span> <span>7% • 5%</span> </div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 16641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-GLUCOSIDASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	18	0
			3709	2421	619	663	6			
1	B	443	Total	C	N	O	S	0	9	0
			3647	2382	603	655	7			
1	C	443	Total	C	N	O	S	0	22	0
			3755	2449	625	675	6			
1	D	443	Total	C	N	O	S	0	15	0
			3671	2393	606	666	6			

- Molecule 2 is (3Z,5S,6R,7S,8R,8AR)-3-(OCTYLIMINO)HEXAHYDRO[1,3]OXAZOLO[3,4-A]PYRIDINE-5,6,7,8-TETROL (three-letter code: LGS) (formula: C<sub>15</sub>H<sub>28</sub>N<sub>2</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			22	15	2	5		
2	B	1	Total	C	N	O	0	0
			22	15	2	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			22	15	2	5		
2	D	1	Total	C	N	O	0	0
			22	15	2	5		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	411	Total	O	0	0
			411	411		
4	B	379	Total	O	0	0
			379	379		
4	C	522	Total	O	0	0
			522	522		
4	D	455	Total	O	0	0
			455	455		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: BETA-GLUCOSIDASE A




#### • Molecule 1: BETA-GLUCOSIDASE A




#### • Molecule 1: BETA-GLUCOSIDASE A



#### • Molecule 1: BETA-GLUCOSIDASE A





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.99Å 73.19Å 138.09Å 90.00° 94.03° 90.00°	Depositor
Resolution (Å)	137.36 – 1.85 19.96 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (137.36-1.85) 99.7 (19.96-1.85)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.18 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.4.0077	Depositor
R, $R_{free}$	0.153 , 0.199 0.153 , 0.199	Depositor DCC
$R_{free}$ test set	7664 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.3	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 60.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 152636 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16641	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.77 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4638e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LGS, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.75	0/3877	0.69	0/5269
1	B	0.68	0/3788	0.68	1/5150 (0.0%)
1	C	0.86	0/3931	0.74	0/5339
1	D	0.75	0/3832	0.72	1/5212 (0.0%)
All	All	0.76	0/15428	0.71	2/20970 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	269	ARG	NE-CZ-NH1	-6.88	116.86	120.30
1	D	77	ARG	NE-CZ-NH2	-6.19	117.20	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3709	0	3609	22	0
1	B	3647	0	3505	28	0
1	C	3755	0	3666	20	0
1	D	3671	0	3512	25	0
2	A	22	0	28	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	22	0	28	0	0
2	C	22	0	28	0	0
2	D	22	0	28	0	0
3	C	4	0	3	1	0
4	A	411	0	0	5	0
4	B	379	0	0	6	0
4	C	522	0	0	6	0
4	D	455	0	0	5	0
All	All	16641	0	14407	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:145[C]:GLU:HG3	4:D:2211:HOH:O	1.38	1.23
1:C:192[B]:ARG:CG	1:C:192[B]:ARG:HH11	1.54	1.20
1:C:148[C]:ARG:NH2	4:C:2237:HOH:O	1.73	1.18
1:C:192[B]:ARG:NH1	4:C:2272:HOH:O	1.66	1.12
1:C:192[B]:ARG:HG2	1:C:192[B]:ARG:NH1	1.50	1.08
1:B:185:ARG:HG3	1:B:185:ARG:HH11	1.20	1.06
1:B:216:LYS:HD3	4:B:2311:HOH:O	1.69	0.91
1:D:370:ASN:HB3	4:D:2401:HOH:O	1.72	0.89
1:C:192[B]:ARG:HG2	1:C:192[B]:ARG:HH11	0.74	0.89
1:B:185:ARG:CG	1:B:185:ARG:HH11	1.90	0.85
1:C:131[B]:LYS:NZ	4:C:2207:HOH:O	2.09	0.84
1:A:333:TRP:CZ2	1:A:337:LYS:HD2	2.17	0.79
1:A:144:ALA:HB2	1:A:203[A]:ARG:HG2	1.65	0.79
1:A:144:ALA:HB2	1:A:203[A]:ARG:CG	2.17	0.75
1:B:333:TRP:CZ2	1:B:337:LYS:HD3	2.23	0.74
1:C:357[A]:ASP:OD1	4:C:2425:HOH:O	2.06	0.72
1:B:246:ASN:HB3	4:B:2239:HOH:O	1.91	0.70
1:D:304:PRO:O	1:D:305:ASP:CB	2.41	0.67
1:B:17:ALA:HB3	1:B:20:GLN:HE21	1.60	0.67
1:A:343[A]:ASN:HB2	4:A:2331:HOH:O	1.94	0.66
1:A:141:ASP:OD1	1:A:203[A]:ARG:NH1	2.29	0.65
1:A:333:TRP:CZ2	1:A:337:LYS:CD	2.80	0.64
1:D:304:PRO:O	1:D:305:ASP:CG	2.36	0.64
1:D:304:PRO:O	1:D:305:ASP:OD1	2.15	0.63
1:B:144:ALA:HB2	1:B:203[A]:ARG:HG2	1.81	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:148[B]:ARG:HG3	1:C:149:VAL:N	2.14	0.63
1:D:370:ASN:CB	4:D:2401:HOH:O	2.39	0.61
1:B:185:ARG:CG	1:B:185:ARG:NH1	2.58	0.58
1:D:7:PRO:HG2	4:D:2417:HOH:O	2.02	0.58
1:C:246:ASN:OD1	3:C:1447:ACT:H1	2.04	0.58
1:D:302:PHE:CE2	1:D:304:PRO:HD3	2.40	0.56
1:D:213:LYS:O	1:D:214:ASP:HB2	2.06	0.55
1:D:5:LYS:HE2	1:D:442:ASN:HB3	1.89	0.55
1:D:383:TRP:O	1:D:387:GLN:HG2	2.07	0.55
1:A:88[B]:GLU:OE2	1:A:92:ARG:NH2	2.38	0.55
1:C:109[B]:LEU:HD11	1:C:157:ARG:HB3	1.89	0.54
1:A:144:ALA:HB2	1:A:203[A]:ARG:HG3	1.89	0.52
1:B:152[B]:GLU:OE1	1:B:152[B]:GLU:CA	2.57	0.52
1:C:274[A]:GLU:O	1:C:275[A]:ASN:HB2	2.10	0.51
1:A:262[A]:GLU:HG3	1:A:263:LEU:N	2.25	0.51
1:B:185:ARG:NH1	1:B:185:ARG:HG3	2.02	0.51
1:A:262[A]:GLU:HG3	1:A:263:LEU:HG	1.93	0.51
1:A:185:ARG:NH1	4:A:2205:HOH:O	2.43	0.51
1:D:236:ILE:O	1:D:240[A]:ARG:HG3	2.12	0.50
1:A:333:TRP:CE2	1:A:337:LYS:HD2	2.46	0.49
1:D:304:PRO:O	1:D:305:ASP:HB2	2.11	0.49
1:B:300:VAL:HG13	1:B:310[B]:VAL:HG23	1.93	0.49
1:C:300[A]:VAL:HG13	1:C:310:VAL:HG13	1.94	0.49
1:B:141:ASP:OD1	1:B:203[B]:ARG:NH2	2.46	0.48
1:A:74:LYS:HD3	4:A:2102:HOH:O	2.14	0.48
1:B:265:LEU:O	1:B:269:ARG:HB2	2.13	0.48
1:A:90:THR:HG22	4:A:2118:HOH:O	2.14	0.47
1:A:20:GLN:O	1:A:403:ASN:HB2	2.13	0.47
1:C:192[B]:ARG:CG	1:C:192[B]:ARG:NH1	2.28	0.46
1:B:144:ALA:HB2	1:B:203[A]:ARG:CG	2.45	0.46
1:B:141:ASP:OD1	1:B:203[A]:ARG:NH1	2.37	0.46
1:B:238:ALA:HB1	1:B:310[A]:VAL:HG12	1.96	0.46
1:B:275:ASN:HB2	4:B:2264:HOH:O	2.16	0.46
1:B:223:ASN:HA	1:B:247:ASN:OD1	2.16	0.46
1:B:121:HIS:O	1:B:122:TRP:HB2	2.16	0.45
1:B:26:LEU:HD21	1:B:32:MET:HG2	1.99	0.45
1:B:4:LYS:HE3	4:B:2004:HOH:O	2.16	0.45
1:B:339[B]:LYS:NZ	4:B:2302:HOH:O	2.49	0.45
1:D:216:LYS:HD2	4:D:2372:HOH:O	2.16	0.45
1:A:307:PRO:C	1:A:309:LYS:H	2.20	0.45
1:A:5:LYS:HE2	1:A:441:ASN:O	2.16	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:109[A]:LEU:HD13	1:C:109[A]:LEU:HA	1.68	0.44
1:A:262[A]:GLU:CG	1:A:263:LEU:N	2.80	0.44
1:B:303:ASP:O	1:B:309:LYS:HA	2.18	0.43
1:C:336:LYS:HE2	1:C:336:LYS:HB3	1.87	0.43
1:D:355:ALA:HA	1:D:413:ARG:O	2.18	0.43
1:B:336:LYS:HE2	1:B:336:LYS:HB3	1.91	0.43
1:A:316:ASP:OD1	4:A:2312:HOH:O	2.22	0.43
1:D:336:LYS:O	1:D:340[A]:GLU:HG3	2.20	0.42
1:D:82:TRP:HB3	1:D:83:PRO:HD3	2.02	0.41
1:C:46:LYS:HD3	4:C:2479:HOH:O	2.19	0.41
1:B:398:TRP:HA	1:B:399:SER:HA	1.85	0.41
1:C:274[A]:GLU:O	1:C:275[A]:ASN:CB	2.68	0.41
1:B:403:ASN:ND2	4:B:2343:HOH:O	2.50	0.41
1:A:68:ILE:HG23	1:A:73:VAL:HB	2.01	0.41
1:C:24:SER:N	1:C:25:PRO:CD	2.84	0.41
1:D:207:VAL:O	1:D:211:THR:HG23	2.20	0.41
1:A:300:VAL:HG13	1:A:310[A]:VAL:CG1	2.51	0.41
1:C:405:GLU:HG3	1:C:405:GLU:O	2.21	0.41
1:C:148[A]:ARG:NH2	4:C:2236:HOH:O	0.69	0.41
1:B:248:TYR:N	1:B:249:PRO:CD	2.84	0.40
1:D:28:ASP:HB3	1:D:96:LYS:HD3	2.03	0.40
1:D:300[A]:VAL:HG13	1:D:310:VAL:HG13	2.03	0.40
1:D:26:LEU:HD21	1:D:32:MET:HG2	2.03	0.40
1:D:307:PRO:O	1:D:308:ALA:HB3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	459/468 (98%)	451 (98%)	8 (2%)	0	<b>100</b> <b>100</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	450/468 (96%)	443 (98%)	7 (2%)	0	100	100
1	C	464/468 (99%)	461 (99%)	2 (0%)	1 (0%)	52	36
1	D	457/468 (98%)	453 (99%)	2 (0%)	2 (0%)	39	22
All	All	1830/1872 (98%)	1808 (99%)	19 (1%)	3 (0%)	52	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	305	ASP
1	D	122	TRP
1	C	122	TRP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/399 (98%)	382 (98%)	7 (2%)	66	52
1	B	376/399 (94%)	372 (99%)	4 (1%)	80	72
1	C	397/399 (100%)	388 (98%)	9 (2%)	58	41
1	D	379/399 (95%)	374 (99%)	5 (1%)	76	65
All	All	1541/1596 (97%)	1516 (98%)	25 (2%)	72	57

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	74	LYS
1	A	337	LYS
1	A	361	SER
1	A	365[A]	ARG
1	A	365[B]	ARG
1	A	403	ASN
1	B	185	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	269	ARG
1	B	317	LEU
1	B	403	ASN
1	C	3	VAL
1	C	185	ARG
1	C	277	LYS
1	C	300[A]	VAL
1	C	300[B]	VAL
1	C	316	ASP
1	C	317[A]	LEU
1	C	317[B]	LEU
1	C	403	ASN
1	D	5	LYS
1	D	203	ARG
1	D	361	SER
1	D	370	ASN
1	D	403	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	ASN
1	A	403	ASN
1	A	437	ASN
1	B	20	GLN
1	B	246	ASN
1	B	275	ASN
1	B	403	ASN
1	B	437	ASN
1	C	129	GLN
1	C	403	ASN
1	D	47	ASN
1	D	160	ASN
1	D	403	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	LGS	A	1446	-	20,23,23	0.97	2 (10%)	21,31,31	1.69	3 (14%)
2	LGS	B	1446	-	20,23,23	0.87	1 (5%)	21,31,31	2.25	5 (23%)
2	LGS	C	1446	-	20,23,23	1.04	2 (10%)	21,31,31	3.11	10 (47%)
3	ACT	C	1447	-	1,3,3	1.20	0	0,3,3	0.00	-
2	LGS	D	1446	-	20,23,23	0.96	1 (5%)	21,31,31	2.18	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LGS	A	1446	-	-	0/7/42/42	0/2/2/2
2	LGS	B	1446	-	-	0/7/42/42	0/2/2/2
2	LGS	C	1446	-	-	0/7/42/42	0/2/2/2
3	ACT	C	1447	-	-	0/0/0/0	0/0/0/0
2	LGS	D	1446	-	-	0/7/42/42	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1446	LGS	C2-C1	2.28	1.55	1.52
2	C	1446	LGS	C2-C1	2.32	1.55	1.52
2	A	1446	LGS	O1-C1	2.65	1.44	1.41
2	D	1446	LGS	O1-C1	2.65	1.44	1.41
2	B	1446	LGS	C2-C1	2.94	1.56	1.52
2	C	1446	LGS	O1-C1	3.19	1.45	1.41

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1446	LGS	C3-C4-C5	-7.53	101.24	111.39
2	D	1446	LGS	C3-C4-C5	-6.53	102.59	111.39
2	C	1446	LGS	C9-C8-N2	-6.34	100.38	110.73
2	B	1446	LGS	C3-C4-C5	-5.98	103.33	111.39
2	A	1446	LGS	C3-C4-C5	-3.99	106.01	111.39
2	D	1446	LGS	O3-C3-C2	-3.83	101.72	110.34
2	C	1446	LGS	O3-C3-C2	-3.24	103.04	110.34
2	C	1446	LGS	O2-C2-C1	-3.20	102.81	109.07
2	C	1446	LGS	C10-C9-C8	-3.15	101.10	113.97
2	C	1446	LGS	C11-C10-C9	-2.45	101.87	114.53
2	C	1446	LGS	C12-C11-C10	-2.45	101.87	114.53
2	B	1446	LGS	C9-C8-N2	-2.39	106.82	110.73
2	B	1446	LGS	O3-C3-C2	-2.35	105.05	110.34
2	C	1446	LGS	C13-C12-C11	-2.30	102.65	114.53
2	D	1446	LGS	C9-C8-N2	-2.01	107.45	110.73
2	A	1446	LGS	C3-C2-C1	2.37	112.90	109.13
2	D	1446	LGS	C6-C5-N1	3.09	102.57	100.00
2	B	1446	LGS	C3-C2-C1	3.26	114.30	109.13
2	C	1446	LGS	C3-C2-C1	3.76	115.11	109.13
2	D	1446	LGS	C3-C2-C1	3.96	115.42	109.13
2	A	1446	LGS	C6-C5-N1	5.35	104.44	100.00
2	C	1446	LGS	C6-C5-N1	5.88	104.89	100.00
2	B	1446	LGS	C6-C5-N1	6.24	105.19	100.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1447	ACT	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	443/468 (94%)	-0.31	5 (1%) 82 82	8, 16, 28, 39	0
1	B	443/468 (94%)	-0.18	6 (1%) 78 78	10, 19, 32, 41	0
1	C	443/468 (94%)	-0.49	8 (1%) 71 71	6, 11, 24, 38	0
1	D	443/468 (94%)	-0.41	4 (0%) 85 85	8, 14, 28, 44	0
All	All	1772/1872 (94%)	-0.35	23 (1%) 79 79	6, 15, 29, 44	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	VAL	5.3
1	D	3	VAL	4.2
1	B	3	VAL	4.1
1	B	307	PRO	3.6
1	D	304	PRO	3.6
1	D	305	ASP	3.3
1	C	3	VAL	3.3
1	A	316	ASP	3.2
1	A	305	ASP	3.0
1	C	304	PRO	2.9
1	C	274[A]	GLU	2.8
1	B	306	ALA	2.6
1	D	214	ASP	2.6
1	A	8	GLU	2.5
1	A	362	GLU	2.4
1	B	305	ASP	2.3
1	B	362	GLU	2.3
1	C	314	GLU	2.2
1	C	231	GLU	2.2
1	C	305	ASP	2.1
1	C	275[A]	ASN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	262	GLU	2.1
1	C	445	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	LGS	D	1446	22/22	0.97	0.07	-0.01	6,10,24,29	0
2	LGS	B	1446	22/22	0.97	0.08	-0.04	11,14,27,33	0
2	LGS	C	1446	22/22	0.97	0.07	-0.18	5,8,20,24	0
2	LGS	A	1446	22/22	0.98	0.06	-0.55	8,11,25,26	0
3	ACT	C	1447	4/4	0.89	0.09	-	23,24,25,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.