



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:09 PM GMT

PDB ID : 3WCO  
Title : Crystal structure of the depentamerized mutant of N-terminal truncated selenocysteine synthase Sela  
Authors : Itoh, Y.; Sekine, S.; Yokoyama, S.  
Deposited on : 2013-05-29  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	<div> <div>7%</div> <div>82%</div> <div>16%</div> <div>..</div> </div>
1	B	392	<div> <div>4%</div> <div>78%</div> <div>19%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	THJ	A	3002	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-seryl-tRNA(Sec) selenium transferase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	385	Total	C	N	O	P	S	Se	0	0	0
			3036	1931	530	564	1	3	7			
1	B	382	Total	C	N	O	P	S	Se	0	0	0
			3012	1915	526	561	1	3	6			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	191	TYR	THR	ENGINEERED MUTATION	UNP O67140
A	192	TYR	THR	ENGINEERED MUTATION	UNP O67140
A	199	ARG	ASP	ENGINEERED MUTATION	UNP O67140
A	220	PRO	TYR	ENGINEERED MUTATION	UNP O67140
B	191	TYR	THR	ENGINEERED MUTATION	UNP O67140
B	192	TYR	THR	ENGINEERED MUTATION	UNP O67140
B	199	ARG	ASP	ENGINEERED MUTATION	UNP O67140
B	220	PRO	TYR	ENGINEERED MUTATION	UNP O67140

- Molecule 2 is THIOSULFATE (three-letter code: THJ) (formula: O<sub>3</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	3	2		
2	A	1	Total	O	S	0	0
			5	3	2		
2	A	1	Total	O	S	0	0
			5	3	2		
2	B	1	Total	O	S	0	0
			5	3	2		
2	B	1	Total	O	S	0	0
			5	3	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	52	Total	O	0	0
			52	52		
3	B	53	Total	O	0	0
			53	53		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-seryl-tRNA(Sec) selenium transferase



- Molecule 1: L-seryl-tRNA(Sec) selenium transferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.19Å 81.29Å 95.54Å 90.00° 92.23° 90.00°	Depositor
Resolution (Å)	41.16 – 2.40 41.16 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.1 (41.16-2.40) 99.2 (41.16-2.40)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.59 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.179 , 0.229 0.188 , 0.240	Depositor DCC
$R_{free}$ test set	1767 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.3	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 53.8	EDS
Estimated twinning fraction	0.074 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 35379 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6178	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, THJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/3043	0.55	0/4075
1	B	0.40	0/3019	0.57	0/4044
All	All	0.40	0/6062	0.56	0/8119

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3036	0	3200	44	0
1	B	3012	0	3169	47	0
2	A	15	0	0	3	0
2	B	10	0	0	1	0
3	A	52	0	0	3	0
3	B	53	0	0	0	0
All	All	6178	0	6369	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:250:SER:HB2	1:B:286:LEU:HD12	1.71	0.73
1:A:118:SER:HB2	2:A:3002:THJ:O2	1.98	0.64
1:B:63:SER:OG	1:B:64:LEU:N	2.29	0.64
1:B:129:GLU:O	1:B:269:ARG:NH2	2.35	0.60
1:A:309:PRO:HG2	1:B:181:LYS:HG3	1.84	0.60
1:A:384:SER:HB3	3:A:3115:HOH:O	2.02	0.60
1:A:199:ARG:NH1	3:A:3120:HOH:O	2.30	0.59
1:B:120:ILE:HG13	1:B:137:PHE:CE1	2.37	0.59
1:B:179:MSE:HE2	1:B:186:LEU:HD21	1.84	0.58
1:B:158:GLU:OE2	1:B:205:ASN:ND2	2.37	0.58
1:B:255:ASN:H	1:B:255:ASN:HD22	1.53	0.56
1:B:255:ASN:ND2	1:B:255:ASN:H	2.04	0.55
1:A:166:LEU:HD12	1:A:169:ILE:HD11	1.88	0.55
1:A:181:LYS:HD2	1:B:309:PRO:HG2	1.89	0.53
1:B:161:ILE:HD13	1:B:166:LEU:HD13	1.91	0.52
2:A:3002:THJ:S2	1:B:174:ARG:NH1	2.74	0.52
1:A:71:ARG:HH22	1:B:102:ASN:HB3	1.75	0.51
1:B:158:GLU:OE1	1:B:187:ARG:NH1	2.44	0.51
1:B:159:VAL:HG22	1:B:210:LEU:HB3	1.91	0.51
1:B:376:ASP:CG	1:B:423:ARG:HH22	2.14	0.51
1:A:316:ILE:HD12	1:A:321:LEU:HB2	1.93	0.50
1:A:178:ILE:HD11	1:B:312:ARG:HB3	1.93	0.49
1:A:444:LYS:HA	1:A:447:GLN:NE2	2.27	0.49
1:B:150:LEU:HD21	1:B:179:MSE:HE3	1.95	0.49
1:A:189:VAL:HG11	1:A:200:TYR:HA	1.93	0.49
1:B:135:SER:HB3	1:B:302:ILE:HG13	1.95	0.48
1:B:91:LYS:HE3	1:B:95:ASN:HD21	1.78	0.48
1:A:437:GLU:H	1:A:437:GLU:CD	2.15	0.48
1:B:125:LYS:O	1:B:129:GLU:HG3	2.13	0.48
1:A:423:ARG:HG2	1:A:424:GLU:HG3	1.95	0.48
1:A:148:LEU:HG	1:A:305:ILE:HG23	1.94	0.48
1:A:72:VAL:HG22	1:A:417:PRO:HB2	1.94	0.48
1:A:313:ALA:HB1	1:B:144:GLY:HA2	1.95	0.48
1:A:174:ARG:HH21	1:B:312:ARG:HD3	1.79	0.47
1:A:376:ASP:CG	1:A:423:ARG:HH22	2.18	0.47
1:A:397:ILE:HG13	1:A:427:LEU:HB2	1.97	0.47
1:A:253:LEU:HD23	1:A:268:PHE:HD2	1.80	0.47
1:A:366:ILE:HA	1:A:367:PRO:HD2	1.60	0.47
1:A:315:ARG:HD3	2:A:3002:THJ:S2	2.56	0.46
1:B:213:LYS:HD3	1:B:230:LEU:HD23	1.98	0.46
1:A:161:ILE:HD13	1:A:166:LEU:HD13	1.98	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:SER:HB2	2:B:502:THJ:O1	2.16	0.45
1:A:369:LEU:HA	1:A:369:LEU:HD23	1.84	0.45
1:A:420:CYS:SG	1:A:427:LEU:HD22	2.56	0.45
1:A:312:ARG:HB3	1:B:178:ILE:HD11	1.98	0.45
1:B:123:ILE:HA	1:B:126:TYR:HD2	1.81	0.45
1:B:212:MSE:HE3	1:B:245:TYR:CE1	2.52	0.45
1:A:244:THR:N	1:A:277:ASP:OD2	2.34	0.45
1:B:254:ILE:HD12	1:B:259:PHE:CE1	2.52	0.44
1:A:399:HIS:CD2	1:A:450:LEU:HD22	2.52	0.44
1:A:212:MSE:HE2	1:A:214:VAL:HG22	1.98	0.44
1:A:144:GLY:HA2	1:B:313:ALA:HB1	1.99	0.44
1:B:422:ILE:HG12	1:B:427:LEU:HD23	1.99	0.44
1:A:142:ASN:HB3	1:A:285:LLP:OP4	2.17	0.43
1:B:169:ILE:HG23	1:B:214:VAL:HG11	2.00	0.43
1:A:148:LEU:HD22	1:A:314:LEU:HD12	2.00	0.43
1:A:402:LEU:HD12	1:A:402:LEU:HA	1.83	0.43
1:A:436:HIS:N	1:A:437:GLU:OE1	2.52	0.43
1:B:107:LEU:HD22	1:B:318:LYS:HB2	2.01	0.42
1:A:406:GLU:O	1:A:410:ARG:HG3	2.19	0.42
1:B:191:TYR:CG	1:B:192:TYR:N	2.87	0.42
1:A:254:ILE:HD12	1:A:259:PHE:HE1	1.84	0.42
1:A:405:GLN:HG2	3:A:3125:HOH:O	2.19	0.42
1:A:87:ALA:HA	1:A:88:PRO:HD2	1.95	0.41
1:B:359:LEU:HB2	1:B:439:LEU:HD22	2.02	0.41
1:B:261:ILE:HG23	1:B:390:LEU:HD21	2.01	0.41
1:A:399:HIS:CE1	1:A:401:ARG:HB2	2.55	0.41
1:B:406:GLU:O	1:B:410:ARG:HG3	2.20	0.41
1:B:147:PHE:CE1	1:B:182:SER:HA	2.54	0.41
1:A:65:MSE:HB3	1:A:65:MSE:HE2	1.97	0.41
1:A:174:ARG:HE	1:B:312:ARG:HB2	1.86	0.41
1:A:175:ILE:HA	1:A:178:ILE:HD12	2.02	0.41
1:B:148:LEU:HG	1:B:305:ILE:HG23	2.01	0.41
1:A:399:HIS:HE1	1:A:401:ARG:HD2	1.86	0.41
1:B:236:LEU:HD23	1:B:236:LEU:HA	1.83	0.41
1:B:370:LYS:HD2	1:B:370:LYS:HA	1.80	0.41
1:B:206:GLN:H	1:B:206:GLN:HG2	1.56	0.41
1:B:162:SER:OG	1:B:195:THR:HG21	2.20	0.41
1:B:142:ASN:HB3	1:B:285:LLP:OP4	2.21	0.40
1:B:388:LEU:HA	1:B:388:LEU:HD12	1.90	0.40
1:B:405:GLN:O	1:B:408:SER:HB2	2.21	0.40
1:A:244:THR:O	1:A:277:ASP:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/392 (97%)	364 (96%)	16 (4%)	0	100	100
1	B	377/392 (96%)	368 (98%)	9 (2%)	0	100	100
All	All	757/784 (97%)	732 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/331 (101%)	322 (97%)	11 (3%)	45	66
1	B	330/331 (100%)	322 (98%)	8 (2%)	57	76
All	All	663/662 (100%)	644 (97%)	19 (3%)	50	71

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	SER
1	A	263	VAL
1	A	264	ASP
1	A	332	PHE
1	A	360	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	384	SER
1	A	400	ASP
1	A	408	SER
1	A	424	GLU
1	A	426	GLN
1	A	437	GLU
1	B	118	SER
1	B	255	ASN
1	B	263	VAL
1	B	316	ILE
1	B	358	ARG
1	B	387	GLU
1	B	389	GLU
1	B	451	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	LLP	A	285	1	23,24,25	1.67	4 (17%)	28,32,34	1.38	6 (21%)
1	LLP	B	285	1	23,24,25	1.53	4 (17%)	28,32,34	1.45	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	285	1	-	0/15/17/19	0/1/1/1
1	LLP	B	285	1	-	0/15/17/19	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	285	LLP	CE-NZ	-2.67	1.41	1.46
1	A	285	LLP	CE-NZ	-2.59	1.41	1.46
1	B	285	LLP	C4'-NZ	2.74	1.35	1.27
1	B	285	LLP	C2'-C2	2.91	1.56	1.50
1	A	285	LLP	C2'-C2	3.02	1.56	1.50
1	A	285	LLP	C4'-NZ	3.12	1.36	1.27
1	B	285	LLP	C4-C4'	3.71	1.53	1.46
1	A	285	LLP	C4-C4'	4.31	1.54	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	285	LLP	CD-CE-NZ	-2.88	106.26	110.98
1	A	285	LLP	C5-C6-N1	-2.65	119.25	123.86
1	B	285	LLP	C5-C6-N1	-2.63	119.30	123.86
1	A	285	LLP	CD-CE-NZ	-2.52	106.86	110.98
1	B	285	LLP	O-C-CA	-2.41	119.22	125.49
1	A	285	LLP	C5-C4-C4'	-2.16	118.41	121.52
1	A	285	LLP	O-C-CA	-2.04	120.17	125.49
1	A	285	LLP	C3-C4-C4'	2.76	123.73	120.16
1	A	285	LLP	OP4-C5'-C5	3.13	114.17	108.99
1	B	285	LLP	OP4-C5'-C5	4.32	116.14	108.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	285	LLP	1	0
1	B	285	LLP	1	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	THJ	A	3001	-	3,4,4	1.01	0	4,6,6	0.78	0
2	THJ	A	3002	-	3,4,4	0.81	0	4,6,6	1.47	1 (25%)
2	THJ	A	3003	-	3,4,4	1.25	0	4,6,6	0.56	0
2	THJ	B	501	-	3,4,4	1.20	0	4,6,6	0.73	0
2	THJ	B	502	-	3,4,4	0.75	0	4,6,6	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	THJ	A	3001	-	-	0/0/0/0	0/0/0/0
2	THJ	A	3002	-	-	0/0/0/0	0/0/0/0
2	THJ	A	3003	-	-	0/0/0/0	0/0/0/0
2	THJ	B	501	-	-	0/0/0/0	0/0/0/0
2	THJ	B	502	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3002	THJ	O2-S1-S2	-2.80	101.58	108.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3002	THJ	3	0
2	B	502	THJ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	377/392 (96%)	0.22	26 (6%) 20 19	35, 62, 101, 136	0
1	B	375/392 (95%)	0.11	15 (4%) 42 43	34, 61, 91, 121	0
All	All	752/784 (95%)	0.17	41 (5%) 29 29	34, 62, 96, 136	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	SER	7.7
1	A	452	ILE	4.4
1	B	320	THR	3.5
1	A	425	ASP	3.5
1	A	189	VAL	3.4
1	B	319	LEU	3.4
1	A	398	ARG	3.3
1	A	424	GLU	3.0
1	B	314	LEU	3.0
1	A	401	ARG	3.0
1	A	140	ASN	2.9
1	A	319	LEU	2.8
1	B	139	VAL	2.8
1	B	140	ASN	2.7
1	A	316	ILE	2.7
1	B	316	ILE	2.7
1	A	320	THR	2.7
1	B	424	GLU	2.6
1	A	202	GLY	2.5
1	A	368	GLY	2.4
1	A	402	LEU	2.4
1	A	370	LYS	2.3
1	B	452	ILE	2.3
1	A	191	TYR	2.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	374	ILE	2.2
1	B	226	GLU	2.2
1	A	194	LYS	2.2
1	A	314	LEU	2.2
1	A	228	VAL	2.2
1	B	317	ASP	2.2
1	A	138	VAL	2.1
1	A	292	ALA	2.1
1	A	294	ILE	2.1
1	A	291	GLN	2.1
1	A	366	ILE	2.1
1	A	290	PRO	2.1
1	B	196	LYS	2.0
1	B	310	ILE	2.0
1	B	146	VAL	2.0
1	B	374	ILE	2.0
1	A	227	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	B	285	24/25	0.97	0.16	-	32,45,55,61	0
1	LLP	A	285	24/25	0.98	0.17	-	34,43,56,63	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	THJ	A	3002	5/5	0.94	0.18	0.31	64,66,74,95	0
2	THJ	B	501	5/5	0.96	0.16	-0.45	73,79,88,98	0
2	THJ	B	502	5/5	0.96	0.14	-0.75	62,69,73,77	0
2	THJ	A	3003	5/5	0.97	0.14	-0.88	84,84,88,93	0
2	THJ	A	3001	5/5	0.97	0.12	-1.26	56,72,78,99	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.