



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:11 PM GMT

PDB ID : 3WE2  
Title : Structure of BLM RQC domain bound to a phosphate ion  
Authors : Kim, S.Y.; Hakoshima, T.; Kitano, K.  
Deposited on : 2013-06-28  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	<div> <div>8%</div> <div>48%</div> <div>32%</div> <div>••</div> <div>18%</div> </div>
1	B	147	<div> <div>7%</div> <div>54%</div> <div>20%</div> <div>•</div> <div>24%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1839 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bloom syndrome protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	121	Total	C	N	O	S	0	0	0
			950	595	169	181	5			
1	B	112	Total	C	N	O	S	0	0	0
			878	553	152	169	4			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1063	GLY	-	EXPRESSION TAG	UNP P54132
A	1064	PRO	-	EXPRESSION TAG	UNP P54132
A	1065	LEU	-	EXPRESSION TAG	UNP P54132
A	1066	GLY	-	EXPRESSION TAG	UNP P54132
A	1067	SER	-	EXPRESSION TAG	UNP P54132
B	1063	GLY	-	EXPRESSION TAG	UNP P54132
B	1064	PRO	-	EXPRESSION TAG	UNP P54132
B	1065	LEU	-	EXPRESSION TAG	UNP P54132
B	1066	GLY	-	EXPRESSION TAG	UNP P54132
B	1067	SER	-	EXPRESSION TAG	UNP P54132

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bloom syndrome protein



- Molecule 1: Bloom syndrome protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.22Å 59.22Å 210.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 19.86 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.70) 93.8 (19.86-2.70)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.05 (at 2.71Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.237 , 0.271 0.238 , 0.272	Depositor DCC
$R_{free}$ test set	541 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	73.9	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 56.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 10768 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	1839	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/962	0.66	0/1289
1	B	0.44	0/888	0.62	0/1188
All	All	0.42	0/1850	0.64	0/2477

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	958	38	1
1	B	878	0	885	27	0
2	A	5	0	0	1	0
3	A	4	0	3	0	0
4	A	2	0	0	0	0
All	All	1839	0	1846	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1162:ASN:HD22	1:B:1168:ILE:HD13	1.35	0.90
1:A:1089:GLN:HB3	1:A:1173:LEU:HD11	1.52	0.89
1:A:1103:VAL:HG12	1:A:1104:GLY:H	1.37	0.87
1:B:1109:PHE:HD2	1:B:1113:MET:HE1	1.46	0.81
1:B:1162:ASN:OD1	1:B:1164:ASN:HB3	1.84	0.77
1:B:1109:PHE:CD2	1:B:1113:MET:HE1	2.21	0.75
1:A:1103:VAL:HG12	1:A:1104:GLY:N	2.02	0.74
1:A:1150:ILE:HA	1:A:1155:LEU:O	1.90	0.72
1:B:1074:THR:HB	1:B:1190:MET:HG2	1.74	0.69
1:B:1089:GLN:CB	1:B:1173:LEU:HD11	2.25	0.66
1:A:1145:LEU:O	1:A:1149:LEU:HG	1.96	0.66
1:B:1162:ASN:ND2	1:B:1168:ILE:HD13	2.09	0.66
1:B:1089:GLN:HB3	1:B:1173:LEU:HD11	1.79	0.65
1:A:1119:LEU:HD11	1:A:1139:ARG:HA	1.82	0.61
1:B:1139:ARG:O	1:B:1143:GLU:HG3	2.00	0.61
1:A:1148:LYS:HE2	1:A:1188:ASP:O	2.00	0.61
1:A:1113:MET:O	1:A:1117:ILE:HG13	2.00	0.60
1:A:1103:VAL:CG1	1:A:1104:GLY:H	2.14	0.60
1:A:1193:GLU:O	1:A:1194:ASN:HB3	2.02	0.59
1:A:1121:SER:HB2	2:A:1301:PO4:O4	2.02	0.59
1:A:1108:ARG:HG2	1:A:1170:TYR:CD1	2.40	0.57
1:B:1148:LYS:HE3	1:B:1188:ASP:O	2.06	0.55
1:A:1104:GLY:HA3	1:A:1108:ARG:HD3	1.89	0.55
1:A:1161:ILE:HA	1:A:1166:GLN:O	2.07	0.54
1:A:1108:ARG:HG3	1:A:1108:ARG:HH11	1.72	0.54
1:B:1084:ILE:O	1:B:1088:VAL:HG23	2.08	0.53
1:A:1077:VAL:O	1:A:1081:VAL:HG23	2.08	0.53
1:A:1175:ASN:O	1:A:1176:LYS:HB2	2.08	0.53
1:B:1085:VAL:CG2	1:B:1181:LEU:HD13	2.38	0.52
1:A:1103:VAL:CG1	1:A:1104:GLY:N	2.73	0.51
1:A:1182:ASN:HB2	1:A:1184:ASN:ND2	2.25	0.51
1:A:1084:ILE:HD13	1:A:1146:PHE:CZ	2.47	0.50
1:B:1105:PRO:HB2	1:B:1170:TYR:HB2	1.93	0.50
1:B:1117:ILE:O	1:B:1131:PHE:HA	2.12	0.49
1:B:1153:LYS:HE2	1:B:1175:ASN:ND2	2.27	0.49
1:A:1108:ARG:HG2	1:A:1170:TYR:HD1	1.77	0.48
1:A:1182:ASN:C	1:A:1184:ASN:H	2.16	0.48
1:B:1153:LYS:HE3	1:B:1156:ASP:OD2	2.14	0.48
1:A:1089:GLN:HB3	1:A:1173:LEU:CD1	2.35	0.47
1:B:1075:ARG:CZ	1:B:1075:ARG:HB2	2.44	0.47
1:B:1133:LYS:HZ3	1:B:1133:LYS:HB2	1.79	0.47
1:A:1192:THR:HG22	1:A:1193:GLU:N	2.28	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1080:ASP:O	1:A:1084:ILE:HG13	2.14	0.47
1:A:1099:ASN:O	1:A:1101:LYS:N	2.48	0.47
1:B:1081:VAL:O	1:B:1085:VAL:HG13	2.16	0.46
1:A:1185:LEU:HD12	1:A:1186:LYS:N	2.31	0.46
1:B:1111:MET:O	1:B:1115:VAL:HG23	2.16	0.45
1:B:1164:ASN:ND2	1:B:1166:GLN:HB2	2.32	0.45
1:A:1144:ARG:NH1	1:A:1189:PHE:CZ	2.85	0.45
1:A:1127:GLN:HG2	1:A:1131:PHE:CE1	2.53	0.44
1:A:1110:THR:OG1	1:A:1113:MET:HG3	2.17	0.44
1:A:1130:ILE:O	1:A:1130:ILE:HG13	2.18	0.43
1:A:1086:ARG:HD3	1:A:1181:LEU:HD22	2.01	0.43
1:A:1103:VAL:HG12	1:A:1107:GLY:HA3	2.01	0.43
1:A:1086:ARG:HD3	1:A:1181:LEU:CD2	2.49	0.42
1:B:1168:ILE:HD12	1:B:1170:TYR:CE2	2.55	0.42
1:B:1153:LYS:O	1:B:1153:LYS:HD2	2.20	0.42
1:A:1127:GLN:HA	1:A:1131:PHE:CD1	2.55	0.42
1:B:1180:VAL:CG2	1:B:1185:LEU:HD23	2.50	0.42
1:A:1108:ARG:HB3	1:A:1170:TYR:HB2	2.02	0.41
1:B:1133:LYS:HB2	1:B:1133:LYS:NZ	2.35	0.41
1:B:1105:PRO:O	1:B:1170:TYR:CD2	2.74	0.41
1:B:1120:GLY:HA2	1:B:1131:PHE:CE1	2.55	0.41
1:A:1192:THR:CG2	1:A:1193:GLU:N	2.84	0.40
1:A:1103:VAL:CG1	1:A:1107:GLY:HA3	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1122:LYS:CG	1:A:1122:LYS:CG[8_665]	2.09	0.11

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/147 (81%)	95 (80%)	19 (16%)	5 (4%)	3	7
1	B	108/147 (74%)	100 (93%)	8 (7%)	0	100	100
All	All	227/294 (77%)	195 (86%)	27 (12%)	5 (2%)	8	22

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1176	LYS
1	A	1120	GLY
1	A	1183	GLY
1	A	1119	LEU
1	A	1100	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/127 (83%)	99 (94%)	6 (6%)	25	53
1	B	96/127 (76%)	92 (96%)	4 (4%)	36	68
All	All	201/254 (79%)	191 (95%)	10 (5%)	30	60

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1079	ASP
1	A	1086	ARG
1	A	1108	ARG
1	A	1156	ASP
1	A	1172	MET
1	A	1176	LYS
1	B	1074	THR
1	B	1089	GLN
1	B	1153	LYS
1	B	1156	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1089	GLN
1	A	1112	ASN
1	A	1175	ASN
1	A	1184	ASN
1	B	1089	GLN
1	B	1164	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	1301	-	4,4,4	1.08	0	6,6,6	0.27	0
3	ACT	A	1302	-	1,3,3	2.30	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1301	-	-	0/0/0/0	0/0/0/0
3	ACT	A	1302	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1302	ACT	CH3-C	2.30	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1301	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	121/147 (82%)	0.42	12 (9%) 9 7	59, 87, 128, 152	0
1	B	112/147 (76%)	0.39	11 (9%) 10 7	58, 81, 120, 136	0
All	All	233/294 (79%)	0.41	23 (9%) 9 7	58, 84, 126, 152	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1105	PRO	9.6
1	B	1104	GLY	6.1
1	A	1096	GLY	4.2
1	B	1194	ASN	4.1
1	A	1099	ASN	3.8
1	A	1102	HIS	3.6
1	B	1163	ALA	3.5
1	B	1093	SER	3.5
1	A	1098	ARG	3.3
1	A	1100	ILE	3.1
1	A	1097	MET	3.1
1	A	1194	ASN	3.0
1	B	1164	ASN	2.9
1	B	1107	GLY	2.7
1	B	1106	SER	2.6
1	A	1084	ILE	2.5
1	B	1160	TYR	2.5
1	A	1081	VAL	2.5
1	B	1122	LYS	2.3
1	A	1193	GLU	2.2
1	A	1101	LYS	2.2
1	A	1190	MET	2.0
1	B	1075	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	A	1301	5/5	0.83	0.26	0.54	115,116,117,117	0
3	ACT	A	1302	4/4	0.91	0.20	0.41	88,88,88,89	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.