



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:12 PM GMT

PDB ID : 3WGW
Title : Structure of PCNA bound to a small molecule inhibitor
Authors : Hashimoto, H.
Deposited on : 2013-08-12
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 99%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 74% 16% • • 5% </div> </div>
1	B	261	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, green 97%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 70% 20% 5% 5% </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proliferating cell nuclear antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1907	1201	313	377	16			
1	B	249	Total	C	N	O	S	0	0	0
			1913	1204	314	379	16			

- Molecule 2 is 4-{4-[(2S)-2-AMINO-3-HYDROXYPROPYL]-2,6-DIIODOPHENOXY}PHE NOL (three-letter code: T2B) (formula: C₁₅H₁₅I₂NO₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	I	N	O	0	0
			21	15	2	1	3		
2	A	1	Total	C	I	N	O	0	0
			21	15	2	1	3		
2	B	1	Total	C	I	N	O	0	0
			21	15	2	1	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	66	Total	O	0	0
			66	66		
4	B	60	Total	O	0	0
			60	60		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proliferating cell nuclear antigen



- Molecule 1: Proliferating cell nuclear antigen



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	192.17Å 192.17Å 192.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 19.82 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-2.80) 99.8 (19.82-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.13 (at 2.79Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.182 , 0.221 0.182 , 0.221	Depositor DCC
R_{free} test set	1532 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	49.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 30251 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4019	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: T2B, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	0/1932	1.18	8/2609 (0.3%)
1	B	0.96	0/1938	1.11	4/2617 (0.2%)
All	All	1.00	0/3870	1.15	12/5226 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	16	LEU	CA-CB-CG	-9.38	93.71	115.30
1	A	162	CYS	CA-CB-SG	-7.27	100.91	114.00
1	A	12	LEU	CB-CG-CD2	-7.14	98.86	111.00
1	A	175	LEU	CA-CB-CG	6.75	130.82	115.30
1	B	235	LEU	CB-CG-CD1	-6.71	99.59	111.00
1	B	162	CYS	CA-CB-SG	-5.94	103.31	114.00
1	A	6	LEU	CB-CG-CD2	-5.90	100.96	111.00
1	B	150	ASP	CB-CG-OD2	-5.76	113.12	118.30
1	A	156	ASP	CB-CA-C	-5.50	99.39	110.40
1	A	170	SER	CB-CA-C	-5.28	100.06	110.10
1	A	54	SER	CB-CA-C	-5.21	100.21	110.10
1	B	175	LEU	CB-CG-CD1	5.00	119.50	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	254	LYS	Peptide
1	B	193	GLU	Peptide
1	B	241	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1907	0	1923	30	0
1	B	1913	0	1928	31	0
2	A	42	0	28	2	0
2	B	21	0	15	0	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
4	A	66	0	0	2	0
4	B	60	0	0	2	0
All	All	4019	0	3894	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:SER:HB3	1:A:177:ASN:HB3	1.35	1.07
1:A:140:PRO:HG3	1:A:193:GLU:HB3	1.44	1.00
1:A:53:ARG:HD3	4:A:424:HOH:O	1.73	0.87
1:A:56:GLY:HA3	1:A:244:MET:HG3	1.66	0.78
1:B:172:SER:HB3	1:B:177:ASN:HB3	1.67	0.74
1:B:13:LYS:NZ	1:B:82:ALA:O	2.21	0.72
1:B:185:THR:HG21	1:B:194:ALA:HB1	1.74	0.69
1:A:53:ARG:NH1	3:A:302:SO4:O1	2.26	0.68
1:B:107:ASN:O	1:B:109:GLU:N	2.27	0.67
1:B:30:ILE:HB	1:B:66:LEU:HD23	1.78	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:LEU:HD23	1:A:175:LEU:C	2.17	0.64
1:A:183:SER:HB2	2:A:303:T2B:C7	2.28	0.64
1:A:107:ASN:O	1:A:109:GLU:N	2.31	0.63
1:A:210:ARG:CG	1:A:210:ARG:HH11	2.14	0.60
1:A:68:MET:HB3	1:A:118:LEU:HD13	1.85	0.58
1:B:185:THR:HG21	1:B:194:ALA:CB	2.34	0.58
1:B:226:THR:HG22	4:B:459:HOH:O	2.02	0.57
1:A:210:ARG:NH1	4:A:407:HOH:O	2.38	0.56
1:A:175:LEU:CD2	1:A:175:LEU:C	2.74	0.55
1:A:16:LEU:HD13	1:A:79:LEU:HD12	1.89	0.55
1:B:25:GLU:HG2	1:B:119:MET:SD	2.47	0.55
1:B:107:ASN:O	1:B:107:ASN:OD1	2.24	0.54
1:B:226:THR:CG2	4:B:459:HOH:O	2.56	0.53
1:B:35:VAL:HB	1:B:52:LEU:HB2	1.91	0.52
1:A:85:GLU:OE2	1:A:85:GLU:HA	2.10	0.52
1:B:124:GLU:O	1:B:124:GLU:HG3	2.09	0.51
1:A:16:LEU:HD13	1:A:79:LEU:CD1	2.40	0.51
1:B:2:PHE:CG	1:B:30:ILE:HG21	2.46	0.50
1:A:9:GLY:HA3	1:A:88:ILE:HG13	1.93	0.49
1:B:41:ASP:O	1:B:43:SER:N	2.43	0.49
1:A:139:MET:CE	1:A:227:LEU:HD11	2.43	0.48
1:B:172:SER:HB3	1:B:177:ASN:CB	2.40	0.48
1:B:37:LEU:HB3	1:B:50:LEU:HB3	1.97	0.46
1:B:125:GLN:O	1:B:126:LEU:HD23	2.15	0.46
1:B:30:ILE:HB	1:B:66:LEU:CD2	2.45	0.46
1:B:105:ALA:HA	1:B:106:PRO:HD2	1.72	0.46
1:B:99:LEU:HD23	1:B:116:MET:CE	2.46	0.45
1:B:131:GLN:OE1	1:B:233:VAL:HG21	2.15	0.45
1:A:105:ALA:HA	1:A:106:PRO:HD3	1.79	0.45
1:B:32:SER:HA	1:B:62:CYS:HB3	1.99	0.45
1:A:2:PHE:CD2	1:A:30:ILE:HG21	2.52	0.44
1:A:140:PRO:HG3	1:A:193:GLU:CB	2.32	0.44
1:A:210:ARG:HG3	1:A:210:ARG:HH11	1.82	0.44
1:B:19:LEU:CD2	1:B:48:VAL:HG11	2.48	0.44
1:B:184:GLN:HG3	1:B:197:ILE:HD12	2.01	0.43
1:A:19:LEU:CD2	1:A:48:VAL:HG11	2.48	0.43
1:B:2:PHE:CD2	1:B:30:ILE:HG21	2.54	0.43
1:A:182:LEU:HA	1:A:182:LEU:HD23	1.78	0.43
1:B:107:ASN:O	1:B:107:ASN:CG	2.56	0.43
1:A:210:ARG:CG	1:A:210:ARG:NH1	2.78	0.43
1:A:199:MET:HG3	1:A:199:MET:O	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:GLY:O	2:A:301:T2B:I3	3.07	0.43
1:B:197:ILE:HG22	1:B:199:MET:HG2	2.00	0.42
1:A:37:LEU:C	1:A:37:LEU:HD23	2.39	0.42
1:B:82:ALA:HB2	1:B:103:PHE:CD2	2.55	0.42
1:B:234:PRO:HA	1:B:253:PRO:HD3	2.02	0.42
1:A:12:LEU:HD23	1:A:12:LEU:HA	1.66	0.42
1:B:231:ALA:O	1:B:233:VAL:HG23	2.20	0.41
1:A:222:SER:HB2	1:A:240:LYS:O	2.20	0.41
1:A:14:LYS:HD3	1:A:220:PRO:HB2	2.01	0.41
1:B:41:ASP:C	1:B:43:SER:H	2.24	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/261 (94%)	236 (97%)	3 (1%)	5 (2%)	9	30
1	B	245/261 (94%)	233 (95%)	6 (2%)	6 (2%)	7	25
All	All	489/522 (94%)	469 (96%)	9 (2%)	11 (2%)	8	28

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	108	GLN
1	B	107	ASN
1	B	108	GLN
1	A	54	SER
1	A	107	ASN
1	B	42	SER
1	B	43	SER
1	A	124	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	106	PRO
1	B	244	MET
1	A	244	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/228 (95%)	195 (90%)	21 (10%)	10	29
1	B	217/228 (95%)	191 (88%)	26 (12%)	6	19
All	All	433/456 (95%)	386 (89%)	47 (11%)	8	23

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LEU
1	A	22	LEU
1	A	27	CYS
1	A	43	SER
1	A	64	ARG
1	A	85	GLU
1	A	108	GLN
1	A	113	ASP
1	A	125	GLN
1	A	130	GLU
1	A	134	SER
1	A	146	ARG
1	A	175	LEU
1	A	177	ASN
1	A	179	ASN
1	A	183	SER
1	A	193	GLU
1	A	199	MET
1	A	200	ASN
1	A	210	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	230	SER
1	B	2	PHE
1	B	24	ASN
1	B	36	ASN
1	B	41	ASP
1	B	49	GLN
1	B	61	ARG
1	B	64	ARG
1	B	97	ASP
1	B	108	GLN
1	B	109	GLU
1	B	110	LYS
1	B	117	LYS
1	B	122	ASP
1	B	123	VAL
1	B	126	LEU
1	B	138	LYS
1	B	143	GLU
1	B	146	ARG
1	B	149	ARG
1	B	175	LEU
1	B	179	ASN
1	B	184	GLN
1	B	185	THR
1	B	193	GLU
1	B	226	THR
1	B	229	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	84	ASN
1	A	125	GLN
1	B	24	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	T2B	A	301	-	22,22,22	1.60	5 (22%)	27,30,30	1.74	6 (22%)
3	SO4	A	302	-	4,4,4	0.88	0	6,6,6	1.09	1 (16%)
2	T2B	A	303	-	22,22,22	1.39	3 (13%)	27,30,30	2.03	7 (25%)
3	SO4	B	301	-	4,4,4	0.61	0	6,6,6	0.82	0
2	T2B	B	302	-	22,22,22	1.29	3 (13%)	27,30,30	2.17	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	T2B	A	301	-	-	0/10/10/10	0/2/2/2
3	SO4	A	302	-	-	0/0/0/0	0/0/0/0
2	T2B	A	303	-	-	0/10/10/10	0/2/2/2
3	SO4	B	301	-	-	0/0/0/0	0/0/0/0
2	T2B	B	302	-	-	0/10/10/10	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	T2B	C5-I1	2.01	2.15	2.10
2	B	302	T2B	C3-C1	2.07	1.43	1.39
2	A	301	T2B	C11-C1	2.15	1.43	1.39
2	A	301	T2B	C7-C9	2.17	1.48	1.40
2	B	302	T2B	C7-C5	2.26	1.48	1.40
2	A	303	T2B	C7-C9	2.29	1.48	1.40
2	A	303	T2B	C7-C5	2.55	1.49	1.40
2	A	301	T2B	C7-C5	2.65	1.49	1.40
2	A	303	T2B	C12-C2	2.95	1.44	1.38
2	A	301	T2B	C17-C15	3.02	1.57	1.52
2	A	301	T2B	C5-I1	3.60	2.19	2.10

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	302	T2B	C1-C13-C15	-5.81	102.89	113.31
2	A	301	T2B	C3-C5-C7	-3.91	113.04	121.67
2	A	303	T2B	C7-C9-I3	-2.72	114.05	119.31
2	A	303	T2B	C12-C10-C8	-2.65	116.80	119.87
2	A	303	T2B	C7-C5-I1	-2.62	114.25	119.31
3	A	302	SO4	O2-S-O1	-2.49	101.62	109.50
2	B	302	T2B	C3-C5-C7	-2.31	116.57	121.67
2	A	301	T2B	C4-C6-C8	2.25	122.47	119.87
2	A	303	T2B	C10-C12-C2	2.26	122.59	119.74
2	A	301	T2B	C2-O2-C7	2.81	123.35	118.46
2	A	301	T2B	C7-C5-I1	2.81	124.75	119.31
2	A	301	T2B	O4-C17-C15	3.37	120.66	111.84
2	A	301	T2B	C1-C13-C15	3.45	119.48	113.31
2	A	303	T2B	C3-C5-I1	3.56	125.35	118.60
2	B	302	T2B	C11-C9-I3	3.81	125.82	118.60
2	B	302	T2B	C3-C5-I1	4.65	127.41	118.60
2	B	302	T2B	C2-O2-C7	4.66	126.57	118.46
2	A	303	T2B	C11-C9-I3	5.02	128.11	118.60
2	A	303	T2B	C2-O2-C7	5.34	127.77	118.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	T2B	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	SO4	1	0
2	A	303	T2B	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/261 (95%)	-0.51	3 (1%) 81 73	22, 35, 72, 111	0
1	B	249/261 (95%)	-0.43	8 (3%) 51 39	25, 40, 75, 123	0
All	All	497/522 (95%)	-0.47	11 (2%) 65 54	22, 37, 75, 123	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	255	ILE	3.6
1	B	232	ASP	3.2
1	B	122	ASP	3.0
1	B	186	SER	3.0
1	A	255	ILE	2.9
1	A	108	GLN	2.6
1	B	106	PRO	2.6
1	B	108	GLN	2.4
1	B	123	VAL	2.4
1	A	125	GLN	2.4
1	B	107	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	B	301	5/5	0.98	0.26	1.66	54,59,64,74	0
2	T2B	A	303	21/21	0.94	0.28	0.95	59,94,114,177	0
3	SO4	A	302	5/5	0.97	0.17	0.67	62,65,69,74	0
2	T2B	B	302	21/21	0.98	0.10	-0.65	34,52,71,103	0
2	T2B	A	301	21/21	0.99	0.10	-0.98	28,34,56,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.