



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:14 PM GMT

PDB ID : 1WMP
Title : Crystal structure of amine oxidase complexed with cobalt ion
Authors : Okajima, T.; Kishishita, S.; Chiu, Y.C.; Murakawa, T.; Kim, M.; Yamaguchi, H.; Hirota, S.; Kuroda, S.; Tanizawa, K.
Deposited on : 2004-07-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	638	<div> <div>5%</div> <div>78%</div> <div>18%</div> <div>••</div> </div>
1	B	638	<div> <div>4%</div> <div>85%</div> <div>12%</div> <div>•</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylethylamine oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	620	Total	C	N	O	S	0	0	0
			4866	3074	855	928	9			
1	B	620	Total	C	N	O	S	0	0	0
			4866	3074	855	928	9			

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Co	0	0
			1	1		
2	A	1	Total	Co	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	442	Total	O	0	0
			442	442		
3	B	533	Total	O	0	0
			533	533		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phenylethylamine oxidase



• Molecule 1: Phenylethylamine oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	157.58Å 63.38Å 184.20Å 90.00° 112.47° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 24.09 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 98.6 (24.09-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 1.99Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.188 , 0.220 0.180 , 0.211	Depositor DCC
R_{free} test set	5587 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.689	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 80.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 112241 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10709	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.17 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.2107e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/4989	0.61	0/6795
1	B	0.35	0/4989	0.62	0/6795
All	All	0.34	0/9978	0.62	0/13590

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4866	0	4688	96	0
1	B	4866	0	4688	51	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	442	0	0	5	0
3	B	533	0	0	4	0
All	All	10709	0	9376	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (140) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:PRO:HG2	1:A:90:LEU:HB2	1.59	0.83
1:A:143:GLU:HG3	1:A:144:TYR:H	1.49	0.77
1:A:104:GLU:O	1:A:107:VAL:HG22	1.85	0.77
1:A:108:VAL:HG22	1:A:179:VAL:HG21	1.67	0.76
1:A:422:LEU:HD11	1:A:428:ALA:HB2	1.70	0.74
1:A:297:PHE:HB2	1:A:301:GLU:HG3	1.68	0.74
1:A:451:GLU:HG3	3:A:1154:HOH:O	1.92	0.70
1:A:324:LEU:HB2	1:A:342:ILE:HB	1.74	0.68
1:A:105:PHE:HD1	1:A:136:PRO:HB2	1.57	0.68
1:A:247:VAL:HG11	1:A:344:MET:CE	2.24	0.68
1:B:324:LEU:HB2	1:B:342:ILE:HB	1.74	0.68
1:A:66:ASP:OD1	1:A:68:SER:HB3	1.94	0.67
1:A:162:PHE:HB2	1:A:163:PRO:HD2	1.77	0.66
1:A:247:VAL:HG11	1:A:344:MET:HE1	1.78	0.66
1:A:483:ALA:HB1	1:A:543:VAL:HB	1.77	0.66
1:A:170:HIS:HD2	1:A:198:PRO:O	1.80	0.65
1:A:247:VAL:HG21	1:A:324:LEU:HD11	1.80	0.64
1:A:206:ASP:OD1	1:A:208:GLU:HG2	1.95	0.64
1:A:505:ARG:HD3	1:A:618:ASP:HB3	1.79	0.64
1:B:72:PRO:HG2	1:B:90:LEU:HB2	1.81	0.63
1:A:105:PHE:CD1	1:A:136:PRO:HB2	2.37	0.60
1:B:599:TRP:CD2	1:B:600:PRO:HA	2.37	0.59
1:A:624:ASP:CG	1:B:214:ARG:HD2	2.22	0.59
1:B:451:GLU:HG3	3:B:1346:HOH:O	2.01	0.59
1:A:347:GLU:OE1	1:A:370:ARG:HD3	2.04	0.58
1:A:16:SER:OG	1:A:19:GLU:HG3	2.04	0.58
1:A:599:TRP:CD2	1:A:600:PRO:HA	2.39	0.57
1:A:78:SER:HB2	1:A:85:ILE:HD11	1.86	0.57
1:A:241:GLU:O	1:A:242:LYS:HB2	2.04	0.56
1:B:241:GLU:O	1:B:242:LYS:HB2	2.05	0.56
1:B:232:VAL:HG22	1:B:238:ILE:CD1	2.36	0.56
1:A:247:VAL:O	1:A:247:VAL:HG23	2.07	0.55
1:A:134:VAL:O	1:A:136:PRO:HD3	2.08	0.54
1:A:280:MET:HG3	1:A:435:PHE:CD2	2.42	0.54
1:B:111:LEU:HD11	1:B:184:LYS:HA	1.90	0.53
1:B:143:GLU:H	1:B:143:GLU:CD	2.12	0.53
1:A:297:PHE:HB2	1:A:301:GLU:CG	2.38	0.53
1:A:370:ARG:O	1:A:370:ARG:HG2	2.09	0.53
1:A:624:ASP:OD2	1:B:214:ARG:HD2	2.09	0.53
1:B:150:ARG:NH2	1:B:185:GLU:OE1	2.42	0.53
1:B:106:GLU:HB3	3:B:1503:HOH:O	2.08	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:318:LEU:HD12	3:B:1511:HOH:O	2.09	0.52
1:A:218:LYS:HD2	1:B:226:GLU:HB2	1.92	0.52
1:A:154:ARG:HD2	1:A:293:TRP:CE3	2.45	0.52
1:B:297:PHE:HB2	1:B:301:GLU:HG3	1.91	0.52
1:A:170:HIS:HE1	3:A:1234:HOH:O	1.93	0.51
1:B:152:ILE:HG22	1:B:153:LEU:N	2.26	0.51
1:B:20:ILE:HD12	1:B:327:VAL:HG12	1.93	0.50
1:A:232:VAL:HG22	1:A:238:ILE:CD1	2.42	0.50
1:A:180:ASP:OD1	1:A:182:VAL:HG22	2.12	0.49
1:A:143:GLU:HG3	1:A:144:TYR:N	2.25	0.49
1:A:95:THR:O	1:A:557:PHE:HB2	2.13	0.48
1:A:292:SER:OG	1:B:466:ARG:HD2	2.13	0.48
1:A:619:ARG:HA	1:B:596:VAL:HG12	1.95	0.48
1:A:460:MET:HE2	1:A:466:ARG:C	2.34	0.48
1:A:504:ASN:HD21	1:A:508:GLU:HB3	1.78	0.48
1:A:26:ILE:HD11	1:A:82:GLY:HA2	1.94	0.47
1:B:142:PHE:HB3	1:B:143:GLU:OE1	2.15	0.47
1:A:154:ARG:HD2	1:A:293:TRP:CD2	2.50	0.47
1:A:36:GLU:O	1:A:67:VAL:HG22	2.14	0.47
1:A:143:GLU:CG	1:A:144:TYR:H	2.23	0.47
1:A:182:VAL:HG22	3:A:1352:HOH:O	2.14	0.47
1:A:28:ARG:HG3	1:A:33:LEU:HD23	1.97	0.46
1:B:125:ARG:HG2	1:B:194:VAL:HG23	1.96	0.46
1:A:182:VAL:HG23	1:A:183:SER:N	2.31	0.46
1:A:152:ILE:HG22	1:A:153:LEU:N	2.31	0.46
1:A:206:ASP:CG	1:A:208:GLU:HG2	2.36	0.46
1:A:323:TYR:HE1	1:A:343:CYS:SG	2.39	0.46
1:A:443:ILE:H	1:A:448:ASN:HD21	1.64	0.46
1:A:20:ILE:HD12	1:A:327:VAL:HG12	1.97	0.46
1:A:321:ILE:HD12	1:A:321:ILE:N	2.29	0.46
1:A:15:ALA:HA	1:A:19:GLU:OE1	2.16	0.46
1:B:323:TYR:CE1	1:B:343:CYS:SG	3.09	0.46
1:A:246:ASP:HB2	1:A:258:HIS:HB2	1.98	0.46
1:A:63:PHE:HB3	1:A:98:LEU:HD13	1.98	0.45
1:B:280:MET:CG	1:B:298:ASP:HB2	2.46	0.45
1:B:451:GLU:HG2	1:B:476:VAL:HG22	1.98	0.45
1:A:280:MET:HE1	1:A:384:TYR:CE2	2.51	0.45
1:A:214:ARG:HE	1:B:624:ASP:CG	2.18	0.45
1:A:140:GLY:HA3	1:A:293:TRP:O	2.16	0.45
1:B:152:ILE:CG2	1:B:153:LEU:N	2.80	0.45
1:A:150:ARG:NH2	1:A:185:GLU:OE1	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:483:ALA:HB1	1:B:543:VAL:HB	1.98	0.45
1:A:183:SER:OG	1:A:185:GLU:HG2	2.17	0.44
1:B:323:TYR:HE1	1:B:343:CYS:SG	2.41	0.44
1:B:13:ARG:NH2	1:B:58:ARG:HG3	2.32	0.44
1:A:302:TYR:O	1:A:303:LEU:HB2	2.18	0.44
1:A:550:GLU:HG2	1:A:570:TYR:CE1	2.53	0.44
1:A:232:VAL:HG22	1:A:238:ILE:HD12	2.00	0.44
1:A:49:ARG:HB2	3:A:1379:HOH:O	2.17	0.44
1:A:152:ILE:CG2	1:A:153:LEU:N	2.80	0.44
1:A:247:VAL:HG21	1:A:324:LEU:CD1	2.46	0.44
1:A:247:VAL:HG11	1:A:344:MET:HE3	1.96	0.44
1:A:317:CYS:HB3	1:A:321:ILE:HG12	2.00	0.44
1:B:542:TRP:CG	1:B:567:LEU:HD13	2.53	0.43
1:A:319:GLY:C	1:A:321:ILE:HD12	2.39	0.43
1:A:238:ILE:HG13	1:A:344:MET:CE	2.49	0.43
1:A:133:ARG:HA	1:A:133:ARG:NE	2.33	0.43
1:A:247:VAL:CG2	1:A:324:LEU:HD11	2.46	0.43
1:A:328:ILE:HD11	1:A:336:ARG:NH2	2.34	0.43
1:A:484:VAL:CG1	1:A:539:LYS:HG3	2.48	0.43
1:A:22:GLU:O	1:A:26:ILE:HG13	2.19	0.43
1:A:590:LEU:HD21	1:A:607:VAL:HB	2.00	0.43
1:B:267:ARG:HG2	1:B:267:ARG:HH11	1.83	0.43
1:A:250:ASP:OD2	1:A:252:ARG:HB3	2.19	0.43
1:B:39:ILE:HD12	1:B:62:VAL:HG11	2.00	0.43
1:B:480:GLU:HB2	1:B:574:ASP:HA	2.00	0.42
1:A:267:ARG:NH2	3:A:1137:HOH:O	2.52	0.42
1:B:250:ASP:OD2	1:B:252:ARG:HB3	2.20	0.42
1:A:486:GLU:OE1	1:A:539:LYS:NZ	2.52	0.42
1:B:302:TYR:O	1:B:303:LEU:HB2	2.19	0.42
1:B:42:LEU:HD23	1:B:42:LEU:C	2.39	0.42
1:A:590:LEU:CD2	1:A:607:VAL:HB	2.50	0.42
1:A:238:ILE:HG13	1:A:344:MET:HE1	2.01	0.42
1:B:271:ILE:HG22	1:B:272:ILE:HG13	2.02	0.42
1:A:245:LEU:HD23	1:A:245:LEU:N	2.34	0.42
1:A:66:ASP:CG	1:A:70:ALA:HB3	2.41	0.41
1:B:103:GLU:O	1:B:106:GLU:HG2	2.20	0.41
1:B:513:LYS:O	1:B:611:LEU:HA	2.20	0.41
1:A:72:PRO:HB2	1:A:98:LEU:HD11	2.02	0.41
1:A:152:ILE:HG23	1:A:179:VAL:O	2.19	0.41
1:B:273:ASN:ND2	3:B:1079:HOH:O	2.40	0.41
1:B:370:ARG:HA	1:B:390:LEU:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:288:SER:HA	1:B:289:PRO:HD3	1.93	0.41
1:A:163:PRO:O	1:B:360:SER:HA	2.20	0.41
1:B:542:TRP:CD2	1:B:567:LEU:HD13	2.55	0.41
1:A:13:ARG:NH2	1:A:58:ARG:HG3	2.35	0.41
1:A:323:TYR:CE1	1:A:343:CYS:SG	3.14	0.41
1:B:143:GLU:CD	1:B:143:GLU:N	2.74	0.41
1:B:162:PHE:HB2	1:B:163:PRO:HD2	2.02	0.41
1:B:510:VAL:HB	1:B:616:PHE:HA	2.03	0.41
1:A:371:MET:HB3	1:A:390:LEU:HB2	2.03	0.41
1:B:281:VAL:HA	1:B:296:TYR:O	2.21	0.41
1:B:401:LYS:HG2	1:B:606:THR:HG22	2.02	0.40
1:B:134:VAL:O	1:B:136:PRO:HD3	2.20	0.40
1:A:507:ASN:HD22	1:A:507:ASN:HA	1.70	0.40
1:A:261:ALA:CB	1:A:268:LEU:HD11	2.51	0.40
1:A:115:ASP:OD1	1:A:117:ARG:HB2	2.22	0.40
1:A:241:GLU:O	1:A:242:LYS:CB	2.69	0.40
1:B:422:LEU:HD11	1:B:428:ALA:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	618/638 (97%)	582 (94%)	32 (5%)	4 (1%)	30	22
1	B	618/638 (97%)	590 (96%)	28 (4%)	0	100	100
All	All	1236/1276 (97%)	1172 (95%)	60 (5%)	4 (0%)	46	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	55	ALA
1	A	143	GLU
1	A	144	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	514/530 (97%)	509 (99%)	5 (1%)	82	85
1	B	514/530 (97%)	510 (99%)	4 (1%)	86	89
All	All	1028/1060 (97%)	1019 (99%)	9 (1%)	84	88

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	ASP
1	A	296	TYR
1	A	370	ARG
1	A	376	PHE
1	A	507	ASN
1	B	143	GLU
1	B	376	PHE
1	B	507	ASN
1	B	541	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	170	HIS
1	A	309	ASN
1	A	448	ASN
1	A	507	ASN
1	A	515	HIS
1	A	573	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	224	GLN
1	B	306	GLN
1	B	421	GLN
1	B	468	ASN
1	B	507	ASN
1	B	519	GLN
1	B	628	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	620/638 (97%)	-0.07	31 (5%) 32 34	16, 28, 48, 93	0
1	B	620/638 (97%)	-0.32	23 (3%) 45 47	13, 22, 40, 80	0
All	All	1240/1276 (97%)	-0.19	54 (4%) 38 39	13, 24, 45, 93	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	52	GLY	13.8
1	A	53	SER	11.3
1	B	52	GLY	10.0
1	B	51	ALA	9.2
1	B	54	GLU	9.2
1	A	148	ARG	9.1
1	A	51	ALA	9.0
1	A	54	GLU	8.6
1	B	55	ALA	8.5
1	A	145	ALA	8.4
1	A	55	ALA	7.7
1	B	53	SER	7.4
1	A	144	TYR	6.0
1	A	50	GLY	5.6
1	A	146	GLU	5.5
1	A	9	ALA	5.2
1	B	50	GLY	5.2
1	B	628	ASN	5.1
1	A	628	ASN	4.6
1	B	266	ASP	4.4
1	A	56	GLU	3.7
1	B	225	PRO	3.6
1	B	212	PRO	3.5
1	B	414	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	56	GLU	3.3
1	A	68	SER	3.3
1	B	207	PRO	3.2
1	A	339	ARG	3.1
1	A	225	PRO	3.0
1	B	462	PRO	3.0
1	A	266	ASP	2.9
1	A	462	PRO	2.9
1	A	208	GLU	2.8
1	A	141	VAL	2.7
1	B	461	GLY	2.7
1	A	507	ASN	2.7
1	A	414	GLY	2.6
1	A	212	PRO	2.6
1	B	413	GLU	2.6
1	A	142	PHE	2.5
1	A	149	GLY	2.4
1	A	413	GLU	2.4
1	A	35	PRO	2.4
1	B	215	THR	2.4
1	A	147	GLU	2.3
1	A	48	ALA	2.3
1	B	145	ALA	2.3
1	B	463	GLY	2.2
1	B	163	PRO	2.2
1	A	265	GLY	2.2
1	B	48	ALA	2.2
1	B	9	ALA	2.2
1	B	343	CYS	2.1
1	A	163	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CO	B	1002	1/1	0.99	0.04	-	49,49,49,49	0
2	CO	A	1001	1/1	0.98	0.04	-	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.