



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:00 PM GMT

PDB ID : 4WNW
Title : Human Cytochrome P450 2D6 Thioridazine Complex
Authors : Wang, A.; Stout, C.D.; Johnson, E.F.
Deposited on : 2014-10-14
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	<div> <div>5%</div> <div>80%</div> <div>16%</div> <div>.</div> </div>
1	B	479	<div> <div>5%</div> <div>74%</div> <div>20%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NA	B	603	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 2D6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	461	Total	C	N	O	S	0	0	0
			3652	2343	646	649	14			
1	B	452	Total	C	N	O	S	0	0	0
			3584	2296	634	640	14			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	initiating methionine	UNP P10635
A	24	ALA	-	expression tag	UNP P10635
A	25	LYS	-	expression tag	UNP P10635
A	26	LYS	-	expression tag	UNP P10635
A	27	THR	-	expression tag	UNP P10635
A	28	SER	-	expression tag	UNP P10635
A	29	SER	-	expression tag	UNP P10635
A	30	LYS	-	expression tag	UNP P10635
A	31	GLY	-	expression tag	UNP P10635
A	32	LYS	-	expression tag	UNP P10635
A	33	LEU	-	expression tag	UNP P10635
A	498	HIS	-	expression tag	UNP P10635
A	499	HIS	-	expression tag	UNP P10635
A	500	HIS	-	expression tag	UNP P10635
A	501	HIS	-	expression tag	UNP P10635
B	23	MET	-	initiating methionine	UNP P10635
B	24	ALA	-	expression tag	UNP P10635
B	25	LYS	-	expression tag	UNP P10635
B	26	LYS	-	expression tag	UNP P10635
B	27	THR	-	expression tag	UNP P10635
B	28	SER	-	expression tag	UNP P10635
B	29	SER	-	expression tag	UNP P10635
B	30	LYS	-	expression tag	UNP P10635
B	31	GLY	-	expression tag	UNP P10635
B	32	LYS	-	expression tag	UNP P10635

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	33	LEU	-	expression tag	UNP P10635
B	498	HIS	-	expression tag	UNP P10635
B	499	HIS	-	expression tag	UNP P10635
B	500	HIS	-	expression tag	UNP P10635
B	501	HIS	-	expression tag	UNP P10635

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 10-{2-[(2R)-1-methylpiperidin-2-yl]ethyl}-2-(methylsulfanyl)-10H-phenothiazine (three-letter code: RTZ) (formula: $C_{21}H_{26}N_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	S	0	1
			50	42	4	4		
3	B	1	Total	C	N	S	0	1
			50	42	4	4		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ni	0	0
			1	1		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

Chain B:

Amino Acid	Frequency (%)	Category
Met	~1.0	Grey
Ala	~1.0	Green
Lys	~1.0	Green
Lys	~1.0	Green
Thr	~1.0	Green
Ser	~1.0	Green
Ser	~1.0	Green
Lys	~1.0	Green
Gly	~1.0	Green
L33	~1.0	Green
P41	~1.0	Green
N45	~1.0	Green
H48	~1.0	Green
V49	~1.0	Green
D50	~1.0	Green
F51	~1.0	Green
D59	~1.0	Green
R62	~1.0	Green
V68	~1.0	Green
P77	~1.0	Green
L84	~1.0	Green
L91	~1.0	Green
D97	~1.0	Green
D100	~1.0	Green
R101	~1.0	Green
V104	~1.0	Green
P105	~1.0	Green
I106	~1.0	Green
L110	~1.0	Green
G111	~1.0	Green
F112	~1.0	Green
G118	~1.0	Green
V119	~1.0	Green
F120	~1.0	Green
L121	~1.0	Green
A122	~1.0	Green
R123	~1.0	Green
Y124	~1.0	Green
R132	~1.0	Green
V136	~1.0	Green
L142	~1.0	Green
G143	~1.0	Green
L144	~1.0	Green
G445	~1.0	Green
E446	~1.0	Green
R450	~1.0	Green
F454	~1.0	Green
L455	~1.0	Green
L460	~1.0	Green
L461	~1.0	Green
Q462	~1.0	Green
H463	~1.0	Green
F464	~1.0	Green
T470	~1.0	Green
G471	~1.0	Green
Q472	~1.0	Green
P487	~1.0	Green
S488	~1.0	Green
P489	~1.0	Green
L492	~1.0	Green
V495	~1.0	Green
P496	~1.0	Green
R497	~1.0	Green
HIS	~1.0	Grey
HIS	~1.0	Grey
HIS	~1.0	Grey
HIS	~1.0	Grey
A407	~1.0	Green
V408	~1.0	Green
N409	~1.0	Green
F410	~1.0	Green
K411	~1.0	Green
R414	~1.0	Green
F415	~1.0	Green
H416	~1.0	Green
H419	~1.0	Green
N285	~1.0	Green
P286	~1.0	Green
F290	~1.0	Green
L295	~1.0	Green
A305	~1.0	Green
V308	~1.0	Green
T309	~1.0	Green
T310	~1.0	Green
R316	~1.0	Green
L319	~1.0	Green
L320	~1.0	Green
G321	~1.0	Green
L322	~1.0	Green
L323	~1.0	Green
H324	~1.0	Green
P325	~1.0	Green
Q328	~1.0	Green
T339	~1.0	Green
G340	~1.0	Green
Q341	~1.0	Green
V342	~1.0	Green
R343	~1.0	Green
H352	~1.0	Green
V359	~1.0	Green
V363	~1.0	Green
S379	~1.0	Green
R380	~1.0	Green
D381	~1.0	Green
G386	~1.0	Green
F387	~1.0	Green
R388	~1.0	Green
T393	~1.0	Green
T397	~1.0	Green
A407	~1.0	Green
V408	~1.0	Green
N409	~1.0	Green
F410	~1.0	Green
K411	~1.0	Green
R414	~1.0	Green
F415	~1.0	Green
H416	~1.0	Green
H419	~1.0	Green
L421	~1.0	Green
D422	~1.0	Green
A423	~1.0	Green
Q424	~1.0	Green
G425	~1.0	Green
H426	~1.0	Green
F427	~1.0	Green
V428	~1.0	Green
G429	~1.0	Green
P430	~1.0	Green
E431	~1.0	Green
A432	~1.0	Green
S437	~1.0	Green
A438	~1.0	Green
G439	~1.0	Green
R440	~1.0	Green
L444	~1.0	Green
G445	~1.0	Green
E446	~1.0	Green
R450	~1.0	Green
F454	~1.0	Green
L455	~1.0	Green
L460	~1.0	Green
L461	~1.0	Green
Q462	~1.0	Green
H463	~1.0	Green
F464	~1.0	Green
T470	~1.0	Green
G471	~1.0	Green

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	194.19 Å 55.35 Å 145.22 Å 90.00° 134.19° 90.00°	Depositor
Resolution (Å)	32.98 – 3.30 32.98 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (32.98-3.30) 99.5 (32.98-3.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 3.32 Å)	Xtriage
Refinement program	phenix	Depositor
R, R_{free}	0.242 , 0.287 0.246 , 0.293	Depositor DCC
R_{free} test set	866 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	75.8	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.1	EDS
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 16992 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7425	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NA, NI, RTZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/3751	0.46	0/5104
1	B	0.22	0/3680	0.46	0/5005
All	All	0.22	0/7431	0.46	0/10109

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3652	0	3632	55	0
1	B	3584	0	3550	64	0
2	A	43	0	30	4	0
2	B	43	0	30	3	0
3	A	50	0	52	2	0
3	B	50	0	52	5	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	1	0	0	0	0
All	All	7425	0	7346	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:LEU:HD12	1:A:240:VAL:HG11	1.75	0.69
1:A:228:PRO:HB3	1:B:50:ASP:HB3	1.75	0.69
2:B:601:HEM:HBC2	2:B:601:HEM:HHD	1.76	0.68
1:A:273:GLU:HA	1:A:276:LEU:HD12	1.75	0.68
1:A:101:ARG:NH2	1:A:120:PHE:O	2.31	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/479 (95%)	428 (94%)	29 (6%)	0	100	100
1	B	446/479 (93%)	409 (92%)	35 (8%)	2 (0%)	39	76
All	All	903/958 (94%)	837 (93%)	64 (7%)	2 (0%)	52	85

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	68	VAL
1	B	106	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	395/409 (97%)	391 (99%)	4 (1%)	82	91
1	B	388/409 (95%)	385 (99%)	3 (1%)	86	93
All	All	783/818 (96%)	776 (99%)	7 (1%)	84	92

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	252	ASP
1	B	193	ARG
1	B	48	HIS
1	A	164	PHE
1	B	120	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	48	HIS
1	B	463	HIS
1	B	416	HIS
1	A	416	HIS
1	B	419	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	601	1	30,50,50	2.44	10 (33%)	24,82,82	2.31	7 (29%)
3	RTZ	A	602[A]	-	27,28,28	0.11	0	36,39,39	0.59	1 (2%)
3	RTZ	A	602[B]	-	27,28,28	0.14	0	36,39,39	0.57	1 (2%)
2	HEM	B	601	1	30,50,50	2.51	8 (26%)	24,82,82	2.20	6 (25%)
3	RTZ	B	602[A]	-	27,28,28	0.11	0	36,39,39	0.57	1 (2%)
3	RTZ	B	602[B]	-	27,28,28	0.11	0	36,39,39	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	601	1	-	0/10/54/54	0/0/8/8
3	RTZ	A	602[A]	-	-	0/7/30/30	0/4/4/4
3	RTZ	A	602[B]	-	-	0/7/30/30	0/4/4/4
2	HEM	B	601	1	-	0/10/54/54	0/0/8/8
3	RTZ	B	602[A]	-	-	0/7/30/30	0/4/4/4
3	RTZ	B	602[B]	-	-	0/7/30/30	0/4/4/4

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	HEM	C3B-C4B	-7.29	1.45	1.51
2	B	601	HEM	C2D-C3D	-7.01	1.33	1.54
2	A	601	HEM	C2D-C3D	-6.99	1.33	1.54
2	A	601	HEM	C3B-C4B	-6.81	1.45	1.51
2	B	601	HEM	C3D-C4D	-4.90	1.45	1.51

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602[A]	RTZ	CAL-CAP-CAG	-2.63	108.61	112.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602[A]	RTZ	CAL-CAP-CAG	-2.59	108.66	112.45
3	A	602[B]	RTZ	CAL-CAP-CAG	-2.32	109.05	112.45
2	A	601	HEM	C3B-C4B-CHC	2.13	126.17	123.16
2	B	601	HEM	CMD-C2D-C3D	2.40	124.95	114.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	HEM	4	0
3	A	602[A]	RTZ	1	0
3	A	602[B]	RTZ	1	0
2	B	601	HEM	3	0
3	B	602[A]	RTZ	3	0
3	B	602[B]	RTZ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	461/479 (96%)	0.39	25 (5%)	29 24	34, 66, 93, 114	0
1	B	452/479 (94%)	0.27	24 (5%)	30 24	37, 60, 88, 105	0
All	All	913/958 (95%)	0.33	49 (5%)	29 24	34, 63, 91, 114	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	232	HIS	4.7
1	A	234	PRO	4.0
1	B	472	GLN	3.6
1	A	474	ARG	3.5
1	B	97	ASP	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NA	B	603	1/1	0.93	0.57	4.47	71,71,71,71	0
2	HEM	A	601	43/43	0.97	0.27	0.13	30,36,42,48	0
3	RTZ	B	602[A]	25/25	0.87	0.28	-0.04	56,63,67,68	25
3	RTZ	B	602[B]	25/25	0.87	0.28	-0.07	58,62,66,68	25
3	RTZ	A	602[B]	25/25	0.93	0.25	-0.10	48,52,55,58	25
3	RTZ	A	602[A]	25/25	0.93	0.25	-0.22	48,53,55,64	25
2	HEM	B	601	43/43	0.97	0.23	-0.55	40,46,51,56	0
5	NI	B	604	1/1	0.98	0.04	-2.73	58,58,58,58	0
4	NA	A	603	1/1	0.92	0.61	-	49,49,49,49	0

6.5 Other polymers

There are no such residues in this entry.