



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:17 PM GMT

PDB ID : 3WRG
Title : The complex structure of HypBA1 with L-arabinose
Authors : Huang, C.H.; Zhu, Z.; Cheng, Y.S.; Chan, H.C.; Ko, T.P.; Chen, C.C.; Wang, I.; Ho, M.R.; Hsu, S.T.; Zeng, Y.F.; Huang, Y.N.; Liu, J.R.; Guo, R.T.
Deposited on : 2014-02-25
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1611 (2.26-2.22)
Clashscore	102246	1764 (2.26-2.22)
Ramachandran outliers	100387	1724 (2.26-2.22)
Sidechain outliers	100360	1724 (2.26-2.22)
RSRZ outliers	91569	1616 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	658	<div> <div>4%</div> <div>80%</div> <div>19%</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-reducing end beta-L-arabinofuranosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	658	5152	3243	884	1002	23	0	0	0

- Molecule 2 is SUGAR (BETA-L-ARABINOFURANOSE) (three-letter code: FUB) (formula: $C_5H_{10}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	5	5	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	308	Total 308	O 308	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Non-reducing end beta-L-arabinofuranosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	77.59 Å 77.59 Å 254.18 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.23 24.90 – 2.24	Depositor EDS
% Data completeness (in resolution range)	91.4 (25.00-2.23) 92.2 (24.90-2.24)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.08 (at 2.24 Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.204 , 0.244 0.201 , 0.243	Depositor DCC
R_{free} test set	2027 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	47.7	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.6	EDS
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 40524 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5471	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FUB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/5274	0.65	1/7173 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	30	ASN	N-CA-C	-7.09	91.86	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5152	0	4914	101	0
2	A	10	0	10	0	0
3	A	1	0	0	0	0
4	A	308	0	0	5	0
All	All	5471	0	4924	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:MET:HE2	1:A:192:ASP:H	0.99	1.08
1:A:118:LEU:H	1:A:147:MET:HE3	1.22	1.02
1:A:144:MET:HE2	1:A:192:ASP:N	1.79	0.95
1:A:118:LEU:N	1:A:147:MET:HE3	1.85	0.91
1:A:144:MET:CE	1:A:192:ASP:H	1.85	0.89
1:A:572:ASN:HD21	1:A:637:ARG:H	1.23	0.86
1:A:159:GLN:HE22	1:A:431:ARG:HH12	1.25	0.84
1:A:362:ALA:HB3	1:A:493:ARG:HD3	1.68	0.76
1:A:599:LEU:HD21	1:A:647:ARG:HD2	1.69	0.73
1:A:208:THR:HG22	1:A:210:GLU:H	1.53	0.73
1:A:469:TRP:CH2	1:A:602:VAL:HG22	2.24	0.72
1:A:4:THR:HB	1:A:452:THR:HG22	1.72	0.71
1:A:158:HIS:HE1	1:A:164:GLU:OE2	1.74	0.70
1:A:393:THR:HB	1:A:402:ARG:NH2	2.06	0.70
1:A:571:ASP:CG	1:A:647:ARG:HH22	1.94	0.69
1:A:515:LEU:N	1:A:515:LEU:HD12	2.07	0.69
1:A:270:HIS:HE1	1:A:322:GLU:OE2	1.77	0.68
1:A:27:GLY:O	1:A:30:ASN:O	2.12	0.68
1:A:392:THR:HG22	4:A:1105:HOH:O	1.95	0.67
1:A:393:THR:HB	1:A:402:ARG:HH22	1.60	0.67
1:A:393:THR:HG22	1:A:396:GLY:N	2.09	0.66
1:A:393:THR:HG22	1:A:396:GLY:H	1.61	0.66
1:A:312:THR:HG21	4:A:862:HOH:O	1.97	0.64
1:A:448:PHE:HA	1:A:463:GLN:HE22	1.64	0.63
1:A:139:GLN:HB2	1:A:231:TYR:CD1	2.33	0.63
1:A:393:THR:CG2	1:A:395:ASP:OD1	2.47	0.63
1:A:503:THR:HB	1:A:535:GLU:HB2	1.80	0.62
1:A:362:ALA:CB	1:A:493:ARG:HD3	2.30	0.61
1:A:595:GLN:HB2	1:A:598:LEU:HB3	1.82	0.61
1:A:571:ASP:OD1	1:A:647:ARG:NH2	2.34	0.61
1:A:29:MET:HG2	1:A:53:ALA:HB3	1.82	0.61
1:A:393:THR:HG23	1:A:395:ASP:OD1	2.01	0.60
1:A:515:LEU:HD12	1:A:515:LEU:H	1.66	0.60
1:A:302:LYS:O	1:A:306:THR:HB	2.02	0.59
1:A:10:TRP:O	1:A:14:ARG:HG3	2.02	0.59
1:A:277:LEU:O	1:A:281:VAL:HG23	2.02	0.58
1:A:159:GLN:HE22	1:A:431:ARG:NH1	2.00	0.57
1:A:158:HIS:HD2	1:A:207:GLU:OE2	1.87	0.57
1:A:613:HIS:HD2	1:A:630:ALA:HB1	1.70	0.56
1:A:538:MET:HB3	1:A:560:ARG:NH1	2.21	0.56
1:A:502:TYR:OH	1:A:515:LEU:HD11	2.07	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:MET:HG3	1:A:192:ASP:HB2	1.91	0.53
1:A:87:TYR:CD2	1:A:428:SER:HB3	2.44	0.53
1:A:264:GLN:O	1:A:299:ARG:NH1	2.42	0.53
1:A:14:ARG:NH2	1:A:430:ASP:OD1	2.42	0.52
1:A:159:GLN:NE2	1:A:431:ARG:HH12	1.99	0.52
1:A:118:LEU:HD23	1:A:119:ASP:N	2.24	0.51
1:A:525:ASN:O	1:A:528:ASP:HB2	2.11	0.51
1:A:147:MET:CE	1:A:173:MET:HG2	2.39	0.51
1:A:208:THR:HG22	1:A:210:GLU:N	2.25	0.51
1:A:475:TYR:HB2	1:A:532:ILE:HB	1.93	0.51
1:A:582:ALA:HB3	1:A:585:VAL:CG1	2.41	0.50
1:A:392:THR:HG22	1:A:393:THR:N	2.27	0.50
1:A:390:LEU:HG	1:A:654:VAL:HG21	1.93	0.49
1:A:572:ASN:HD21	1:A:637:ARG:N	2.03	0.49
1:A:582:ALA:HB3	1:A:585:VAL:HG11	1.95	0.49
1:A:469:TRP:HH2	1:A:602:VAL:HG22	1.76	0.49
1:A:229:GLN:O	1:A:233:LYS:HG3	2.12	0.49
1:A:20:SER:HB3	1:A:382:LYS:HE3	1.94	0.49
1:A:606:ASP:HB3	1:A:635:THR:CG2	2.43	0.49
1:A:507:ASN:ND2	1:A:531:GLU:H	2.09	0.49
1:A:331:PRO:O	1:A:335:MET:HB3	2.12	0.48
1:A:418:CYS:HB2	1:A:419:PRO:HD3	1.95	0.48
1:A:312:THR:HG23	1:A:314:ALA:H	1.78	0.48
1:A:639:VAL:HG13	1:A:640:PRO:HD2	1.94	0.48
1:A:117:TYR:HA	1:A:147:MET:HE3	1.97	0.47
1:A:538:MET:O	1:A:560:ARG:NH1	2.46	0.47
1:A:512:VAL:HG22	1:A:512:VAL:O	2.14	0.47
1:A:595:GLN:HG3	1:A:604:THR:HG23	1.97	0.47
1:A:379:LEU:H	1:A:646:ASN:HD21	1.60	0.47
1:A:75:ASP:HB2	1:A:146:VAL:HG21	1.96	0.46
1:A:239:ASN:OD1	1:A:239:ASN:O	2.34	0.46
1:A:167:LEU:O	1:A:171:LYS:HG3	2.15	0.46
1:A:147:MET:HE1	1:A:173:MET:HG2	1.97	0.45
1:A:642:TYR:C	1:A:642:TYR:CD1	2.90	0.45
1:A:452:THR:HG23	4:A:1036:HOH:O	2.16	0.45
1:A:198:GLU:HB2	4:A:810:HOH:O	2.16	0.45
1:A:590:ALA:HB2	1:A:607:LEU:HD21	1.99	0.45
1:A:379:LEU:H	1:A:646:ASN:ND2	2.15	0.44
1:A:335:MET:O	1:A:404:HIS:HE1	2.00	0.44
1:A:507:ASN:HD21	1:A:531:GLU:H	1.66	0.44
1:A:505:THR:OG1	1:A:533:ALA:HB3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:616:GLU:HA	1:A:616:GLU:OE1	2.18	0.43
1:A:448:PHE:CA	1:A:463:GLN:HE22	2.28	0.43
1:A:270:HIS:HD2	1:A:273:ARG:H	1.66	0.43
1:A:21:VAL:O	1:A:25:GLN:HG3	2.17	0.43
1:A:538:MET:HB3	1:A:560:ARG:HH12	1.84	0.43
1:A:467:PHE:CE2	1:A:562:PRO:HG3	2.55	0.42
1:A:8:PRO:O	1:A:12:ARG:HG3	2.20	0.42
1:A:75:ASP:HB2	1:A:146:VAL:CG2	2.50	0.41
1:A:648:GLU:HB2	4:A:900:HOH:O	2.20	0.41
1:A:131:ASP:O	1:A:133:PRO:HD3	2.20	0.41
1:A:459:LEU:HD12	1:A:478:SER:O	2.21	0.41
1:A:147:MET:HE2	1:A:173:MET:HG2	2.03	0.40
1:A:118:LEU:C	1:A:118:LEU:CD2	2.89	0.40
1:A:404:HIS:H	1:A:404:HIS:CD2	2.39	0.40
1:A:607:LEU:HA	1:A:608:PRO:HD3	1.95	0.40
1:A:71:MET:H	1:A:74:GLN:NE2	2.19	0.40
1:A:123:GLN:O	1:A:125:LYS:HG3	2.21	0.40
1:A:340:CYS:SG	1:A:418:CYS:HA	2.61	0.40
1:A:83:GLU:HB2	1:A:153:ALA:HA	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	656/658 (100%)	628 (96%)	23 (4%)	5 (1%)	24 20

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	196	GLU
1	A	483	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	484	THR
1	A	482	SER
1	A	512	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	538/538 (100%)	522 (97%)	16 (3%)	48	58

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	THR
1	A	19	GLU
1	A	28	VAL
1	A	29	MET
1	A	74	GLN
1	A	118	LEU
1	A	218	GLN
1	A	253	THR
1	A	293	LEU
1	A	393	THR
1	A	467	PHE
1	A	468	PRO
1	A	523	VAL
1	A	562	PRO
1	A	602	VAL
1	A	639	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	45	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	56	ASN
1	A	74	GLN
1	A	92	HIS
1	A	158	HIS
1	A	159	GLN
1	A	180	ASN
1	A	218	GLN
1	A	239	ASN
1	A	264	GLN
1	A	265	GLN
1	A	270	HIS
1	A	404	HIS
1	A	447	GLN
1	A	463	GLN
1	A	507	ASN
1	A	572	ASN
1	A	613	HIS
1	A	646	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FUB	A	701	-	10,10,10	0.68	0	12,14,14	1.37	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FUB	A	701	-	-	0/2/18/18	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	701	FUB	C1'-C2'-C3'	-3.62	97.66	102.45

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	658/658 (100%)	0.00	25 (3%)	44 46	36, 51, 69, 87	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	483	ALA	8.2
1	A	482	SER	5.9
1	A	484	THR	5.4
1	A	485	ASP	5.1
1	A	512	VAL	4.0
1	A	65	ASP	3.0
1	A	480	PRO	2.7
1	A	34	ASP	2.7
1	A	584	GLY	2.6
1	A	240	GLY	2.6
1	A	239	ASN	2.4
1	A	628	PRO	2.4
1	A	197	ILE	2.4
1	A	127	GLY	2.4
1	A	144	MET	2.3
1	A	481	ALA	2.3
1	A	513	GLY	2.3
1	A	185	GLU	2.3
1	A	128	VAL	2.2
1	A	445	SER	2.2
1	A	596	ALA	2.2
1	A	64	LEU	2.1
1	A	492	LEU	2.1
1	A	631	GLY	2.1
1	A	36	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FUB	A	701	10/10	0.96	0.12	-0.25	42,42,44,44	0
3	ZN	A	702	1/1	0.99	0.09	-	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.