



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2016 – 03:42 PM EST

PDB ID : 5WSI
Title : Crystal structure of ethanol modified signalling protein from buffalo (SPB-40) at 1.49 Å resolution
Authors : Singh, P.K.; Chaudhary, A.; Tyagi, T.K.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2016-12-07
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chitinase-3-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	361	2884	1846	499	530	9	0	2	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASP	deletion	UNP Q7YS85

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	8	6	2	0	0

- Molecule 3 is ETHANOL (three-letter code: EOH) (formula: C_2H_6O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	2	1		

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	412	Total 412	O 412	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chitinase-3-like protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.46 Å 66.52 Å 105.85 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.32 – 1.49 28.09 – 1.49	Depositor EDS
% Data completeness (in resolution range)	98.5 (56.32-1.49) 98.5 (28.09-1.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 1.50 Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.148 , 0.175 0.148 , 0.176	Depositor DCC
R_{free} test set	3479 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3335	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, EOH, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.80	57/2972 (1.9%)	1.71	64/4031 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (57) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	36	SER	CB-OG	-12.48	1.26	1.42
1	A	7	TYR	CZ-OH	11.10	1.56	1.37
1	A	56	TYR	CE2-CZ	-8.70	1.27	1.38
1	A	191	TRP	CE3-CZ3	8.63	1.53	1.38
1	A	9	SER	CA-CB	-8.47	1.40	1.52
1	A	91	LYS	CE-NZ	-8.44	1.27	1.49
1	A	81	GLY	N-CA	7.92	1.57	1.46
1	A	357	ASP	CB-CG	7.70	1.68	1.51
1	A	227	GLY	CA-C	-7.70	1.39	1.51
1	A	1	TYR	CB-CG	7.66	1.63	1.51
1	A	191	TRP	CE2-CZ2	7.58	1.52	1.39
1	A	303	GLN	CD-OE1	7.30	1.40	1.24
1	A	35	TYR	CE1-CZ	-7.25	1.29	1.38
1	A	174	SER	CB-OG	-7.12	1.32	1.42
1	A	361	GLY	CA-C	6.96	1.62	1.51
1	A	85	PHE	CD1-CE1	-6.95	1.25	1.39
1	A	44	GLU	CG-CD	-6.90	1.41	1.51
1	A	225	ARG	CG-CD	6.89	1.69	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	78	TRP	CB-CG	-6.82	1.38	1.50
1	A	6	TYR	CE2-CZ	-6.78	1.29	1.38
1	A	20	CYS	C-O	6.72	1.36	1.23
1	A	49	GLU	CD-OE1	-6.59	1.18	1.25
1	A	82	SER	CB-OG	6.58	1.50	1.42
1	A	63	LYS	CD-CE	-6.51	1.34	1.51
1	A	331	TRP	CE3-CZ3	6.33	1.49	1.38
1	A	84	ARG	CZ-NH2	-6.25	1.25	1.33
1	A	265	GLN	CD-OE1	6.17	1.37	1.24
1	A	35	TYR	CE2-CZ	-6.13	1.30	1.38
1	A	311[A]	GLU	CD-OE1	6.12	1.32	1.25
1	A	311[B]	GLU	CD-OE1	6.12	1.32	1.25
1	A	244	TYR	CE1-CZ	-6.02	1.30	1.38
1	A	280	ASP	CB-CG	-5.99	1.39	1.51
1	A	142	ARG	CZ-NH1	5.92	1.40	1.33
1	A	261	GLY	N-CA	5.87	1.54	1.46
1	A	222	TYR	CE2-CZ	-5.85	1.30	1.38
1	A	286	THR	CB-CG2	-5.79	1.33	1.52
1	A	94	SER	CB-OG	5.66	1.49	1.42
1	A	123	TRP	CZ3-CH2	-5.62	1.31	1.40
1	A	142	ARG	CD-NE	5.58	1.55	1.46
1	A	297	TYR	CE1-CZ	-5.58	1.31	1.38
1	A	119	LEU	C-O	5.44	1.33	1.23
1	A	303	GLN	CD-NE2	-5.41	1.19	1.32
1	A	300	LYS	N-CA	5.39	1.57	1.46
1	A	142	ARG	CZ-NH2	5.37	1.40	1.33
1	A	19	SER	CB-OG	5.36	1.49	1.42
1	A	85	PHE	CD2-CE2	5.32	1.49	1.39
1	A	142	ARG	NE-CZ	5.30	1.40	1.33
1	A	118	TRP	C-O	-5.29	1.13	1.23
1	A	15	GLU	CD-OE2	5.27	1.31	1.25
1	A	84	ARG	C-O	5.26	1.33	1.23
1	A	142	ARG	CG-CD	5.25	1.65	1.51
1	A	178	ASP	CG-OD1	5.24	1.37	1.25
1	A	225	ARG	NE-CZ	5.17	1.39	1.33
1	A	20	CYS	CB-SG	-5.16	1.73	1.81
1	A	118	TRP	CG-CD1	5.14	1.44	1.36
1	A	289	ARG	NE-CZ	5.07	1.39	1.33
1	A	63	LYS	CB-CG	-5.01	1.39	1.52

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	323	ARG	NE-CZ-NH1	-16.60	112.00	120.30
1	A	124	ARG	NE-CZ-NH2	16.45	128.53	120.30
1	A	225	ARG	NE-CZ-NH2	13.74	127.17	120.30
1	A	178	ASP	CB-CG-OD2	13.50	130.45	118.30
1	A	84	ARG	NE-CZ-NH2	-12.62	113.99	120.30
1	A	168	TYR	CB-CG-CD1	-9.56	115.26	121.00
1	A	107	ARG	NE-CZ-NH1	-9.49	115.55	120.30
1	A	175	ARG	NE-CZ-NH1	9.49	125.05	120.30
1	A	124	ARG	NE-CZ-NH1	-9.36	115.62	120.30
1	A	91	LYS	CD-CE-NZ	8.56	131.38	111.70
1	A	84	ARG	NE-CZ-NH1	8.48	124.54	120.30
1	A	252	ASP	CB-CG-OD2	-8.26	110.86	118.30
1	A	168	TYR	CB-CG-CD2	7.91	125.75	121.00
1	A	26	ASP	CB-CG-OD2	-7.90	111.19	118.30
1	A	142	ARG	NE-CZ-NH2	-7.87	116.36	120.30
1	A	252	ASP	CB-CG-OD1	7.80	125.33	118.30
1	A	84	ARG	O-C-N	-7.58	110.58	122.70
1	A	56	TYR	CB-CG-CD1	7.54	125.52	121.00
1	A	207	ASP	CB-CG-OD1	7.50	125.05	118.30
1	A	192	ARG	C-N-CA	-7.50	102.95	121.70
1	A	56	TYR	CB-CG-CD2	-7.41	116.55	121.00
1	A	142	ARG	CD-NE-CZ	7.32	133.85	123.60
1	A	127	ARG	NE-CZ-NH2	-7.25	116.67	120.30
1	A	26	ASP	CB-CG-OD1	7.19	124.77	118.30
1	A	280	ASP	CB-CG-OD1	6.71	124.34	118.30
1	A	95	ARG	NE-CZ-NH1	6.67	123.63	120.30
1	A	276	TYR	CZ-CE2-CD2	-6.60	113.86	119.80
1	A	336	ASP	CB-CG-OD2	-6.58	112.37	118.30
1	A	124	ARG	CD-NE-CZ	6.41	132.58	123.60
1	A	143	GLU	OE1-CD-OE2	-6.41	115.61	123.30
1	A	318	ARG	NE-CZ-NH1	-6.40	117.10	120.30
1	A	280	ASP	CB-CG-OD2	-6.40	112.54	118.30
1	A	300	LYS	CG-CD-CE	6.19	130.48	111.90
1	A	80	PHE	CB-CG-CD1	5.96	124.97	120.80
1	A	300	LYS	CD-CE-NZ	-5.95	98.01	111.70
1	A	235	MET	CG-SD-CE	-5.94	90.69	100.20
1	A	319	TYR	CB-CG-CD2	-5.93	117.44	121.00
1	A	60	ASN	CB-CA-C	5.93	122.25	110.40
1	A	289	ARG	NE-CZ-NH1	-5.86	117.37	120.30
1	A	17	ASP	CB-CG-OD2	-5.80	113.08	118.30
1	A	127	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	A	185	TYR	CB-CG-CD1	5.71	124.43	121.00
1	A	318	ARG	NE-CZ-NH2	5.66	123.13	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	52	ASP	CB-CG-OD1	5.66	123.39	118.30
1	A	210	SER	O-C-N	-5.64	113.67	122.70
1	A	3	LEU	O-C-N	5.60	131.67	122.70
1	A	307	TYR	CB-CG-CD1	5.57	124.34	121.00
1	A	52	ASP	CB-CG-OD2	-5.55	113.31	118.30
1	A	85	PHE	CG-CD1-CE1	5.54	126.90	120.80
1	A	68	LYS	CD-CE-NZ	5.50	124.36	111.70
1	A	323	ARG	NE-CZ-NH2	5.45	123.03	120.30
1	A	28	PHE	CB-CG-CD1	5.42	124.59	120.80
1	A	205	GLN	CB-CA-C	5.40	121.20	110.40
1	A	275	TYR	CB-CG-CD2	-5.36	117.78	121.00
1	A	142	ARG	CA-CB-CG	5.36	125.19	113.40
1	A	96	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	49	GLU	CG-CD-OE1	5.19	128.69	118.30
1	A	80	PHE	CB-CG-CD2	-5.18	117.17	120.80
1	A	7	TYR	CB-CG-CD2	-5.17	117.90	121.00
1	A	17	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	225	ARG	NE-CZ-NH1	-5.14	117.73	120.30
1	A	337	ASP	CB-CG-OD2	-5.12	113.69	118.30
1	A	23	ASP	CB-CG-OD2	-5.08	113.73	118.30
1	A	61	THR	OG1-CB-CG2	5.04	121.60	110.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	205	GLN	Peptide
1	A	209	SER	Peptide
1	A	210	SER	Mainchain
1	A	80	PHE	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2884	0	2811	16	0
2	A	8	0	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3	0	6	1	0
4	A	28	0	24	0	0
5	A	412	0	0	8	3
All	All	3335	0	2855	17	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:ASP:O	1:A:61:THR:HG23	1.92	0.69
1:A:310:GLN:HE22	1:A:345:GLN:HE22	1.40	0.67
1:A:198:HIS:CD2	1:A:198:HIS:H	2.16	0.64
1:A:262:ILE:H	1:A:303:GLN:HE22	1.46	0.61
1:A:357:ASP:HB2	5:A:680:HOH:O	2.05	0.57
1:A:315:ASN:HD22	1:A:318:ARG:HH11	1.53	0.57
1:A:198:HIS:HE1	5:A:586:HOH:O	1.90	0.55
1:A:269:GLU:OE2	5:A:501:HOH:O	2.17	0.55
1:A:262:ILE:H	1:A:303:GLN:NE2	2.08	0.52
1:A:70:LYS:HE2	5:A:597:HOH:O	2.09	0.51
1:A:188:HIS:HE1	1:A:194:THR:O	1.94	0.49
1:A:206:GLU:HA	1:A:207:ASP:HA	1.59	0.48
3:A:402:EOH:H11	5:A:561:HOH:O	2.14	0.47
1:A:91:LYS:HE2	5:A:782:HOH:O	2.15	0.47
1:A:286:THR:HG22	1:A:288:HIS:NE2	2.30	0.47
1:A:198:HIS:HD2	5:A:542:HOH:O	2.00	0.45
1:A:188:HIS:HD2	5:A:754:HOH:O	2.05	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:509:HOH:O	5:A:675:HOH:O[4_546]	2.03	0.17
5:A:675:HOH:O	5:A:831:HOH:O[4_446]	2.11	0.09
5:A:703:HOH:O	5:A:785:HOH:O[4_456]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/361 (100%)	350 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/302 (101%)	301 (99%)	3 (1%)	82	62

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	LYS
1	A	85	PHE
1	A	184	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	188	HIS
1	A	193	GLN
1	A	198	HIS
1	A	294	GLN
1	A	303	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	315	ASN
1	A	345	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MPD	A	401	-	6,7,7	1.00	0	6,10,10	1.43	1 (16%)
3	EOH	A	402	-	2,2,2	0.94	0	1,1,1	0.46	0
4	NAG	A	403	4	14,14,15	0.90	0	15,19,21	2.05	3 (20%)
4	NAG	A	404	1,4	14,14,15	3.10	6 (42%)	15,19,21	2.73	8 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	A	401	-	-	0/5/5/5	0/0/0/0
3	EOH	A	402	-	-	0/0/0/0	0/0/0/0
4	NAG	A	403	4	-	0/6/23/26	0/1/1/1
4	NAG	A	404	1,4	-	0/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	404	NAG	O5-C1	-6.93	1.32	1.43
4	A	404	NAG	C1-C2	-5.82	1.44	1.52
4	A	404	NAG	C4-C5	-3.88	1.44	1.53
4	A	404	NAG	C3-C2	2.24	1.57	1.52
4	A	404	NAG	O5-C5	3.18	1.50	1.43
4	A	404	NAG	O4-C4	4.39	1.53	1.43

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	404	NAG	O5-C5-C4	-5.64	100.79	110.13
4	A	403	NAG	O4-C4-C3	-4.73	99.69	110.36
4	A	404	NAG	C4-C3-C2	-4.39	104.52	111.34
4	A	404	NAG	C2-N2-C7	-3.98	117.93	123.11
4	A	403	NAG	C4-C3-C2	-3.94	105.22	111.34
4	A	404	NAG	C1-O5-C5	-3.14	107.52	112.14
4	A	404	NAG	O4-C4-C5	-2.92	101.53	109.23
4	A	404	NAG	O7-C7-C8	-2.41	117.64	122.07
4	A	404	NAG	O4-C4-C3	-2.12	105.57	110.36
2	A	401	MPD	C1-C2-C3	2.16	121.54	109.98
4	A	404	NAG	C6-C5-C4	2.44	119.11	112.99
4	A	403	NAG	C1-O5-C5	2.91	116.41	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	EOH	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	361/361 (100%)	0.32	26 (7%)	18 18	14, 23, 44, 112	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	362	VAL	11.7
1	A	207	ASP	10.9
1	A	209	SER	10.9
1	A	208	ALA	9.8
1	A	206	GLU	8.2
1	A	78	TRP	6.3
1	A	81	GLY	5.3
1	A	205	GLN	4.3
1	A	13	TYR	3.7
1	A	80	PHE	3.6
1	A	291	ARG	3.5
1	A	253	VAL	3.2
1	A	193	GLN	3.1
1	A	210	SER	3.1
1	A	20	CYS	3.0
1	A	15	GLU	3.0
1	A	79	ASN	2.8
1	A	361	GLY	2.8
1	A	16	GLY	2.8
1	A	82	SER	2.8
1	A	21	PHE	2.6
1	A	18	GLY	2.2
1	A	19	SER	2.1
1	A	238	PRO	2.0
1	A	204	GLY	2.0
1	A	252	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EOH	A	402	3/3	0.91	0.15	0.50	32,32,34,35	0
4	NAG	A	404	14/15	0.93	0.09	0.28	25,31,37,41	0
2	MPD	A	401	8/8	0.93	0.09	-0.64	22,25,27,28	0
4	NAG	A	403	14/15	0.82	0.35	-	40,53,60,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.