



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:23 AM GMT

PDB ID : 2WVH
Title : STRUCTURAL INSIGHTS INTO THE PRE-REACTION STATE OF
PYRUVATE DECARBOXYLASE FROM ZYMOMONAS MOBILIS
Authors : Pei, X.Y.; Erixon, K.; Luisi, B.F.; Leeper, F.J.
Deposited on : 2009-10-16
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	<div> <div>2%</div> <div>87% 12%</div> </div>
1	B	568	<div> <div>4%</div> <div>88% 11%</div> </div>
1	E	568	<div> <div>3%</div> <div>88% 11%</div> </div>
1	F	568	<div> <div>%</div> <div>86% 13%</div> </div>
1	V	568	<div> <div>3%</div> <div>88% 12%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	X	568	 4% 87% 12% •
1	Y	568	 2% 88% 11% •
1	Z	568	 3% 88% 11% •

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 38331 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PYRUVATE DECARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	566	Total	C	N	O	S	0	1	0
			4281	2724	730	808	19			
1	B	565	Total	C	N	O	S	0	1	0
			4276	2720	730	808	18			
1	E	565	Total	C	N	O	S	0	1	0
			4273	2719	729	807	18			
1	F	565	Total	C	N	O	S	0	1	0
			4273	2719	729	807	18			
1	V	565	Total	C	N	O	S	0	1	0
			4273	2719	729	807	18			
1	X	564	Total	C	N	O	S	0	1	0
			4264	2713	727	806	18			
1	Y	565	Total	C	N	O	S	0	1	0
			4273	2719	729	807	18			
1	Z	565	Total	C	N	O	S	0	1	0
			4273	2719	729	807	18			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	614	Total	O	0	0
			614	614		
2	B	479	Total	O	0	0
			479	479		
2	E	546	Total	O	0	0
			546	546		
2	F	522	Total	O	0	0
			522	522		
2	V	482	Total	O	0	0
			482	482		
2	X	486	Total	O	0	0
			486	486		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	Y	503	Total 503	O 503	0	0
2	Z	513	Total 513	O 513	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PYRUVATE DECARBOXYLASE





• Molecule 1: PYRUVATE DECARBOXYLASE





• Molecule 1: PYRUVATE DECARBOXYLASE







● Molecule 1: PYRUVATE DECARBOXYLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	70.90Å 111.17Å 167.12Å 89.89° 90.87° 101.27°	Depositor
Resolution (Å)	16.97 – 2.30 47.66 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.6 (16.97-2.30) 98.1 (47.66-2.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.08 (at 2.29Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.171 , 0.194 0.167 , 0.188	Depositor DCC
R_{free} test set	10929 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.5	EDS
Estimated twinning fraction	0.337 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 218204 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	38331	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/4377	0.46	0/5950
1	B	0.36	1/4369 (0.0%)	0.48	1/5940 (0.0%)
1	E	0.38	0/4369	0.46	0/5940
1	F	0.41	0/4369	0.48	0/5940
1	V	0.34	0/4369	0.46	0/5940
1	X	0.34	0/4360	0.46	0/5929
1	Y	0.35	0/4369	0.46	0/5940
1	Z	0.36	0/4369	0.46	0/5940
All	All	0.36	1/34951 (0.0%)	0.47	1/47519 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	200	GLU	CB-CG	-5.75	1.41	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	565	ASN	O-C-N	5.26	131.12	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4281	0	4255	58	0
1	B	4276	0	4242	56	0
1	E	4273	0	4243	53	0
1	F	4273	0	4243	68	0
1	V	4273	0	4243	62	0
1	X	4264	0	4230	69	0
1	Y	4273	0	4243	53	0
1	Z	4273	0	4243	57	0
2	A	614	0	0	13	0
2	B	479	0	0	13	0
2	E	546	0	0	12	0
2	F	522	0	0	25	0
2	V	482	0	0	15	0
2	X	486	0	0	11	0
2	Y	503	0	0	12	0
2	Z	513	0	0	12	0
All	All	38331	0	33942	429	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 429 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:72:THR:CG2	1:X:75:VAL:HG23	1.69	1.23
1:X:72:THR:HG23	1:X:75:VAL:CG2	1.77	1.14
1:X:72:THR:CG2	1:X:75:VAL:CG2	2.26	1.12
1:F:308:GLU:HG3	2:F:2343:HOH:O	1.52	1.07
1:F:380:PRO:HB2	1:X:431:GLU:HB3	1.43	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	565/568 (100%)	553 (98%)	12 (2%)	0	100	100
1	B	564/568 (99%)	554 (98%)	10 (2%)	0	100	100
1	E	564/568 (99%)	554 (98%)	10 (2%)	0	100	100
1	F	564/568 (99%)	553 (98%)	11 (2%)	0	100	100
1	V	564/568 (99%)	555 (98%)	9 (2%)	0	100	100
1	X	563/568 (99%)	553 (98%)	10 (2%)	0	100	100
1	Y	564/568 (99%)	554 (98%)	10 (2%)	0	100	100
1	Z	564/568 (99%)	555 (98%)	9 (2%)	0	100	100
All	All	4512/4544 (99%)	4431 (98%)	81 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	436/437 (100%)	435 (100%)	1 (0%)	95	98
1	B	435/437 (100%)	430 (99%)	5 (1%)	80	90
1	E	435/437 (100%)	433 (100%)	2 (0%)	92	97
1	F	435/437 (100%)	431 (99%)	4 (1%)	84	93
1	V	435/437 (100%)	434 (100%)	1 (0%)	95	98
1	X	434/437 (99%)	432 (100%)	2 (0%)	92	97
1	Y	435/437 (100%)	432 (99%)	3 (1%)	88	95
1	Z	435/437 (100%)	433 (100%)	2 (0%)	92	97
All	All	3480/3496 (100%)	3460 (99%)	20 (1%)	90	96

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	308	GLU
1	F	541	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Y	200	GLU
1	E	474	VAL
1	F	58	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	256	HIS
1	V	208	ASN
1	Z	208	ASN
1	F	323	HIS
1	F	565	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	566/568 (99%)	-0.15	13 (2%)	64	72	15, 21, 39, 65	1 (0%)
1	B	565/568 (99%)	0.01	24 (4%)	40	49	14, 27, 49, 73	1 (0%)
1	E	565/568 (99%)	-0.19	15 (2%)	58	67	15, 23, 41, 73	1 (0%)
1	F	565/568 (99%)	-0.25	8 (1%)	78	83	15, 22, 42, 71	1 (0%)
1	V	565/568 (99%)	-0.15	15 (2%)	58	67	15, 27, 48, 66	1 (0%)
1	X	564/568 (99%)	0.01	24 (4%)	39	48	15, 27, 48, 67	1 (0%)
1	Y	565/568 (99%)	-0.00	12 (2%)	67	74	16, 27, 45, 67	1 (0%)
1	Z	565/568 (99%)	-0.09	17 (3%)	54	63	16, 26, 45, 63	1 (0%)
All	All	4520/4544 (99%)	-0.10	128 (2%)	56	66	14, 25, 47, 73	8 (0%)

The worst 5 of 128 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	502	TYR	7.8
1	X	76	GLY	5.0
1	Y	502	TYR	4.9
1	X	75	VAL	4.8
1	A	475	MET	4.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.