



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:20 PM GMT

PDB ID : 3WXW
Title : Crystal structure of the human adiponectin receptor 2
Authors : Tanabe, H.; Fujii, Y.; Nakamura, Y.; Hosaka, T.; Okada-Iwabu, M.; Iwabu, M.; Kimura-Someya, T.; Shirouzu, M.; Yamauchi, T.; Kadowaki, T.; Yokoyama, S.
Deposited on : 2014-08-11
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	<div> <div>11%</div> <div> <div></div> <div>77%</div> <div>20%</div> <div></div> </div> </div>
2	H	119	<div> <div>5%</div> <div> <div></div> <div>80%</div> <div>20%</div> </div> </div>
3	L	107	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>14%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adiponectin receptor protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	283	2286	1534	369	366	17	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP Q86V24
A	-3	GLY	-	EXPRESSION TAG	UNP Q86V24
A	-2	SER	-	EXPRESSION TAG	UNP Q86V24
A	-1	GLU	-	EXPRESSION TAG	UNP Q86V24
A	0	PHE	-	EXPRESSION TAG	UNP Q86V24

- Molecule 2 is a protein called V region heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	119	927	592	149	182	4	0	0	0

- Molecule 3 is a protein called V region light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	107	820	517	137	163	3	0	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total 28	O 28	0	0
5	H	13	Total 13	O 13	0	0
5	L	11	Total 11	O 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Adiponectin receptor protein 2



• Molecule 2: V region heavy chain



• Molecule 3: V region light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	74.58Å 101.03Å 108.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.52 – 2.40 19.52 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.2 (19.52-2.40) 98.3 (19.52-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.41Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1690)	Depositor
R, R_{free}	0.248 , 0.290 0.248 , 0.290	Depositor DCC
R_{free} test set	1607 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	47.2	Xtriage
Anisotropy	0.653	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 64.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 32174 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4086	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2367	0.42	0/3213
2	H	0.25	0/951	0.43	0/1291
3	L	0.25	0/840	0.40	0/1140
All	All	0.25	0/4158	0.42	0/5644

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2286	0	2255	39	0
2	H	927	0	886	14	0
3	L	820	0	786	10	0
4	A	1	0	0	0	0
5	A	28	0	0	1	0
5	H	13	0	0	0	0
5	L	11	0	0	0	0
All	All	4086	0	3927	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:36:TYR:HE2	3:L:89:GLN:HG2	1.54	0.72
3:L:78:LEU:HD13	3:L:106:ILE:HD12	1.75	0.67
1:A:300:GLU:HB3	1:A:304:LYS:HD2	1.79	0.64
2:H:67:LYS:NZ	2:H:84:ARG:O	2.34	0.61
1:A:311:ILE:HG21	1:A:369:LEU:HD12	1.82	0.60
1:A:283:LEU:HD11	5:A:518:HOH:O	2.03	0.57
1:A:191:ALA:HA	1:A:226:LEU:HD13	1.87	0.57
1:A:127:PRO:HG3	2:H:101:GLY:HA3	1.88	0.56
1:A:166:TYR:OH	1:A:370:GLN:OE1	2.23	0.55
1:A:227:ILE:HG21	1:A:285:LEU:HG	1.88	0.55
1:A:0:PHE:HE1	3:L:93:SER:HB2	1.73	0.52
1:A:145:GLU:O	1:A:149:ILE:HG12	2.09	0.52
1:A:308:ILE:HG13	1:A:311:ILE:HG23	1.94	0.50
1:A:334:GLU:OE2	1:A:347:SER:N	2.42	0.50
1:A:156:CYS:SG	1:A:196:SER:OG	2.70	0.49
1:A:237:TYR:HB3	1:A:369:LEU:HD21	1.95	0.49
2:H:100:THR:OG1	2:H:101:GLY:N	2.42	0.49
1:A:239:PHE:HB2	1:A:246:CYS:HB2	1.95	0.49
1:A:320:LEU:HD13	1:A:358:GLY:HA2	1.95	0.49
1:A:316:LEU:HD22	1:A:361:VAL:HG11	1.96	0.48
1:A:313:TRP:CH2	1:A:362:HIS:HA	2.49	0.47
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.97	0.47
1:A:111:LEU:HD11	1:A:116:LYS:HG3	1.96	0.47
1:A:227:ILE:HD12	1:A:285:LEU:HD23	1.97	0.47
3:L:3:GLN:HB3	3:L:26:SER:HB3	1.98	0.46
3:L:15:VAL:HA	3:L:78:LEU:HD12	1.98	0.46
1:A:269:PHE:CZ	1:A:278:ARG:HG2	2.51	0.46
1:A:334:GLU:CD	1:A:347:SER:H	2.18	0.45
1:A:313:TRP:CZ3	1:A:362:HIS:HA	2.51	0.45
1:A:291:ILE:HB	1:A:292:PRO:HD3	1.97	0.45
1:A:301:GLY:H	1:A:304:LYS:HD2	1.81	0.45
2:H:103:ALA:HB3	3:L:91:PHE:HB2	1.98	0.44
2:H:71:THR:OG1	2:H:80:TYR:HB2	2.18	0.44
2:H:39:GLN:HB2	2:H:45:LEU:HD23	2.00	0.43
2:H:67:LYS:HE2	2:H:67:LYS:HB2	1.82	0.43
1:A:290:ILE:HG13	1:A:291:ILE:N	2.33	0.43
3:L:91:PHE:HA	3:L:96:TYR:CD1	2.53	0.43
1:A:0:PHE:O	1:A:102:ARG:HG3	2.18	0.43
1:A:111:LEU:HD13	1:A:115:LEU:HB3	2.00	0.43
1:A:266:TRP:O	1:A:267:ASP:HB3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:6:GLN:NE2	2:H:112:THR:HG23	2.34	0.43
1:A:373:ARG:NH2	1:A:376:ILE:HD11	2.34	0.42
2:H:51:PHE:CE2	2:H:72:VAL:HB	2.54	0.42
1:A:184:PHE:HE1	1:A:237:TYR:HE2	1.68	0.42
1:A:115:LEU:HA	1:A:115:LEU:HD12	1.80	0.41
1:A:176:ALA:HA	1:A:177:PRO:HD3	1.85	0.41
1:A:249:TYR:HA	1:A:252:VAL:HG22	2.01	0.41
2:H:43:LYS:HG2	2:H:44:SER:H	1.86	0.41
1:A:283:LEU:HA	1:A:283:LEU:HD12	1.63	0.41
2:H:20:ILE:HG23	2:H:112:THR:HG21	2.02	0.41
1:A:189:LEU:O	1:A:193:LEU:HG	2.20	0.41
1:A:175:VAL:HG21	1:A:370:GLN:NE2	2.35	0.41
3:L:33:LEU:HG	3:L:34:ALA:N	2.36	0.41
2:H:99:GLU:HB3	2:H:103:ALA:HA	2.02	0.40
1:A:158:PHE:HD1	1:A:159:PHE:CD1	2.39	0.40
3:L:35:TRP:CD2	3:L:73:LEU:HB2	2.56	0.40
1:A:180:GLU:HA	1:A:183:VAL:HG22	2.04	0.40
1:A:252:VAL:HA	1:A:255:VAL:HG12	2.02	0.40
3:L:35:TRP:CE2	3:L:73:LEU:HB2	2.57	0.40
2:H:104:TRP:CZ3	2:H:106:ALA:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/292 (95%)	261 (94%)	16 (6%)	0	100	100
2	H	117/119 (98%)	114 (97%)	3 (3%)	0	100	100
3	L	105/107 (98%)	102 (97%)	3 (3%)	0	100	100
All	All	499/518 (96%)	477 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/246 (98%)	239 (99%)	2 (1%)	86	94
2	H	99/99 (100%)	96 (97%)	3 (3%)	48	70
3	L	90/90 (100%)	87 (97%)	3 (3%)	45	66
All	All	430/435 (99%)	422 (98%)	8 (2%)	65	83

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	PHE
1	A	293	THR
2	H	21	THR
2	H	52	ASN
2	H	62	GLN
3	L	11	LEU
3	L	23	CYS
3	L	89	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	370	GLN
3	L	89	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/292 (96%)	0.77	33 (11%) 6 6	40, 69, 103, 145	0
2	H	119/119 (100%)	0.44	6 (5%) 32 33	45, 59, 79, 87	0
3	L	107/107 (100%)	0.15	2 (1%) 70 69	41, 56, 75, 87	0
All	All	509/518 (98%)	0.56	41 (8%) 15 14	40, 63, 93, 145	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	382	GLU	12.8
1	A	384	ASP	11.9
1	A	174	PHE	11.3
1	A	383	GLU	7.8
1	A	0	PHE	6.1
1	A	381	SER	6.0
1	A	-1	GLU	5.4
2	H	42	GLY	4.6
1	A	380	CYS	4.5
1	A	313	TRP	4.4
1	A	173	SER	4.1
1	A	375	MET	3.9
1	A	169	ARG	3.7
1	A	244	GLN	3.4
2	H	36	TRP	3.3
1	A	273	GLN	3.2
1	A	247	PHE	3.1
1	A	235	LEU	3.1
1	A	236	TYR	3.1
2	H	96	CYS	3.1
1	A	243	PRO	3.0
1	A	178	LEU	2.9
1	A	266	TRP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	168	PHE	2.8
1	A	307	THR	2.7
1	A	213	ARG	2.6
1	A	306	ALA	2.6
1	A	237	TYR	2.5
1	A	172	ILE	2.4
1	A	267	ASP	2.4
1	A	181	LYS	2.4
1	A	312	GLY	2.4
2	H	94	TYR	2.4
2	H	66	ASP	2.3
1	A	308	ILE	2.3
3	L	40	GLN	2.3
3	L	42	LYS	2.2
1	A	177	PRO	2.1
2	H	37	VAL	2.1
1	A	370	GLN	2.0
1	A	234	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ZN	A	401	1/1	0.97	0.20	1.22	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.