



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2016 – 04:50 PM EDT

PDB ID : 4X1T
Title : The crystal structure of Arabidopsis thaliana galactolipid synthase MGD1 in complex with UDP
Authors : Rocha, J.; Breton, C.
Deposited on : 2014-11-25
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	408	<div> <div>9%</div> <div>61%</div> <div>17%</div> <div>•</div> <div>21%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	602	-	-	-	X
3	EDO	A	605	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	606	-	-	-	X
3	EDO	A	607	-	-	-	X
3	EDO	A	608	-	-	-	X
3	EDO	A	609	-	-	-	X
3	EDO	A	610	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5230 atoms, of which 2638 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Monogalactosyldiacylglycerol synthase 1, chloroplastic.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	321	5047	1579	2573	432	445	18	0	2	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	534	ALA	-	expression tag	UNP O81770
A	535	ALA	-	expression tag	UNP O81770
A	536	ALA	-	expression tag	UNP O81770
A	537	LEU	-	expression tag	UNP O81770
A	538	GLU	-	expression tag	UNP O81770
A	539	HIS	-	expression tag	UNP O81770
A	540	HIS	-	expression tag	UNP O81770
A	541	HIS	-	expression tag	UNP O81770
A	542	HIS	-	expression tag	UNP O81770
A	543	HIS	-	expression tag	UNP O81770
A	544	HIS	-	expression tag	UNP O81770

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			36	9	11	2	12	2		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	57	Total	O	0	0
			57	57		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Monogalactosyldiacylglycerol synthase 1, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	139.25Å 139.25Å 38.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.58 – 2.25 45.58 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.0 (45.58-2.25) 99.4 (45.58-2.25)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 2.24Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.199 , 0.216 0.202 , 0.220	Depositor DCC
R_{free} test set	1045 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	51.2	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5230	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2528	0.49	2/3409 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	329	LYS	C-N-CD	5.92	140.83	128.40
1	A	297	CYS	C-N-CD	5.64	140.24	128.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	154	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2474	2573	2570	64	2
2	A	25	11	11	0	0
3	A	36	54	54	9	2
4	A	57	0	0	2	0
All	All	2592	2638	2635	65	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:LEU:HD13	1:A:289:HIS:ND1	1.80	0.97
1:A:150:ASP:O	1:A:152:GLY:N	2.11	0.84
1:A:156:ARG:HD3	1:A:160:GLU:OE2	1.77	0.84
1:A:259:LEU:CD1	1:A:289:HIS:ND1	2.46	0.79
1:A:290:LYS:O	1:A:315:GLN:NE2	2.18	0.75
1:A:156:ARG:CG	1:A:160:GLU:OE2	2.41	0.69
1:A:499:ASN:OD1	1:A:502:ARG:NH1	2.27	0.68
1:A:156:ARG:CD	1:A:160:GLU:OE2	2.42	0.67
1:A:156:ARG:HG3	1:A:160:GLU:OE2	1.97	0.65
1:A:257:VAL:HB	1:A:258:PRO:HD3	1.81	0.63
1:A:469:GLY:N	1:A:496:MET:HE2	2.15	0.62
1:A:147:LEU:HD13	1:A:254:MET:HB3	1.82	0.62
1:A:252:PRO:O	1:A:286:THR:OG1	2.12	0.60
1:A:151:THR:O	1:A:151:THR:HG22	2.01	0.59
1:A:371:ASP:HB3	3:A:609:EDO:H21	1.86	0.58
1:A:371:ASP:CB	3:A:609:EDO:H21	2.34	0.57
1:A:150:ASP:C	1:A:152:GLY:H	2.05	0.56
1:A:163:ARG:HG3	1:A:176:VAL:CG1	2.35	0.56
1:A:294:ARG:HG3	1:A:519:LEU:HD11	1.89	0.55
1:A:363:GLU:OE1	4:A:730:HOH:O	2.18	0.55
1:A:150:ASP:C	1:A:152:GLY:N	2.59	0.54
1:A:492:GLU:O	1:A:496:MET:HG3	2.07	0.54
1:A:252:PRO:O	1:A:286:THR:CG2	2.57	0.53
1:A:147:LEU:CD1	1:A:254:MET:HB3	2.39	0.52
1:A:259:LEU:HD13	1:A:289:HIS:CE1	2.45	0.52
1:A:268:LEU:HD23	1:A:269:LYS:N	2.24	0.51
1:A:332:ARG:NH1	1:A:340:GLU:OE1	2.42	0.51
1:A:163:ARG:HG3	1:A:176:VAL:HG13	1.93	0.51
1:A:143:LYS:HB2	1:A:244:PRO:HA	1.94	0.50
1:A:246:ILE:CD1	1:A:516:MET:HB3	2.43	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:VAL:HG12	1:A:276:VAL:HB	1.96	0.48
1:A:321:LEU:HD22	1:A:322:PRO:HD2	1.94	0.48
1:A:487:GLY:CA	3:A:610:EDO:H11	2.44	0.47
1:A:252:PRO:HD3	1:A:287:TRP:CZ2	2.50	0.47
1:A:251:HIS:ND1	1:A:252:PRO:HD2	2.30	0.46
1:A:394:LYS:HG2	3:A:604:EDO:H11	1.96	0.46
1:A:153:GLY:O	1:A:154:GLY:C	2.53	0.46
1:A:149:SER:HB3	1:A:251:HIS:CD2	2.51	0.46
1:A:147:LEU:CD1	1:A:254:MET:CB	2.94	0.46
1:A:398:LYS:HB2	3:A:604:EDO:H12	1.97	0.46
1:A:268:LEU:HD23	1:A:268:LEU:C	2.37	0.45
1:A:276:VAL:HG11	1:A:322:PRO:HG3	1.98	0.45
1:A:524:ASN:O	1:A:525:SER:HB2	2.16	0.44
1:A:272:VAL:HG22	1:A:523:LYS:HG3	1.99	0.44
1:A:163:ARG:HG3	1:A:176:VAL:HG11	1.99	0.44
1:A:339:ARG:NE	1:A:345:GLU:OE2	2.48	0.44
1:A:441:ILE:HA	1:A:504:ALA:HB1	2.01	0.43
1:A:325:PRO:HA	1:A:328:VAL:HG22	2.01	0.42
1:A:446:ILE:O	1:A:469:GLY:HA3	2.19	0.42
1:A:524:ASN:O	1:A:525:SER:CB	2.67	0.42
1:A:251:HIS:CE1	1:A:253:LEU:HB2	2.54	0.42
1:A:147:LEU:HD13	1:A:254:MET:SD	2.59	0.42
1:A:163:ARG:CG	1:A:176:VAL:HG11	2.50	0.42
1:A:320:GLY:CA	1:A:505:LYS:HD3	2.49	0.42
1:A:251:HIS:CG	1:A:252:PRO:HD2	2.55	0.41
1:A:507:GLU:HB2	3:A:602:EDO:H21	2.02	0.41
1:A:445:PRO:HG2	1:A:493:LEU:HD12	2.02	0.41
1:A:147:LEU:HD13	1:A:254:MET:CB	2.50	0.41
1:A:460:VAL:N	1:A:461:PRO:CD	2.84	0.41
1:A:400:SER:CB	3:A:608:EDO:H12	2.50	0.41
3:A:609:EDO:H11	4:A:714:HOH:O	2.20	0.41
1:A:303:ALA:HB2	1:A:318:VAL:HG22	2.03	0.40
1:A:487:GLY:HA2	3:A:610:EDO:H11	2.02	0.40
1:A:284:HIS:O	1:A:286:THR:N	2.55	0.40
1:A:469:GLY:N	1:A:496:MET:CE	2.83	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:THR:CG2	3:A:606:EDO:O2[2_565]	2.00	0.20

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:THR:HG21	3:A:606:EDO:O2[2_565]	1.45	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	319/408 (78%)	304 (95%)	14 (4%)	1 (0%)	46 52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	285	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/343 (79%)	265 (98%)	5 (2%)	65 75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150	ASP
1	A	156	ARG
1	A	329	LYS
1	A	330	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	524	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UDP	A	601	-	19,26,26	0.99	0	22,40,40	0.82	0
3	EDO	A	602	-	3,3,3	0.44	0	2,2,2	0.37	0
3	EDO	A	603	-	3,3,3	0.44	0	2,2,2	0.38	0
3	EDO	A	604	-	3,3,3	0.41	0	2,2,2	0.49	0
3	EDO	A	605	-	3,3,3	0.46	0	2,2,2	0.32	0
3	EDO	A	606	-	3,3,3	0.31	0	2,2,2	0.35	0
3	EDO	A	607	-	3,3,3	0.44	0	2,2,2	0.36	0
3	EDO	A	608	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	A	609	-	3,3,3	0.35	0	2,2,2	0.29	0
3	EDO	A	610	-	3,3,3	0.45	0	2,2,2	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	601	-	-	0/12/32/32	0/2/2/2
3	EDO	A	602	-	-	0/1/1/1	0/0/0/0
3	EDO	A	603	-	-	0/1/1/1	0/0/0/0
3	EDO	A	604	-	-	0/1/1/1	0/0/0/0
3	EDO	A	605	-	-	0/1/1/1	0/0/0/0
3	EDO	A	606	-	-	0/1/1/1	0/0/0/0
3	EDO	A	607	-	-	0/1/1/1	0/0/0/0
3	EDO	A	608	-	-	0/1/1/1	0/0/0/0
3	EDO	A	609	-	-	0/1/1/1	0/0/0/0
3	EDO	A	610	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	EDO	1	0
3	A	604	EDO	2	0
3	A	606	EDO	0	2
3	A	608	EDO	1	0
3	A	609	EDO	3	0
3	A	610	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/408 (78%)	0.80	38 (11%) 6 6	19, 52, 116, 142	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	ILE	10.6
1	A	267	LEU	10.5
1	A	262	LEU	8.9
1	A	261	VAL	7.7
1	A	521	ARG	7.7
1	A	331	VAL	7.5
1	A	523	LYS	6.7
1	A	265	LYS	6.0
1	A	524	ASN	5.5
1	A	142	LYS	5.4
1	A	269	LYS	5.0
1	A	264	SER	4.7
1	A	143	LYS	4.4
1	A	270	LYS	4.3
1	A	266	GLY	4.0
1	A	525	SER	3.9
1	A	257	VAL	3.9
1	A	152	GLY	3.8
1	A	243	GLN	3.8
1	A	245	ASP	3.8
1	A	268	LEU	3.8
1	A	174	TYR	3.7
1	A	378	LEU	3.7
1	A	244	PRO	3.7
1	A	522	LYS	3.5
1	A	259	LEU	3.3
1	A	177	PHE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	HIS	3.1
1	A	246	ILE	2.9
1	A	260	ARG	2.7
1	A	153	GLY	2.6
1	A	288	PHE	2.6
1	A	291	LEU	2.6
1	A	254	MET	2.4
1	A	263	ARG	2.3
1	A	144	VAL	2.3
1	A	453	ALA	2.2
1	A	176	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	A	610	4/4	0.64	0.55	29.49	51,62,67,74	0
3	EDO	A	609	4/4	0.61	0.33	13.54	43,55,64,66	0
3	EDO	A	606	4/4	0.74	0.38	4.80	59,71,78,80	0
3	EDO	A	608	4/4	0.73	0.24	4.78	58,69,74,75	0
3	EDO	A	605	4/4	0.89	0.29	4.21	59,71,78,81	0
3	EDO	A	607	4/4	0.82	0.24	3.55	56,73,83,88	0
3	EDO	A	602	4/4	0.91	0.24	3.21	50,60,62,63	0
2	UDP	A	601	25/25	0.95	0.15	0.23	35,38,65,73	0
3	EDO	A	603	4/4	0.90	0.14	-1.27	45,54,60,63	0
3	EDO	A	604	4/4	0.81	0.33	-	63,76,84,88	0

6.5 Other polymers

There are no such residues in this entry.