



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:59 PM GMT

PDB ID : 3X3Y  
Title : Copper amine oxidase from *Arthrobacter globiformis* anaerobically reduced by histamine  
Authors : Okajima, T.; Nakanishi, S.; Murakawa, T.; Kataoka, M.; Hayashi, H.; Hamaguchi, A.; Nakai, T.; Kawano, Y.; Yamaguchi, H.; Tanizawa, K.  
Deposited on : 2015-03-10  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	620	<div> <div>2%</div> <div>93%</div> <div>7%</div> </div>
1	B	620	<div> <div>3%</div> <div>93%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	702	-	-	-	X
3	GOL	A	704	-	-	-	X
3	GOL	A	705	-	-	-	X
3	GOL	A	706	-	-	-	X
3	GOL	A	707	-	-	-	X
3	GOL	A	708	-	-	-	X
3	GOL	B	702	-	-	-	X
3	GOL	B	703	-	-	-	X
3	GOL	B	704	-	-	-	X
4	NA	B	706	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11310 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylethylamine oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	620	Total	C	N	O	S	0	17	0
			5008	3159	885	953	11			
1	B	620	Total	C	N	O	S	0	16	0
			5001	3151	883	957	10			

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cu	0	0
			1	1		
2	A	1	Total	Cu	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Na 1 1	0	0
4	A	1	Total Na 1 1	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total K 1 1	0	0
5	A	1	Total K 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	661	Total 661	O 661	0	0
6	B	568	Total 568	O 568	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylethylamine oxidase



- Molecule 1: Phenylethylamine oxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	193.47Å 63.25Å 157.91Å 90.00° 117.73° 90.00°	Depositor
Resolution (Å)	36.46 – 1.50 43.04 – 1.50	Depositor EDS
% Data completeness (in resolution range)	97.5 (36.46-1.50) 91.0 (43.04-1.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.66 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.162 , 0.177 0.158 , 0.172	Depositor DCC
$R_{free}$ test set	12384 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.6	Xtriage
Anisotropy	0.975	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 51.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 264317 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11310	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.29 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5897e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, K, TYQ, CU, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/5115	0.55	0/6962
1	B	0.31	0/5108	0.53	0/6951
All	All	0.32	0/10223	0.54	0/13913

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5008	0	4828	47	0
1	B	5001	0	4806	34	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	42	0	56	0	0
3	B	24	0	32	4	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	661	0	0	12	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	568	0	0	7	0
All	All	11310	0	9722	81	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:GLU:HG3	1:A:55:ALA:N	1.81	0.93
1:A:267:ARG:NH1	6:A:1289:HOH:O	2.06	0.87
1:B:312:GLU:OE1	6:B:1296:HOH:O	1.94	0.85
1:B:188:ARG:NH1	6:B:1329:HOH:O	2.06	0.85
1:A:54:GLU:HG3	1:A:55:ALA:H	1.39	0.84
1:A:71[B]:ARG:NH1	6:A:1403:HOH:O	2.12	0.81
1:A:501:GLU:OE2	6:A:1364:HOH:O	2.02	0.78
1:A:54:GLU:CG	1:A:55:ALA:H	2.00	0.73
1:A:54:GLU:CG	1:A:55:ALA:N	2.54	0.71
1:A:13:ARG:NH2	1:A:56:GLU:OE2	2.18	0.69
1:A:36:GLU:OE1	6:A:1402:HOH:O	2.10	0.69
1:B:422:LEU:HD11	1:B:428:ALA:HB2	1.74	0.68
1:A:238:ILE:HG21	1:A:371[B]:MET:HE2	1.76	0.67
1:B:599:TRP:CD2	1:B:600:PRO:HA	2.29	0.67
1:A:599:TRP:CD2	1:A:600:PRO:HA	2.29	0.67
1:B:482:GLU:OE1	6:B:1239:HOH:O	2.14	0.65
1:A:382:TYQ:HD2	1:A:403:THR:O	1.96	0.65
1:B:20:ILE:HD12	1:B:327[A]:VAL:HG12	1.79	0.65
1:B:59:ARG:NE	1:B:85[A]:ILE:HD13	2.12	0.65
1:A:200:GLU:OE1	6:A:1337:HOH:O	2.13	0.64
1:B:280:MET:HG3	1:B:298:ASP:HB2	1.80	0.64
1:B:54:GLU:HG2	1:B:55:ALA:O	1.98	0.63
1:A:240:TRP:HB2	1:A:371[B]:MET:HE1	1.80	0.63
1:A:62[B]:VAL:HG22	1:A:75:VAL:HB	1.83	0.60
1:B:103:GLU:HB3	6:B:1326:HOH:O	2.03	0.59
1:B:36[A]:GLU:OE2	6:B:1226:HOH:O	2.17	0.59
1:B:490[B]:ARG:NH2	6:B:1305:HOH:O	2.35	0.59
1:A:263:ARG:NH2	6:A:1354:HOH:O	2.36	0.58
1:B:382:TYQ:HD2	1:B:403:THR:O	2.03	0.58
1:B:459:THR:HA	3:B:703:GOL:H31	1.86	0.58
1:B:59:ARG:CZ	1:B:85[A]:ILE:HD13	2.34	0.58
1:A:188[B]:ARG:NH1	6:A:1300:HOH:O	2.37	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:SER:HB2	1:B:85[A]:ILE:HD11	1.86	0.57
1:A:422:LEU:HD11	1:A:428:ALA:HB2	1.87	0.56
3:B:705:GOL:H31	6:B:1347:HOH:O	2.06	0.55
1:A:23:VAL:HG22	1:A:79[B]:VAL:HG12	1.89	0.55
1:A:382:TYQ:OZ	6:A:1188:HOH:O	2.15	0.55
1:B:280:MET:CG	1:B:298:ASP:HB2	2.37	0.54
1:B:20:ILE:CD1	1:B:327[B]:VAL:HG23	2.38	0.53
1:A:28:ARG:NH1	6:A:1201:HOH:O	2.32	0.53
1:A:23:VAL:HG22	1:A:79[A]:VAL:HG22	1.89	0.53
1:A:23:VAL:HG22	1:A:79[B]:VAL:CG1	2.39	0.53
1:A:23:VAL:CG2	1:A:79[B]:VAL:CG1	2.87	0.53
1:A:251:VAL:CG2	1:A:306:GLN:HB3	2.40	0.52
1:A:230:PHE:HB3	1:A:371[B]:MET:HE1	1.92	0.51
1:A:238:ILE:HG21	1:A:371[B]:MET:CE	2.43	0.49
1:B:20:ILE:HD12	1:B:327[A]:VAL:CG1	2.42	0.49
1:B:263[B]:ARG:HG2	1:B:268:LEU:HD13	1.95	0.49
1:A:365:THR:HB	1:B:315[B]:CYS:SG	2.52	0.49
1:B:222:ILE:HD12	1:B:318:LEU:HD23	1.94	0.48
1:A:238:ILE:CG2	1:A:371[B]:MET:CE	2.91	0.48
1:B:71:ARG:NH1	1:B:91:ASP:OD2	2.46	0.48
1:A:71[B]:ARG:HE	1:A:89:GLU:HG2	1.79	0.48
1:B:20:ILE:CD1	1:B:327[A]:VAL:HG12	2.44	0.48
1:A:238:ILE:CG2	1:A:371[B]:MET:HE2	2.42	0.47
1:A:230:PHE:CB	1:A:371[B]:MET:HE1	2.45	0.47
1:B:458:GLN:O	3:B:703:GOL:H31	2.15	0.47
1:B:51:ALA:HB1	1:B:54:GLU:OE2	2.15	0.46
1:A:60:PHE:CD1	1:A:79[A]:VAL:HG21	2.51	0.46
1:B:36[A]:GLU:HG3	1:B:66:ASP:OD2	2.16	0.46
1:B:20:ILE:HD12	1:B:327[B]:VAL:HG23	1.99	0.45
1:A:382:TYQ:N5	1:A:602:MET:SD	2.90	0.45
1:A:146[B]:GLU:CD	1:A:146[B]:GLU:H	2.19	0.45
1:A:62[B]:VAL:CG2	1:A:75:VAL:HB	2.45	0.45
1:A:491[B]:THR:HG23	6:A:1119:HOH:O	2.17	0.45
1:B:10:SER:HA	1:B:11:PRO:HD3	1.88	0.44
1:A:60:PHE:CD1	1:A:79[A]:VAL:CG2	3.00	0.44
1:B:20:ILE:HD11	1:B:327[B]:VAL:HG23	1.99	0.44
1:A:71[B]:ARG:HG2	6:A:906:HOH:O	2.17	0.43
1:B:107:VAL:HG12	1:B:111:LEU:HD12	2.00	0.43
1:A:350:GLY:HA2	1:A:367:ARG:NH2	2.33	0.43
1:A:280:MET:CG	1:A:298:ASP:HB2	2.48	0.43
1:A:23:VAL:CG2	1:A:79[B]:VAL:HG12	2.49	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:505:ARG:HD3	1:A:618:ASP:HB3	2.01	0.42
1:B:26:ILE:HD11	1:B:82:GLY:HA2	2.02	0.42
1:B:391:TYR:CE2	3:B:702:GOL:H2	2.55	0.41
1:B:599:TRP:CG	1:B:600:PRO:HA	2.56	0.41
1:A:556:ASP:OD1	6:A:1191:HOH:O	2.22	0.40
1:A:42:LEU:HD23	1:A:42:LEU:C	2.41	0.40
1:A:232:VAL:HG22	1:A:238:ILE:CD1	2.52	0.40
1:A:280:MET:HG3	1:A:298:ASP:HB2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1358:HOH:O	6:A:1403:HOH:O[4_556]	2.12	0.08

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	634/620 (102%)	614 (97%)	20 (3%)	0	100	100
1	B	633/620 (102%)	612 (97%)	21 (3%)	0	100	100
All	All	1267/1240 (102%)	1226 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	530/513 (103%)	528 (100%)	2 (0%)	93	84
1	B	529/513 (103%)	525 (99%)	4 (1%)	86	70
All	All	1059/1026 (103%)	1053 (99%)	6 (1%)	90	78

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	336	ARG
1	A	376	PHE
1	B	54	GLU
1	B	56	GLU
1	B	376	PHE
1	B	541	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TYQ	A	382	1	12,14,15	3.62	4 (33%)	15,19,21	1.29	2 (13%)
1	TYQ	B	382	1,2	12,14,15	3.58	5 (41%)	15,19,21	1.29	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TYQ	A	382	1	-	0/4/6/8	0/1/1/1
1	TYQ	B	382	1,2	-	0/4/6/8	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	382	TYQ	CD1-CG	-2.38	1.36	1.40
1	B	382	TYQ	CD1-CG	-2.06	1.37	1.40
1	B	382	TYQ	CD2-CE2	-2.02	1.37	1.40
1	B	382	TYQ	CD2-CG	5.58	1.49	1.39
1	A	382	TYQ	CD2-CG	5.78	1.49	1.39
1	B	382	TYQ	CE2-CZ	6.76	1.49	1.40
1	A	382	TYQ	CE2-CZ	6.88	1.49	1.40
1	A	382	TYQ	CE1-CD1	7.69	1.49	1.38
1	B	382	TYQ	CE1-CD1	7.74	1.49	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	382	TYQ	CE2-CD2-CG	-3.31	119.56	122.44
1	A	382	TYQ	CE2-CD2-CG	-2.78	120.02	122.44
1	B	382	TYQ	OH-CZ-CE2	2.30	120.01	116.36
1	A	382	TYQ	OH-CZ-CE2	2.83	120.85	116.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	382	TYQ	3	0
1	B	382	TYQ	1	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	702	-	5,5,5	0.43	0	5,5,5	0.61	0
3	GOL	A	703	-	5,5,5	0.39	0	5,5,5	0.24	0
3	GOL	A	704	-	5,5,5	0.35	0	5,5,5	0.23	0
3	GOL	A	705	-	5,5,5	0.38	0	5,5,5	0.24	0
3	GOL	A	706	-	5,5,5	0.41	0	5,5,5	0.20	0
3	GOL	A	707	-	5,5,5	0.41	0	5,5,5	0.38	0
3	GOL	A	708	-	5,5,5	0.37	0	5,5,5	0.95	0
3	GOL	B	702	-	5,5,5	0.37	0	5,5,5	0.31	0
3	GOL	B	703	-	5,5,5	0.36	0	5,5,5	0.83	0
3	GOL	B	704	-	5,5,5	0.39	0	5,5,5	0.41	0
3	GOL	B	705	-	5,5,5	0.38	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	702	-	-	0/4/4/4	0/0/0/0
3	GOL	A	703	-	-	0/4/4/4	0/0/0/0
3	GOL	A	704	-	-	0/4/4/4	0/0/0/0
3	GOL	A	705	-	-	0/4/4/4	0/0/0/0
3	GOL	A	706	-	-	0/4/4/4	0/0/0/0
3	GOL	A	707	-	-	0/4/4/4	0/0/0/0
3	GOL	A	708	-	-	0/4/4/4	0/0/0/0
3	GOL	B	702	-	-	0/4/4/4	0/0/0/0
3	GOL	B	703	-	-	0/4/4/4	0/0/0/0
3	GOL	B	704	-	-	0/4/4/4	0/0/0/0
3	GOL	B	705	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	702	GOL	1	0
3	B	703	GOL	2	0
3	B	705	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	619/620 (99%)	-0.15	15 (2%) 62 66	6, 14, 27, 67	0
1	B	619/620 (99%)	-0.06	17 (2%) 58 61	8, 18, 33, 70	0
All	All	1238/1240 (99%)	-0.10	32 (2%) 59 63	6, 15, 32, 70	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	53	SER	11.7
1	A	51	ALA	8.1
1	B	52	GLY	7.8
1	B	51	ALA	7.7
1	A	52	GLY	7.5
1	B	53	SER	7.0
1	B	54	GLU	6.7
1	A	55	ALA	5.3
1	B	55	ALA	5.1
1	A	50	GLY	5.0
1	A	54	GLU	4.9
1	B	313	LEU	4.6
1	B	50	GLY	4.6
1	B	266	ASP	4.3
1	A	315[A]	CYS	3.7
1	B	314	GLY	3.6
1	A	313	LEU	3.5
1	B	9	ALA	3.3
1	A	628	ASN	3.3
1	B	628	ASN	3.2
1	B	315[A]	CYS	3.2
1	B	48	ALA	3.1
1	A	225	PRO	2.9
1	A	266	ASP	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	56	GLU	2.5
1	B	462	PRO	2.5
1	A	208	GLU	2.4
1	A	56	GLU	2.2
1	A	215	THR	2.0
1	A	48	ALA	2.0
1	B	267	ARG	2.0
1	B	80	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TYQ	A	382	14/15	0.95	0.11	-	9,16,28,33	0
1	TYQ	B	382	14/15	0.97	0.08	-	9,17,29,30	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	708	6/6	0.67	0.22	23.47	39,40,42,43	0
3	GOL	A	707	6/6	0.62	0.27	20.93	34,46,50,51	0
3	GOL	B	703	6/6	0.43	0.33	11.46	47,54,57,58	0
3	GOL	A	706	6/6	0.81	0.21	9.57	29,38,39,40	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	B	702	6/6	0.80	0.17	4.94	29,35,37,45	0
3	GOL	A	705	6/6	0.78	0.17	3.98	26,31,36,43	0
3	GOL	A	704	6/6	0.83	0.15	3.17	24,27,28,29	0
3	GOL	B	704	6/6	0.88	0.13	2.89	25,27,27,31	0
3	GOL	A	702	6/6	0.89	0.17	2.83	18,25,32,34	0
4	NA	B	706	1/1	0.99	0.16	2.78	17,17,17,17	0
4	NA	A	709	1/1	1.00	0.10	1.17	7,7,7,7	0
3	GOL	A	703	6/6	0.99	0.06	-0.10	9,10,10,10	0
5	K	A	710	1/1	0.98	0.07	-0.21	16,16,16,16	0
5	K	B	707	1/1	0.98	0.05	-1.46	38,38,38,38	0
3	GOL	B	705	6/6	0.74	0.39	-	59,62,64,64	0
2	CU	A	701	1/1	1.00	0.06	-	11,11,11,11	0
2	CU	B	701	1/1	1.00	0.07	-	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.