



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:28 AM GMT

PDB ID : 2X85  
Title : Tailspike protein of E. coli bacteriophage HK620 in complex with hexasaccharide  
Authors : Lorenzen, N.K.; Mueller, J.J.; Heinemann, U.; Seckler, R.; Barbirz, S.  
Deposited on : 2010-03-05  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NA	A	1710	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NA	A	1713	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TAILSPIKE PROTEIN HK620.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	597	Total	C	N	O	S	0	23	0
			4607	2890	787	906	24			

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Na	0	0
			4	4		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	6	Total	C	N	O	0	0
			72	40	2	30		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	691	Total	O	0	0
			691	691		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TAILSPIKE PROTEIN HK620



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.25Å 74.25Å 174.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.13 – 1.50 36.31 – 1.50	Depositor EDS
% Data completeness (in resolution range)	90.4 (37.13-1.50) 90.0 (36.31-1.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.02 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.165 , 0.198 0.185 , 0.216	Depositor DCC
$R_{free}$ test set	4087 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	12.0	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 35.3	EDS
Estimated twinning fraction	0.056 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	1 of 81060 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5382	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RAM, NAG, GLA, GLC, NA, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/4807	0.65	0/6551

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4607	0	4342	29	0
2	A	4	0	0	0	0
3	A	8	0	12	0	0
4	A	72	0	59	0	0
5	A	691	0	0	4	0
All	All	5382	0	4413	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:303[A]:ASN:OD1	5:A:2258:HOH:O	1.76	1.02
1:A:637[B]:THR:HG22	5:A:2615:HOH:O	1.85	0.76
1:A:637[B]:THR:HG23	5:A:2601:HOH:O	1.97	0.64
1:A:409:TYR:HA	1:A:440:ASN:O	2.01	0.60
1:A:626:VAL:HG12	1:A:628[B]:VAL:CG1	2.31	0.60
1:A:381:ARG:HD2	1:A:414[B]:GLU:OE2	2.04	0.57
1:A:626:VAL:HG12	1:A:628[A]:VAL:HG23	1.86	0.57
1:A:241:PHE:O	1:A:283:ARG:HA	2.11	0.50
1:A:371:VAL:O	1:A:392:ALA:HA	2.14	0.48
1:A:288:PHE:HB3	1:A:291:SER:HB3	1.98	0.45
1:A:537:ASN:HA	1:A:569:ASN:O	2.16	0.45
1:A:428:GLY:HA2	1:A:480:ILE:HD11	1.98	0.44
1:A:626:VAL:HG12	1:A:628[B]:VAL:HG13	1.99	0.44
1:A:440:ASN:HA	1:A:481:GLN:O	2.18	0.44
1:A:528:PRO:HD3	5:A:2507:HOH:O	2.17	0.44
1:A:414[A]:GLU:HG2	1:A:445[A]:SER:OG	2.19	0.43
1:A:297:THR:HA	1:A:324:ARG:O	2.18	0.43
1:A:465:ILE:O	1:A:502:ILE:HA	2.17	0.43
1:A:398:ALA:HB3	1:A:432:THR:HG23	2.01	0.43
1:A:284:ASN:ND2	1:A:287:ALA:HB2	2.33	0.43
1:A:600:ILE:HB	1:A:628[A]:VAL:HG22	2.01	0.42
1:A:252:PRO:HB2	1:A:314:TRP:CE2	2.54	0.42
1:A:320:ASN:HA	1:A:349:TYR:O	2.18	0.42
1:A:341:SER:HA	1:A:369:CYS:O	2.19	0.41
1:A:288:PHE:HB3	1:A:291:SER:CB	2.50	0.41
1:A:621:GLN:O	1:A:707:ARG:HD3	2.20	0.41
1:A:599:LYS:HA	1:A:627:TYR:O	2.21	0.40
1:A:294:CYS:O	1:A:321:CYS:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	618/600 (103%)	603 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	512/492 (104%)	507 (99%)	5 (1%)	82	62

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	ARG
1	A	440	ASN
1	A	517	ASN
1	A	669[A]	LEU
1	A	669[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	RAM	A	1715	2,4	10,10,11	0.54	0	14,14,16	0.89	0
4	GLC	A	1716	4	11,11,12	0.77	0	14,15,17	1.19	2 (14%)
4	GLA	A	1717	4	11,11,12	1.56	3 (27%)	14,15,17	2.59	3 (21%)
4	NAG	A	1718	4	14,14,15	0.45	0	15,19,21	0.77	0
4	NAG	A	1719	2,4	15,15,15	0.36	0	17,21,21	1.08	2 (11%)
4	GLC	A	1720	2,4	11,11,12	0.81	1 (9%)	14,15,17	1.17	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RAM	A	1715	2,4	-	0/0/17/20	0/1/1/1
4	GLC	A	1716	4	-	0/2/19/22	0/1/1/1
4	GLA	A	1717	4	-	0/2/19/22	0/1/1/1
4	NAG	A	1718	4	-	0/6/23/26	0/1/1/1
4	NAG	A	1719	2,4	-	0/6/26/26	0/1/1/1
4	GLC	A	1720	2,4	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1717	GLA	C4-C5	-3.47	1.45	1.53
4	A	1717	GLA	O4-C4	-2.80	1.36	1.43
4	A	1720	GLC	O5-C1	-2.30	1.39	1.43
4	A	1717	GLA	O5-C1	-2.01	1.40	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1717	GLA	O4-C4-C5	-7.79	88.58	109.24
4	A	1716	GLC	C1-C2-C3	-2.73	106.31	109.54

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1717	GLA	C1-O5-C5	-2.34	109.28	112.25
4	A	1716	GLC	C1-O5-C5	-2.07	109.62	112.25
4	A	1719	NAG	C3-C2-N2	2.00	114.81	110.66
4	A	1719	NAG	O5-C5-C4	2.60	114.57	109.68
4	A	1717	GLA	C6-C5-C4	2.73	119.76	113.02
4	A	1720	GLC	C1-O5-C5	2.77	115.76	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	TRS	A	1714	-	7,7,7	0.67	0	9,9,9	1.61	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	1714	-	-	0/9/9/9	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	A	1714	TRS	C2-C-C1	-2.81	104.71	110.78
3	A	1714	TRS	O1-C1-C	2.25	115.72	111.18
3	A	1714	TRS	C1-C-N	2.25	112.19	108.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	597/600 (99%)	0.09	13 (2%) 65 68	7, 13, 23, 41	9 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	534	VAL	6.3
1	A	114	PHE	4.6
1	A	434	THR	4.3
1	A	529	GLY	4.1
1	A	533	PRO	3.9
1	A	433	ALA	3.7
1	A	530	THR	3.1
1	A	113	GLN	3.1
1	A	435	VAL	2.4
1	A	532	SER	2.3
1	A	489[A]	SER	2.2
1	A	258	LEU	2.1
1	A	669[A]	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	1719	15/15	0.94	0.10	0.86	11,13,17,18	0
4	RAM	A	1715	10/11	0.93	0.10	0.62	17,20,21,22	0
4	GLA	A	1717	11/12	0.94	0.09	0.06	9,13,17,17	0
4	NAG	A	1718	14/15	0.95	0.08	-0.11	12,14,25,26	0
4	GLC	A	1720	11/12	0.97	0.07	-0.45	11,13,14,14	0
4	GLC	A	1716	11/12	0.94	0.09	-	15,18,21,22	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NA	A	1713	1/1	0.97	0.26	7.10	26,26,26,26	0
2	NA	A	1710	1/1	0.97	0.15	2.60	16,16,16,16	0
3	TRS	A	1714	8/8	0.93	0.12	1.20	13,14,15,16	0
2	NA	A	1712	1/1	0.85	0.15	-	23,23,23,23	0
2	NA	A	1711	1/1	0.88	0.16	-	26,26,26,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.