



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:30 AM GMT

PDB ID : 2XAO  
Title : INOSITOL 1,3,4,5,6-PENTAKISPHOSPHATE 2-KINASE FROM A.  
THALIANA IN COMPLEX WITH IP5  
Authors : Gonzalez, B.; Banos-Sanz, J.I.; Villate, M.; Brearley, C.A.; Sanz-Aparicio, J.  
Deposited on : 2010-03-31  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 18%, green 68%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>3%</span> <span>68%</span> <span>20%</span> <span>• 10%</span> </div> </div>
1	B	451	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 18%, green 73%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>3%</span> <span>73%</span> <span>18%</span> <span>• 6%</span> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INOSITOL-PENTAKISPHOSPHATE 2-KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	0	0
			3224	2052	545	614	13			
1	B	424	Total	C	N	O	S	0	0	0
			3378	2148	572	645	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	SER	ALA	CONFLICT	UNP Q93YN9
A	90	GLN	LYS	CONFLICT	UNP Q93YN9
A	157	THR	SER	SEE REMARK 999	UNP Q93YN9
A	204	ILE	ASN	CONFLICT	UNP Q93YN9
A	224	ARG	SER	CONFLICT	UNP Q93YN9
A	321	CYS	SER	CONFLICT	UNP Q93YN9
A	325	ILE	LEU	CONFLICT	UNP Q93YN9
A	337	ARG	LYS	SEE REMARK 999	UNP Q93YN9
B	54	SER	ALA	CONFLICT	UNP Q93YN9
B	90	GLN	LYS	CONFLICT	UNP Q93YN9
B	157	THR	SER	SEE REMARK 999	UNP Q93YN9
B	204	ILE	ASN	CONFLICT	UNP Q93YN9
B	224	ARG	SER	CONFLICT	UNP Q93YN9
B	321	CYS	SER	CONFLICT	UNP Q93YN9
B	325	ILE	LEU	CONFLICT	UNP Q93YN9
B	337	ARG	LYS	CONFLICT	UNP Q93YN9

- Molecule 2 is MYO-INOSITOL-(1,3,4,5,6)-PENTAKISPHOSPHATE (three-letter code: 5MY) (formula: C<sub>6</sub>H<sub>17</sub>O<sub>21</sub>P<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			32	6	21	5		
2	B	1	Total	C	O	P	0	0
			32	6	21	5		


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: INOSITOL-PENTAKISPHOSPHATE 2-KINASE



#### • Molecule 1: INOSITOL-PENTAKISPHOSPHATE 2-KINASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.76Å 112.74Å 144.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.74 – 2.90 72.23 – 2.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (88.74-2.90) 100.0 (72.23-2.90)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.225 , 0.291 0.221 , 0.284	Depositor DCC
$R_{free}$ test set	1141 reflections (5.48%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.7	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 33.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 21980 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	6668	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 5MY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/3280	0.64	0/4416
1	B	0.53	2/3441 (0.1%)	0.65	0/4640
All	All	0.52	2/6721 (0.0%)	0.65	0/9056

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	153	SER	C-O	8.22	1.39	1.23
1	B	171	CYS	CB-SG	-5.56	1.72	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3224	0	3240	56	0
1	B	3378	0	3393	53	0
2	A	32	0	7	1	0
2	B	32	0	7	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
All	All	6668	0	6647	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:HIS:CD2	1:A:303:ARG:HH22	1.75	1.03
1:B:40:ARG:HD3	1:B:146:LEU:CD1	2.10	0.81
1:A:126:ARG:HH21	1:A:131:VAL:HA	1.49	0.77
1:A:67:HIS:O	1:A:70:ARG:HD2	1.88	0.73
1:A:15:TYR:HA	1:A:25:LEU:HD12	1.71	0.72
1:A:319:ILE:HD13	1:A:420:TYR:HB2	1.70	0.72
1:B:8:LYS:H	1:B:8:LYS:HD2	1.54	0.71
1:A:101:HIS:HD2	1:A:303:ARG:HH22	1.31	0.71
1:A:101:HIS:CD2	1:A:303:ARG:NH2	2.56	0.70
1:B:40:ARG:HD3	1:B:146:LEU:HD11	1.72	0.70
1:B:277:GLN:H	1:B:397:GLN:HE22	1.38	0.70
1:A:153:SER:HB3	1:A:402:LYS:NZ	2.07	0.69
1:B:348:LEU:HG	1:B:352:GLU:HG2	1.75	0.69
1:B:204:ILE:HG22	1:B:204:ILE:O	1.94	0.67
1:A:126:ARG:NH2	1:A:131:VAL:HA	2.09	0.67
1:B:14:ILE:HD12	1:B:15:TYR:H	1.60	0.66
1:B:222:LYS:HB3	1:B:222:LYS:NZ	2.10	0.66
1:B:101:HIS:HD2	1:B:303:ARG:NH2	1.94	0.65
1:B:226:LEU:HA	1:B:292:SER:OG	1.97	0.64
1:A:422:LEU:O	1:A:426:ILE:HG12	1.96	0.64
1:B:345:LEU:O	1:B:348:LEU:HB2	1.98	0.63
1:B:74:GLU:N	1:B:74:GLU:OE1	2.33	0.62
1:B:16:ARG:NH2	1:B:148:ASP:OD1	2.32	0.62
1:A:295:VAL:HG13	1:A:301:LEU:HD22	1.82	0.61
1:B:101:HIS:HD2	1:B:303:ARG:HH22	1.49	0.60
1:A:129:TRP:HB2	1:B:178:ARG:HG2	1.82	0.60
1:B:215:LEU:O	1:B:219:SER:HB3	2.01	0.60
1:A:47:ASN:OD1	1:B:188:THR:HG21	2.03	0.59
1:B:113:GLU:O	1:B:117:CYS:HB2	2.03	0.58
1:A:275:PHE:HD1	1:A:276:ILE:HD12	1.68	0.58
1:A:101:HIS:HD2	1:A:303:ARG:NH2	1.99	0.58
1:A:325:ILE:HD11	1:A:354:LEU:HA	1.86	0.57
1:A:97:LEU:HG	1:A:307:ILE:CD1	2.38	0.54
1:A:153:SER:HB3	1:A:402:LYS:HZ1	1.71	0.54
1:B:277:GLN:H	1:B:397:GLN:NE2	2.05	0.54
1:B:39:ILE:HD12	1:B:41:ILE:HG13	1.90	0.53
1:B:52:LYS:HB2	1:B:58:VAL:N	2.23	0.53

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:SER:HB3	1:A:402:LYS:HZ2	1.73	0.53
1:A:190:VAL:HB	1:A:195:MET:HE2	1.91	0.53
1:A:215:LEU:O	1:A:219:SER:HB3	2.09	0.52
2:A:500:5MY:O35	2:A:500:5MY:O34	2.28	0.52
1:A:18:GLU:OE1	1:A:126:ARG:HG2	2.11	0.51
1:B:393:LYS:N	1:B:394:PRO:HD2	2.26	0.51
1:B:39:ILE:CD1	1:B:41:ILE:HG13	2.41	0.50
1:A:168:LYS:HE3	1:A:170:LYS:HD3	1.92	0.50
1:A:94:ILE:HD13	1:A:102:VAL:HB	1.92	0.50
1:B:251:GLY:HA2	1:B:257:THR:HG21	1.94	0.50
1:B:198:LEU:HD12	1:B:430:TYR:HD1	1.75	0.50
1:A:295:VAL:HG12	1:A:296:TYR:N	2.28	0.49
1:A:319:ILE:HD13	1:A:420:TYR:CB	2.41	0.49
1:A:112:LYS:HG3	1:A:138:THR:O	2.13	0.49
1:B:222:LYS:HB3	1:B:222:LYS:HZ2	1.78	0.48
1:A:325:ILE:HG12	1:A:354:LEU:HD23	1.95	0.48
1:B:300:VAL:HG11	1:B:403:VAL:HG21	1.94	0.48
1:B:345:LEU:HA	1:B:348:LEU:HD22	1.95	0.48
1:A:199:LEU:HD21	1:A:425:LYS:HE3	1.95	0.48
1:B:172:GLY:HA3	1:B:218:PHE:CD2	2.49	0.47
1:B:25:LEU:HB2	1:B:39:ILE:CG2	2.45	0.47
1:A:197:GLN:HB3	1:A:209:GLU:HG3	1.97	0.47
1:B:341:ALA:HA	1:B:344:SER:HB2	1.97	0.47
1:A:72:ASN:HB3	1:A:74:GLU:OE2	2.15	0.47
1:B:39:ILE:HD11	1:B:143:ALA:HB1	1.97	0.47
1:A:119:ASP:HB2	1:A:136:VAL:CG2	2.45	0.46
1:B:40:ARG:CD	1:B:146:LEU:CD1	2.88	0.46
1:B:184:ASN:O	1:B:187:LYS:HG2	2.15	0.45
1:B:226:LEU:HA	1:B:292:SER:HG	1.80	0.45
1:A:200:LYS:HB3	1:A:206:ILE:HG12	1.98	0.45
1:A:170:LYS:HA	1:A:367:LYS:O	2.17	0.45
1:B:337:ARG:HA	1:B:338:PRO:HD3	1.65	0.45
1:B:325:ILE:HD12	1:B:354:LEU:HD23	2.00	0.44
1:B:308:GLN:NE2	1:B:367:LYS:HA	2.32	0.44
1:A:46:ARG:NH1	1:B:182:LYS:HG2	2.32	0.44
1:B:341:ALA:O	1:B:345:LEU:N	2.49	0.44
1:B:351:ASP:HA	1:B:354:LEU:HD12	1.99	0.44
1:B:186:LEU:HA	1:B:186:LEU:HD12	1.90	0.44
1:A:216:ASP:OD1	1:A:224:ARG:NE	2.45	0.44
1:A:186:LEU:HD12	1:A:186:LEU:HA	1.83	0.44
1:A:300:VAL:HG11	1:A:403:VAL:HG11	2.00	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:PRO:HG2	1:A:83:VAL:HG21	1.99	0.43
1:A:163:ILE:HG13	1:A:275:PHE:CD1	2.54	0.43
1:B:165:VAL:HG22	1:B:242:VAL:HG22	2.01	0.43
1:B:204:ILE:O	1:B:204:ILE:CG2	2.65	0.43
1:B:97:LEU:HD13	1:B:307:ILE:HG13	2.01	0.43
1:A:172:GLY:HA3	1:A:218:PHE:CD2	2.53	0.43
1:A:166:GLU:HB3	1:A:241:ARG:HB2	2.00	0.42
1:A:93:ILE:HG23	1:A:97:LEU:HD12	2.00	0.42
1:A:38:VAL:HG13	1:A:148:ASP:HB2	2.00	0.42
1:B:16:ARG:HB3	1:B:24:VAL:O	2.20	0.42
1:B:198:LEU:HD12	1:B:430:TYR:CD1	2.52	0.42
1:B:212:TYR:CE2	1:B:214:PRO:HG3	2.55	0.42
1:A:322:TYR:HE2	1:A:417:GLU:HG3	1.84	0.42
1:A:73:ASN:O	1:A:76:ILE:HG13	2.19	0.42
1:A:212:TYR:CE2	1:A:214:PRO:HG3	2.54	0.42
1:B:74:GLU:O	1:B:87:ARG:NH1	2.51	0.41
1:A:18:GLU:OE2	1:A:126:ARG:NH1	2.53	0.41
1:A:185:MET:O	1:A:188:THR:HB	2.20	0.41
1:A:342:GLU:N	1:A:342:GLU:OE1	2.54	0.41
1:A:295:VAL:HG23	1:A:373:ILE:HD11	2.03	0.41
1:A:251:GLY:HA2	1:A:257:THR:HG21	2.03	0.41
1:A:322:TYR:CE2	1:A:417:GLU:HG3	2.56	0.41
1:B:25:LEU:HB2	1:B:39:ILE:HG22	2.01	0.41
1:B:101:HIS:CD2	1:B:303:ARG:HH22	2.34	0.41
1:A:97:LEU:HD22	1:A:405:PHE:CE1	2.56	0.41
1:B:313:LEU:O	1:B:314:ASP:HB3	2.21	0.41
1:B:36:GLY:HA2	1:B:151:LEU:HD11	2.03	0.41
1:A:287:PHE:O	1:A:290:LEU:HB3	2.21	0.41
1:A:90:GLN:HA	1:A:94:ILE:HG13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/451 (87%)	374 (95%)	16 (4%)	2 (0%)	34	71
1	B	418/451 (93%)	396 (95%)	21 (5%)	1 (0%)	52	84
All	All	810/902 (90%)	770 (95%)	37 (5%)	3 (0%)	39	74

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	49	LYS
1	A	70	ARG
1	A	177	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/399 (91%)	334 (92%)	28 (8%)	16	42
1	B	378/399 (95%)	336 (89%)	42 (11%)	8	22
All	All	740/798 (93%)	670 (90%)	70 (10%)	11	31

All (70) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	16	ARG
1	A	25	LEU
1	A	40	ARG
1	A	72	ASN
1	A	74	GLU
1	A	96	LEU
1	A	164	SER
1	A	174	LEU
1	A	176	THR
1	A	186	LEU
1	A	188	THR
1	A	206	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	207	SER
1	A	223	GLU
1	A	234	SER
1	A	238	ASN
1	A	244	LEU
1	A	247	SER
1	A	276	ILE
1	A	295	VAL
1	A	301	LEU
1	A	310	LEU
1	A	343	LEU
1	A	348	LEU
1	A	350	LEU
1	A	351	ASP
1	A	372	MET
1	B	4	ILE
1	B	5	LEU
1	B	8	LYS
1	B	16	ARG
1	B	39	ILE
1	B	52	LYS
1	B	75	LEU
1	B	83	VAL
1	B	97	LEU
1	B	103	ASP
1	B	117	CYS
1	B	120	LYS
1	B	125	GLN
1	B	145	ILE
1	B	151	LEU
1	B	189	SER
1	B	196	HIS
1	B	198	LEU
1	B	202	GLU
1	B	207	SER
1	B	208	GLU
1	B	234	SER
1	B	237	GLN
1	B	253	SER
1	B	285	GLU
1	B	290	LEU
1	B	292	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	300	VAL
1	B	301	LEU
1	B	334	LYS
1	B	337	ARG
1	B	344	SER
1	B	348	LEU
1	B	352	GLU
1	B	355	LYS
1	B	371	ILE
1	B	372	MET
1	B	377	SER
1	B	379	ASN
1	B	398	THR
1	B	418	SER
1	B	437	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	A	47	ASN
1	A	91	ASN
1	A	101	HIS
1	A	238	ASN
1	B	22	ASN
1	B	147	ASN
1	B	397	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	5MY	A	500	-	32,32,32	0.58	0	43,53,53	1.21	5 (11%)
2	5MY	B	500	-	32,32,32	0.63	0	43,53,53	1.20	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5MY	A	500	-	-	0/25/49/49	0/1/1/1
2	5MY	B	500	-	-	0/25/49/49	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	5MY	O15-C5-C6	-2.51	102.64	108.47
2	A	500	5MY	O15-C5-C6	-2.32	103.09	108.47
2	A	500	5MY	O43-P3-O23	2.19	115.73	107.38
2	A	500	5MY	O45-P5-O25	2.42	116.60	107.38
2	A	500	5MY	O24-P4-O44	2.44	116.67	107.38
2	A	500	5MY	O21-P1-O31	2.49	116.85	107.38
2	B	500	5MY	O24-P4-O44	2.72	117.74	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	5MY	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/451 (89%)	-0.08	3 (0%) 89 88	24, 38, 57, 69	0
1	B	424/451 (94%)	0.02	15 (3%) 48 40	24, 38, 61, 81	0
All	All	828/902 (91%)	-0.03	18 (2%) 65 60	24, 38, 58, 81	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	47	ASN	6.1
1	B	381	TRP	4.4
1	B	254	GLY	4.2
1	B	48	ASP	3.8
1	B	50	ALA	3.4
1	B	338	PRO	2.8
1	B	340	GLU	2.8
1	B	67	HIS	2.8
1	B	253	SER	2.6
1	A	48	ASP	2.6
1	B	280	ASP	2.5
1	A	343	LEU	2.4
1	B	124	LYS	2.3
1	B	256	SER	2.2
1	B	49	LYS	2.2
1	B	153	SER	2.2
1	A	47	ASN	2.1
1	B	339	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	5MY	B	500	32/32	0.98	0.09	-1.63	17,21,26,27	0
2	5MY	A	500	32/32	0.98	0.08	-2.27	13,21,26,28	0
3	ZN	A	700	1/1	0.99	0.05	-2.49	40,40,40,40	0
3	ZN	B	700	1/1	0.97	0.07	-	45,45,45,45	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.