



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:18 PM GMT

PDB ID : 4XAQ  
Title : mGluR2 ECD and mGluR3 ECD with ligands  
Authors : Clawson, D.K.  
Deposited on : 2014-12-15  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	
1	B	503	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7532 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metabotropic glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	0	0
			3475	2206	623	633	13			
1	B	442	Total	C	N	O	S	0	4	0
			3491	2215	624	639	13			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP Q14416
A	0	ALA	-	expression tag	UNP Q14416
A	1	LEU	-	expression tag	UNP Q14416
A	234	SER	CYS	conflict	UNP Q14416
A	302	GLU	SER	conflict	UNP Q14416
A	494	GLU	-	expression tag	UNP Q14416
A	495	GLY	-	expression tag	UNP Q14416
A	496	HIS	-	expression tag	UNP Q14416
A	497	HIS	-	expression tag	UNP Q14416
A	498	HIS	-	expression tag	UNP Q14416
A	499	HIS	-	expression tag	UNP Q14416
A	500	HIS	-	expression tag	UNP Q14416
A	501	HIS	-	expression tag	UNP Q14416
B	-1	MET	-	initiating methionine	UNP Q14416
B	0	ALA	-	expression tag	UNP Q14416
B	1	LEU	-	expression tag	UNP Q14416
B	234	SER	CYS	conflict	UNP Q14416
B	302	GLU	SER	conflict	UNP Q14416
B	494	GLU	-	expression tag	UNP Q14416
B	495	GLY	-	expression tag	UNP Q14416
B	496	HIS	-	expression tag	UNP Q14416
B	497	HIS	-	expression tag	UNP Q14416
B	498	HIS	-	expression tag	UNP Q14416
B	499	HIS	-	expression tag	UNP Q14416
B	500	HIS	-	expression tag	UNP Q14416

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	501	HIS	-	expression tag	UNP Q14416

- Molecule 2 is (1S,2S,5R,6S)-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid (three-letter code: 40F) (formula: C<sub>8</sub>H<sub>11</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			13	8	1	4		
2	B	1	Total	C	N	O	0	0
			13	8	1	4		

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	A	2	Total	Cl	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	226	Total	O	0	0
			226	226		
5	B	301	Total	O	0	0
			301	301		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metabotropic glutamate receptor 2



- Molecule 1: Metabotropic glutamate receptor 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.64Å 159.65Å 93.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.52 – 2.21 27.44 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.6 (25.52-2.21) 99.6 (27.44-2.21)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 2.22Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, $R_{free}$	0.186 , 0.225 0.193 , 0.229	Depositor DCC
$R_{free}$ test set	1898 reflections (3.16%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.1	Xtriage
Anisotropy	0.612	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 57.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 61915 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7532	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 40F, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/3559	0.64	0/4824
1	B	0.54	0/3573	0.68	1/4842 (0.0%)
All	All	0.52	0/7132	0.66	1/9666 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	444	ASP	CB-CG-OD1	5.54	123.29	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3475	0	3362	20	0
1	B	3491	0	3368	26	0
2	A	13	0	8	0	0
2	B	13	0	8	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	2	0	0	1	0
4	B	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	226	0	0	1	0
5	B	301	0	0	2	0
All	All	7532	0	6746	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:399:LEU:HD21	1:B:417:ARG:HH22	1.49	0.75
1:B:396:HIS:HB2	1:B:410:MET:HE1	1.77	0.67
1:B:95:ASP:HB2	1:B:150:GLN:HG3	1.79	0.65
1:A:221:ILE:HG12	1:A:269:PHE:HB2	1.79	0.64
1:A:210:VAL:HB	1:A:268:LEU:HD22	1.83	0.59
1:B:199:LEU:HD13	1:B:207:VAL:HG11	1.85	0.58
1:A:236:ALA:HB1	1:A:260:LYS:HG2	1.85	0.58
1:B:56:HIS:HD2	5:B:828:HOH:O	1.87	0.58
1:A:24:LYS:HD3	1:A:343:PRO:HB2	1.87	0.57
1:B:210:VAL:HG11	1:B:278:LEU:HD11	1.89	0.55
1:A:199:LEU:HD13	1:A:207:VAL:HG11	1.89	0.54
1:B:370:VAL:HB	1:B:371:PRO:CD	2.36	0.54
1:A:478:LEU:HD11	1:A:480:LEU:HD23	1.91	0.53
1:B:210:VAL:HB	1:B:268:LEU:HD22	1.92	0.51
1:A:26:LEU:HD13	1:A:62:LEU:HD11	1.92	0.51
1:B:336:PRO:HG3	1:B:349:TRP:CD2	2.47	0.49
1:A:32:LEU:HG	1:A:406:LEU:HD11	1.94	0.49
1:B:56:HIS:CD2	5:B:828:HOH:O	2.64	0.48
1:A:271:ARG:HB3	4:A:604:CL:CL	2.50	0.48
1:B:225:GLU:HG2	1:B:235:VAL:HG21	1.95	0.47
1:A:386:TYR:CG	1:A:431:PRO:HG3	2.50	0.47
5:A:818:HOH:O	1:B:177:ARG:HD2	2.14	0.46
1:B:171:LYS:HE3	1:B:219:THR:HG21	1.98	0.46
1:B:199:LEU:HD22	1:B:204:TRP:HE3	1.81	0.45
1:B:396:HIS:CB	1:B:410:MET:HE1	2.43	0.45
1:A:177:ARG:HB3	1:A:177:ARG:HH11	1.80	0.45
1:B:370:VAL:HB	1:B:371:PRO:HD2	1.98	0.45
1:B:149:ILE:HG13	1:B:172:LEU:HD21	1.98	0.45
1:B:452:ARG:HH11	1:B:475:ALA:HB1	1.82	0.45
1:B:212:SER:HA	1:B:241:VAL:HG23	1.99	0.44
1:B:163:ILE:HA	1:B:182:ALA:O	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:ARG:CZ	1:B:177:ARG:HD3	2.48	0.43
1:A:210:VAL:HG22	1:A:239:GLU:HB2	2.02	0.42
1:B:228:ALA:HB1	1:B:233:ILE:HB	2.02	0.42
1:B:30:GLY:HA2	1:B:83:ARG:HG2	2.01	0.41
1:A:205:THR:HG23	1:A:232:ASN:O	2.20	0.41
1:A:95:ASP:CG	1:A:243:ARG:HH22	2.24	0.41
1:B:189:PHE:CE2	1:B:193:LYS:HE2	2.56	0.41
1:A:59:ILE:HG13	1:A:348:PHE:HB2	2.02	0.41
1:A:163:ILE:HA	1:A:182:ALA:O	2.20	0.41
1:A:228:ALA:HB1	1:A:233:ILE:HB	2.03	0.41
1:A:314:ALA:O	1:A:457:THR:HA	2.21	0.41
1:A:159:GLN:O	1:A:415:GLY:HA3	2.21	0.41
1:B:221:ILE:HD13	1:B:269:PHE:HB2	2.03	0.41
1:B:159:GLN:O	1:B:415:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/503 (87%)	420 (96%)	18 (4%)	1 (0%)	52	59
1	B	440/503 (88%)	424 (96%)	14 (3%)	2 (0%)	34	34
All	All	879/1006 (87%)	844 (96%)	32 (4%)	3 (0%)	46	50

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	135	ALA
1	B	109	SER
1	A	451	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/402 (88%)	345 (98%)	8 (2%)	58	70
1	B	355/402 (88%)	347 (98%)	8 (2%)	58	70
All	All	708/804 (88%)	692 (98%)	16 (2%)	61	70

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	A	71	ARG
1	A	177	ARG
1	A	269	PHE
1	A	318	GLU
1	A	335	ASP
1	A	417	ARG
1	A	442	ARG
1	B	215	ASP
1	B	269	PHE
1	B	278	LEU
1	B	318[A]	GLU
1	B	318[B]	GLU
1	B	329	SER
1	B	359	GLN
1	B	433	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	40F	A	601	-	6,14,14	1.12	0	2,23,23	0.45	0
3	SO4	A	602	-	4,4,4	0.14	0	6,6,6	0.23	0
2	40F	B	601	-	6,14,14	1.52	1 (16%)	2,23,23	0.37	0
3	SO4	B	602	-	4,4,4	0.32	0	6,6,6	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	40F	A	601	-	-	0/0/31/31	0/1/2/2
3	SO4	A	602	-	-	0/0/0/0	0/0/0/0
2	40F	B	601	-	-	0/0/31/31	0/1/2/2
3	SO4	B	602	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	40F	C7-C1	3.25	1.58	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	443/503 (88%)	0.42	57 (12%) 5 4	23, 45, 76, 118	0
1	B	442/503 (87%)	0.01	29 (6%) 22 21	22, 38, 65, 102	0
All	All	885/1006 (87%)	0.22	86 (9%) 10 9	22, 42, 73, 118	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	22	ALA	7.9
1	A	358	ARG	6.9
1	A	462	GLY	6.2
1	A	438	HIS	6.2
1	A	461	ALA	5.5
1	A	359	GLN	5.0
1	A	360	ARG	4.9
1	A	337	TRP	4.8
1	B	262	SER	4.7
1	A	140	ILE	4.6
1	B	402	ASN	4.6
1	A	464	GLY	4.6
1	A	437	THR	4.2
1	A	364	ALA	4.0
1	A	465	ARG	4.0
1	A	435	ALA	3.9
1	A	47	ALA	3.9
1	B	434	PRO	3.8
1	A	463	SER	3.8
1	A	46	PRO	3.6
1	A	460	ARG	3.5
1	B	261	PRO	3.5
1	B	439	ASN	3.5
1	A	434	PRO	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	262	SER	3.4
1	B	292	VAL	3.3
1	A	362	CYS	3.3
1	A	361	ASP	3.2
1	A	363	ALA	3.2
1	B	140	ILE	3.2
1	A	369	ALA	3.2
1	A	385	VAL	3.2
1	A	325	SER	3.1
1	A	232	ASN	3.0
1	A	402	ASN	2.9
1	B	139	VAL	2.9
1	A	74	ARG	2.8
1	A	459	LEU	2.8
1	B	268	LEU	2.8
1	A	77	HIS	2.8
1	A	467	ARG	2.8
1	B	293	ALA	2.8
1	B	489	SER	2.7
1	B	435	ALA	2.7
1	A	134	THR	2.7
1	B	47	ALA	2.7
1	A	357	PHE	2.7
1	A	371	PRO	2.6
1	A	353	PHE	2.6
1	A	356	SER	2.6
1	B	408	ASP	2.6
1	A	108	ALA	2.5
1	A	139	VAL	2.5
1	A	206	TYR	2.5
1	A	261	PRO	2.5
1	B	229	ARG	2.4
1	A	436	ASP	2.4
1	A	486	PRO	2.4
1	A	466	TYR	2.4
1	A	76	PRO	2.4
1	B	388	MET	2.4
1	A	292	VAL	2.4
1	B	315	ILE	2.4
1	B	48	GLU	2.3
1	A	35	GLY	2.3
1	A	349	TRP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	325	SER	2.3
1	A	388	MET	2.3
1	B	381	VAL	2.3
1	B	385	VAL	2.3
1	B	438	HIS	2.3
1	A	37	LEU	2.3
1	A	315	ILE	2.2
1	A	473	TYR	2.2
1	B	232	ASN	2.2
1	B	258	LEU	2.2
1	B	110	LEU	2.2
1	A	293	ALA	2.2
1	B	384	ALA	2.2
1	B	206	TYR	2.2
1	B	316	THR	2.2
1	A	165	TYR	2.1
1	A	483	SER	2.1
1	A	62	LEU	2.1
1	A	368	ARG	2.0
1	B	108	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	40F	B	601	13/13	0.98	0.13	-0.18	20,23,26,27	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	40F	A	601	13/13	0.98	0.12	-0.39	25,29,33,34	0
4	CL	A	603	1/1	0.98	0.14	-0.71	35,35,35,35	0
4	CL	A	604	1/1	0.97	0.08	-0.89	68,68,68,68	0
4	CL	B	603	1/1	0.99	0.10	-1.33	28,28,28,28	0
3	SO4	B	602	5/5	0.95	0.10	-	60,61,65,69	0
3	SO4	A	602	5/5	0.97	0.08	-	67,69,71,73	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.