



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 11:29 PM GMT

PDB ID : 1XCB  
Title : X-ray Structure of a Rex-Family Repressor/NADH Complex from *Thermus Aquaticus*  
Authors : Sickmier, E.A.; Brekasis, D.; Paranawithana, S.; Bonanno, J.B.; Burley, S.K.; Paget, M.S.; Kielkopf, C.L.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2004-09-01  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>2%</div> <div>47% 39% 11% .</div> </div>
1	B	211	<div> <div>53% 35% 5% . 7%</div> </div>
1	C	211	<div> <div>2%</div> <div>52% 38% 6% .</div> </div>
1	D	211	<div> <div>%</div> <div>46% 41% 6% 7%</div> </div>
1	E	211	<div> <div>%</div> <div>53% 40% . . .</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	211	
1	G	211	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CA	A	212	-	-	-	X
2	CA	B	212	-	-	-	X
2	CA	D	212	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Redox-sensing transcriptional repressor rex.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	205	Total	C	N	O	S	Se	0	0	0
			1591	1025	283	281	1	1			
1	B	197	Total	C	N	O	S	Se	0	0	0
			1530	986	271	271	1	1			
1	C	203	Total	C	N	O	S	Se	0	0	0
			1566	1008	277	279	1	1			
1	D	197	Total	C	N	O	S	Se	0	0	0
			1531	989	268	272	1	1			
1	E	203	Total	C	N	O	S	Se	0	0	0
			1565	1007	276	279	1	2			
1	F	200	Total	C	N	O	S	Se	0	0	0
			1558	1005	274	277	1	1			
1	G	192	Total	C	N	O	S	Se	0	2	0
			1505	973	263	267	1	1			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
A	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
A	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
A	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
B	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
B	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
B	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
C	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
C	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
C	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
D	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
D	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
E	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
E	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
E	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
E	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
F	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
F	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
F	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
F	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
G	1	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
G	88	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
G	209	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5
G	210	MSE	MET	MODIFIED RESIDUE	UNP Q9X2V5

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	D	2	Total Ca 2 2	0	0
2	C	1	Total Ca 1 1	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	F	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	G	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Redox-sensing transcriptional repressor rex



- Molecule 1: Redox-sensing transcriptional repressor rex




- Molecule 1: Redox-sensing transcriptional repressor rex







- Molecule 1: Redox-sensing transcriptional repressor rex




- Molecule 1: Redox-sensing transcriptional repressor rex




- Molecule 1: Redox-sensing transcriptional repressor rex



- Molecule 1: Redox-sensing transcriptional repressor rex







## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	189.24Å 88.97Å 112.91Å 90.00° 106.09° 90.00°	Depositor
Resolution (Å)	20.00 – 2.90 19.98 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.90) 98.3 (19.98-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.87 (at 2.88Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.228 , 0.276 0.232 , 0.280	Depositor DCC
$R_{free}$ test set	2737 reflections (6.95%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.8	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 78045 reflections	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	11159	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.37	0/1621	0.67	0/2194
1	B	0.38	0/1558	0.65	0/2107
1	C	0.35	0/1594	0.64	0/2157
1	D	0.36	0/1560	0.64	0/2111
1	E	0.42	0/1593	0.71	0/2156
1	F	0.35	0/1587	0.63	0/2148
1	G	0.65	5/1532 (0.3%)	0.77	7/2073 (0.3%)
All	All	0.42	5/11045 (0.0%)	0.67	7/14946 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	188	ASP	C-N	-8.03	1.15	1.34
1	G	190[A]	LEU	C-N	-7.87	1.16	1.34
1	G	190[B]	LEU	C-N	-7.87	1.16	1.34
1	G	191	ALA	N-CA	-7.66	1.31	1.46
1	G	188	ASP	CA-CB	-5.59	1.41	1.53

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	188	ASP	CA-C-N	-10.65	93.77	117.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	191	ALA	CB-CA-C	7.24	120.95	110.10
1	G	190[A]	LEU	CA-C-O	5.41	131.46	120.10
1	G	190[B]	LEU	CA-C-O	5.41	131.46	120.10
1	G	169	LEU	N-CA-C	-5.16	97.07	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	188	ASP	Mainchain
1	G	190[A]	LEU	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1591	0	1649	121	0
1	B	1530	0	1583	98	0
1	C	1566	0	1626	102	0
1	D	1531	0	1579	109	0
1	E	1565	0	1625	89	0
1	F	1558	0	1606	118	0
1	G	1505	0	1557	98	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	44	0	26	4	0
3	B	44	0	26	3	0
3	C	44	0	26	4	0
3	D	44	0	26	1	0
3	E	44	0	26	2	0
3	F	44	0	26	3	0
3	G	44	0	26	5	0
All	All	11159	0	11407	679	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 679 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:132:LEU:H	1:E:132:LEU:HD13	1.15	1.06
1:E:154:GLN:NE2	1:E:177:GLU:HB3	1.71	1.06
1:G:122:VAL:HG12	1:G:123:ARG:H	1.20	1.06
1:C:132:LEU:H	1:C:132:LEU:HD12	1.19	1.01
1:A:60:VAL:HG23	1:A:61:GLY:H	1.26	1.01

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/211 (96%)	170 (84%)	22 (11%)	11 (5%)	2	7
1	B	193/211 (92%)	165 (86%)	23 (12%)	5 (3%)	7	26
1	C	201/211 (95%)	161 (80%)	33 (16%)	7 (4%)	4	18
1	D	193/211 (92%)	169 (88%)	19 (10%)	5 (3%)	7	26
1	E	201/211 (95%)	177 (88%)	16 (8%)	8 (4%)	4	15
1	F	196/211 (93%)	156 (80%)	27 (14%)	13 (7%)	1	4
1	G	190/211 (90%)	153 (80%)	31 (16%)	6 (3%)	5	20
All	All	1377/1477 (93%)	1151 (84%)	171 (12%)	55 (4%)	4	15

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	VAL
1	A	118	VAL
1	A	123	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	103	GLU
1	B	153	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/167 (99%)	147 (89%)	19 (11%)	7	21
1	B	160/167 (96%)	146 (91%)	14 (9%)	12	35
1	C	164/167 (98%)	152 (93%)	12 (7%)	17	45
1	D	160/167 (96%)	144 (90%)	16 (10%)	9	28
1	E	164/167 (98%)	155 (94%)	9 (6%)	27	61
1	F	163/167 (98%)	150 (92%)	13 (8%)	15	40
1	G	158/167 (95%)	149 (94%)	9 (6%)	25	59
All	All	1135/1169 (97%)	1043 (92%)	92 (8%)	15	39

5 of 92 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	150	ARG
1	D	131	ASP
1	G	62	TYR
1	C	168	ILE
1	D	67	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	129	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAD	A	213	-	38,48,48	2.44	9 (23%)	47,73,73	2.65	13 (27%)
3	NAD	B	213	-	38,48,48	2.28	8 (21%)	47,73,73	2.42	9 (19%)
3	NAD	C	213	-	38,48,48	2.41	9 (23%)	47,73,73	2.57	15 (31%)
3	NAD	D	214	-	38,48,48	2.29	10 (26%)	47,73,73	2.72	13 (27%)
3	NAD	E	212	-	38,48,48	2.31	9 (23%)	47,73,73	2.66	14 (29%)
3	NAD	F	212	-	38,48,48	2.33	9 (23%)	47,73,73	2.56	12 (25%)
3	NAD	G	212	-	38,48,48	2.29	8 (21%)	47,73,73	2.56	15 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	213	-	-	0/22/62/62	0/5/5/5
3	NAD	B	213	-	-	0/22/62/62	0/5/5/5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	C	213	-	-	0/22/62/62	0/5/5/5
3	NAD	D	214	-	-	0/22/62/62	0/5/5/5
3	NAD	E	212	-	-	0/22/62/62	0/5/5/5
3	NAD	F	212	-	-	0/22/62/62	0/5/5/5
3	NAD	G	212	-	-	0/22/62/62	0/5/5/5

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	212	NAD	C2N-C3N	-3.21	1.34	1.39
3	F	212	NAD	C2N-C3N	-3.05	1.34	1.39
3	B	213	NAD	C2N-C3N	-3.02	1.34	1.39
3	D	214	NAD	C2N-C3N	-2.54	1.35	1.39
3	C	213	NAD	C2N-C3N	-2.34	1.35	1.39

The worst 5 of 91 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	213	NAD	C5N-C4N-C3N	-9.97	107.80	120.33
3	E	212	NAD	C5N-C4N-C3N	-9.94	107.83	120.33
3	B	213	NAD	N3A-C2A-N1A	-9.68	121.48	128.89
3	D	214	NAD	C5N-C4N-C3N	-9.65	108.20	120.33
3	D	214	NAD	N3A-C2A-N1A	-9.34	121.74	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	213	NAD	4	0
3	B	213	NAD	3	0
3	C	213	NAD	4	0
3	D	214	NAD	1	0
3	E	212	NAD	2	0
3	F	212	NAD	3	0
3	G	212	NAD	5	0



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	204/211 (96%)	-0.06	4 (1%) 68 64	11, 31, 54, 66	0
1	B	196/211 (92%)	-0.29	1 (0%) 91 90	11, 25, 47, 59	0
1	C	202/211 (95%)	-0.03	4 (1%) 68 64	13, 32, 71, 77	0
1	D	196/211 (92%)	-0.16	3 (1%) 76 74	10, 29, 64, 74	0
1	E	201/211 (95%)	-0.34	3 (1%) 76 74	10, 22, 42, 64	0
1	F	199/211 (94%)	0.11	4 (2%) 68 64	10, 42, 71, 80	0
1	G	191/211 (90%)	0.09	7 (3%) 45 38	11, 29, 91, 98	0
All	All	1389/1477 (94%)	-0.10	26 (1%) 70 66	10, 30, 69, 98	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	4	PRO	6.5
1	C	57	THR	6.2
1	G	26	VAL	4.3
1	G	24	GLN	3.9
1	G	25	GLY	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	A	212	1/1	0.88	0.40	5.75	58,58,58,58	0
2	CA	B	212	1/1	0.93	0.25	2.83	45,45,45,45	0
2	CA	D	212	1/1	0.91	0.24	2.79	47,47,47,47	0
3	NAD	D	214	44/44	0.96	0.14	-0.32	10,20,25,31	0
3	NAD	G	212	44/44	0.95	0.15	-0.32	16,27,37,39	0
3	NAD	F	212	44/44	0.94	0.16	-0.44	20,28,35,37	0
3	NAD	E	212	44/44	0.96	0.14	-0.48	10,15,23,28	0
3	NAD	C	213	44/44	0.95	0.15	-0.61	21,28,40,41	0
3	NAD	B	213	44/44	0.95	0.13	-0.94	14,22,30,34	0
3	NAD	A	213	44/44	0.93	0.16	-0.96	20,27,32,37	0
2	CA	D	213	1/1	0.90	0.24	-	57,57,57,57	0
2	CA	C	212	1/1	0.89	0.28	-	64,64,64,64	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.