



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:28 PM GMT

PDB ID : 1XD9
Title : Crystal Structure of the Nitrogenase Fe protein Asp39Asn with MgADP bound
Authors : Jang, S.B.; Jeong, M.S.; Seefeldt, L.C.; Peters, J.W.
Deposited on : 2004-09-05
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div></div> <div>69%</div> <div>26%</div> <div>..</div> </div>
1	B	289	<div> <div>3%</div> <div>71%</div> <div>24%</div> <div>..</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrogenase iron protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total	C	N	O	S	0	0	0
			2187	1364	370	432	21			
1	B	289	Total	C	N	O	S	0	0	0
			2187	1364	370	432	21			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASN	ASP	ENGINEERED	UNP P00459
B	39	ASN	ASP	ENGINEERED	UNP P00459

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	Fe	S		0	0
			8	4	4			

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitrogenase iron protein 1



- Molecule 1: Nitrogenase iron protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.05Å 92.21Å 61.38Å 90.00° 101.23° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 19.71 – 2.70	Depositor EDS
% Data completeness (in resolution range)	85.2 (20.00-2.80) 89.7 (19.71-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.71Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.229 , 0.299 0.244 , 0.301	Depositor DCC
R_{free} test set	664 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	62.6	Xtriage
Anisotropy	0.407	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 44.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 15393 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4438	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/2211	1.72	21/2977 (0.7%)
1	B	0.70	0/2211	1.75	21/2977 (0.7%)
All	All	0.70	0/4422	1.73	42/5954 (0.7%)

There are no bond length outliers.

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	245	ARG	NE-CZ-NH1	23.02	131.81	120.30
1	B	100	ARG	NE-CZ-NH1	-22.02	109.29	120.30
1	B	245	ARG	NE-CZ-NH2	21.55	131.08	120.30
1	B	100	ARG	NE-CZ-NH2	21.54	131.07	120.30
1	A	191	ARG	NE-CZ-NH2	21.32	130.96	120.30
1	A	191	ARG	NE-CZ-NH1	-21.00	109.80	120.30
1	B	191	ARG	NE-CZ-NH2	-20.86	109.87	120.30
1	A	100	ARG	NE-CZ-NH2	-20.29	110.15	120.30
1	B	191	ARG	NE-CZ-NH1	20.19	130.40	120.30
1	B	245	ARG	NE-CZ-NH1	-19.35	110.62	120.30
1	A	100	ARG	NE-CZ-NH1	19.01	129.80	120.30
1	A	245	ARG	NE-CZ-NH2	-18.50	111.05	120.30
1	A	245	ARG	CD-NE-CZ	18.13	148.98	123.60
1	B	245	ARG	CD-NE-CZ	17.24	147.73	123.60
1	B	191	ARG	CD-NE-CZ	16.92	147.29	123.60
1	B	100	ARG	CD-NE-CZ	16.39	146.55	123.60
1	A	191	ARG	CD-NE-CZ	15.31	145.04	123.60
1	A	100	ARG	CD-NE-CZ	15.02	144.62	123.60
1	A	187	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	A	187	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	A	213	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	118	ASP	CA-C-N	-6.48	102.94	117.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	95	VAL	CA-CB-CG2	-6.34	101.39	110.90
1	B	213	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	B	187	ARG	NE-CZ-NH1	6.31	123.45	120.30
1	B	118	ASP	O-C-N	6.29	132.76	122.70
1	B	118	ASP	CA-C-N	-6.25	103.45	117.20
1	B	187	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	B	46	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	A	118	ASP	O-C-N	5.85	132.07	122.70
1	A	95	VAL	CA-CB-CG2	-5.77	102.25	110.90
1	B	22	LEU	CA-CB-CG	5.54	128.05	115.30
1	A	124	TYR	CB-CG-CD2	-5.39	117.77	121.00
1	A	22	LEU	CA-CB-CG	5.37	127.65	115.30
1	B	285	THR	CA-CB-CG2	5.29	119.81	112.40
1	A	190	ASP	N-CA-C	5.28	125.26	111.00
1	A	118	ASP	CB-CG-OD1	5.28	123.05	118.30
1	B	190	ASP	N-CA-C	5.23	125.13	111.00
1	B	100	ARG	CB-CG-CD	-5.23	98.01	111.60
1	A	285	THR	CA-CB-CG2	5.18	119.65	112.40
1	A	46	ARG	NE-CZ-NH1	5.05	122.82	120.30
1	B	4	GLN	CA-C-N	-5.01	106.18	117.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2187	0	2199	48	0
1	B	2187	0	2199	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	0	0	0
4	A	27	0	12	3	0
4	B	27	0	12	1	0
All	All	4438	0	4422	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:CYS:HG	1:A:148:TYR:HE2	1.30	0.77
1:B:76:LEU:HD21	1:B:84:LYS:HB3	1.70	0.72
1:B:165:SER:HA	1:B:168:ILE:HD12	1.78	0.65
1:A:165:SER:HA	1:A:168:ILE:HD12	1.78	0.65
1:A:57:ILE:HD11	1:A:75:VAL:HG13	1.79	0.64
1:A:76:LEU:HD21	1:A:84:LYS:HB3	1.79	0.63
1:B:57:ILE:HD11	1:B:75:VAL:HG13	1.82	0.61
1:B:20:GLN:HE22	1:B:47:LEU:H	1.48	0.61
1:B:284:LYS:HD3	1:B:288:GLU:HB3	1.83	0.60
1:A:284:LYS:HD3	1:A:288:GLU:HB3	1.83	0.60
1:B:255:ILE:H	1:B:255:ILE:HD12	1.66	0.60
1:B:268:LEU:HB3	1:B:274:MET:HB2	1.83	0.60
1:A:92:GLU:HB3	1:A:95:VAL:HG12	1.85	0.59
1:B:213:ARG:HD2	4:B:292:ADP:C2	2.36	0.59
1:A:268:LEU:HB3	1:A:274:MET:HB2	1.85	0.59
1:A:213:ARG:HD2	4:A:291:ADP:C2	2.38	0.58
1:A:255:ILE:HD12	1:A:255:ILE:H	1.68	0.58
1:A:20:GLN:HE22	1:A:47:LEU:H	1.50	0.58
1:B:187:ARG:HA	1:B:213:ARG:NH1	2.20	0.56
1:B:92:GLU:HB3	1:B:95:VAL:HG12	1.88	0.56
1:B:238:ASP:HA	1:B:241:ARG:HB2	1.88	0.55
1:A:187:ARG:HA	1:A:213:ARG:NH1	2.22	0.55
1:A:238:ASP:HA	1:A:241:ARG:HB2	1.88	0.55
1:A:33:VAL:HG22	1:A:121:PHE:HB2	1.89	0.54
1:B:13:ILE:HD12	1:B:151:CYS:HA	1.90	0.54
1:B:279:GLU:O	1:B:282:VAL:HG22	2.08	0.53
1:B:27:ALA:HA	1:B:31:LYS:H	1.74	0.52
1:B:56:THR:HA	1:B:87:GLU:O	2.10	0.52
1:A:279:GLU:O	1:A:282:VAL:HG22	2.09	0.52
1:A:13:ILE:HD12	1:A:151:CYS:HA	1.92	0.51
1:B:105:ALA:O	1:B:109:LEU:HD23	2.11	0.50
1:B:33:VAL:HG22	1:B:121:PHE:HB2	1.94	0.50
1:A:49:LEU:HD13	1:A:85:CYS:SG	2.53	0.49
1:B:3:ARG:HH12	1:B:248:VAL:HA	1.78	0.49
1:A:56:THR:HA	1:A:87:GLU:O	2.13	0.49
1:A:27:ALA:HA	1:A:31:LYS:H	1.78	0.48
1:A:149:ILE:HG21	1:A:161:ALA:HA	1.93	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:ILE:HG22	1:A:289:VAL:HG21	1.95	0.48
1:B:211:VAL:HA	1:B:212:PRO:HD3	1.78	0.48
1:B:260:THR:HB	1:B:263:GLU:H	1.79	0.48
1:B:5:CYS:SG	1:B:146:GLU:HB2	2.53	0.48
1:B:281:ILE:HG22	1:B:289:VAL:HG21	1.95	0.47
1:B:148:TYR:CE1	1:B:180:GLY:HA3	2.50	0.47
1:A:284:LYS:O	1:B:223:ARG:HD3	2.14	0.46
1:A:260:THR:HB	1:A:263:GLU:H	1.80	0.46
1:B:155:MET:SD	1:B:264:LEU:HD21	2.55	0.46
1:A:105:ALA:O	1:A:109:LEU:HD23	2.16	0.46
1:A:95:VAL:HG23	1:B:171:TYR:OH	2.16	0.46
1:A:168:ILE:HD13	1:A:179:LEU:HB2	1.99	0.45
1:B:102:VAL:O	1:B:106:ILE:HG12	2.16	0.45
1:B:49:LEU:HD13	1:B:85:CYS:SG	2.57	0.45
1:B:235:LYS:O	1:B:239:GLU:HG2	2.16	0.45
1:A:3:ARG:HH12	1:A:248:VAL:HA	1.82	0.45
1:A:148:TYR:CE1	1:A:180:GLY:HA3	2.52	0.45
1:B:149:ILE:HG21	1:B:161:ALA:HA	1.97	0.45
1:A:182:LEU:O	1:A:208:ILE:HG22	2.17	0.44
1:A:49:LEU:CD1	1:A:85:CYS:SG	3.05	0.44
1:A:5:CYS:SG	1:A:146:GLU:HB2	2.57	0.44
1:B:187:ARG:HA	1:B:213:ARG:HH12	1.81	0.44
1:A:155:MET:SD	1:A:264:LEU:HD21	2.58	0.44
1:B:193:ASP:O	1:B:197:ILE:HG13	2.17	0.44
1:B:4:GLN:HB2	1:B:145:GLN:HE21	1.83	0.44
1:A:187:ARG:HA	1:A:213:ARG:HH12	1.82	0.43
1:A:10:LYS:O	1:A:13:ILE:HG12	2.18	0.43
1:B:168:ILE:HD13	1:B:179:LEU:HB2	2.00	0.43
1:B:162:ASN:O	1:B:165:SER:HB2	2.18	0.43
1:B:5:CYS:SG	1:B:148:TYR:HE2	2.41	0.43
1:A:235:LYS:HE3	1:A:235:LYS:HB3	1.87	0.43
1:A:162:ASN:HD21	1:A:259:ILE:H	1.67	0.43
1:B:66:THR:HG23	1:B:68:GLU:H	1.83	0.42
1:B:10:LYS:O	1:B:13:ILE:HG12	2.19	0.42
1:A:4:GLN:HB2	1:A:145:GLN:HE21	1.84	0.42
1:A:66:THR:HG23	1:A:68:GLU:H	1.84	0.42
1:A:193:ASP:O	1:A:197:ILE:HG13	2.20	0.42
1:B:103:ILE:HA	1:B:103:ILE:HD12	1.88	0.41
1:A:235:LYS:O	1:A:239:GLU:HG2	2.20	0.41
1:A:223:ARG:NE	1:B:281:ILE:HB	2.34	0.41
1:B:61:ALA:HA	1:B:70:LEU:HD11	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ILE:HG22	1:A:207:MET:HG3	2.02	0.41
1:A:236:GLN:NE2	1:A:236:GLN:HA	2.36	0.41
1:B:139:ILE:HG22	1:B:171:TYR:CD2	2.56	0.41
1:A:103:ILE:HA	1:A:103:ILE:HD12	1.91	0.41
1:A:14:GLY:HA2	4:A:291:ADP:O1A	2.21	0.41
1:A:159:TYR:O	1:A:162:ASN:HB3	2.21	0.40
1:A:14:GLY:HA2	4:A:291:ADP:PA	2.61	0.40
1:B:235:LYS:HE3	1:B:235:LYS:HB3	1.87	0.40
1:A:61:ALA:HA	1:A:70:LEU:HD11	2.03	0.40
1:A:54:GLN:HE22	1:A:86:VAL:HA	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/289 (99%)	262 (91%)	23 (8%)	2 (1%)	26	62
1	B	287/289 (99%)	262 (91%)	23 (8%)	2 (1%)	26	62
All	All	574/578 (99%)	524 (91%)	46 (8%)	4 (1%)	26	62

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	ASP
1	B	190	ASP
1	A	285	THR
1	B	285	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233 / 233 (100%)	214 (92%)	19 (8%)	14	38
1	B	233 / 233 (100%)	211 (91%)	22 (9%)	11	31
All	All	466 / 466 (100%)	425 (91%)	41 (9%)	12	35

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	66	THR
1	A	73	GLU
1	A	86	VAL
1	A	92	GLU
1	A	95	VAL
1	A	120	ASP
1	A	156	MET
1	A	178	ARG
1	A	187	ARG
1	A	189	THR
1	A	190	ASP
1	A	201	ASN
1	A	218	GLN
1	A	226	THR
1	A	235	LYS
1	A	238	ASP
1	A	241	ARG
1	A	255	ILE
1	B	22	LEU
1	B	66	THR
1	B	73	GLU
1	B	86	VAL
1	B	92	GLU
1	B	95	VAL
1	B	120	ASP
1	B	156	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	178	ARG
1	B	187	ARG
1	B	189	THR
1	B	190	ASP
1	B	191	ARG
1	B	201	ASN
1	B	218	GLN
1	B	226	THR
1	B	235	LYS
1	B	238	ASP
1	B	241	ARG
1	B	245	ARG
1	B	255	ILE
1	B	260	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	54	GLN
1	A	145	GLN
1	A	162	ASN
1	A	185	ASN
1	B	20	GLN
1	B	50	HIS
1	B	54	GLN
1	B	55	ASN
1	B	107	ASN
1	B	145	GLN
1	B	162	ASN
1	B	173	ASN
1	B	185	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SF4	A	290	1	0,12,12	0.00	-	0,24,24	0.00	-
4	ADP	A	291	-	22,29,29	1.71	2 (9%)	27,45,45	1.56	7 (25%)
4	ADP	B	292	-	22,29,29	1.96	3 (13%)	27,45,45	1.70	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	A	290	1	-	0/0/48/48	4/6/5/5
4	ADP	A	291	-	-	0/12/32/32	0/3/3/3
4	ADP	B	292	-	-	0/12/32/32	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	292	ADP	C3'-C4'	2.76	1.60	1.53
4	A	291	ADP	O4'-C1'	3.66	1.45	1.41
4	B	292	ADP	O4'-C1'	4.38	1.46	1.41
4	A	291	ADP	PB-O3B	5.63	1.74	1.54
4	B	292	ADP	PB-O3B	6.30	1.77	1.54

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	292	ADP	C4'-O4'-C1'	-3.31	106.08	109.72
4	B	292	ADP	C2'-C3'-C4'	-3.06	96.32	102.61
4	A	291	ADP	C4'-O4'-C1'	-2.91	106.53	109.72
4	A	291	ADP	C2'-C3'-C4'	-2.75	96.96	102.61
4	B	292	ADP	PA-O3A-PB	-2.74	123.47	132.67
4	A	291	ADP	PA-O3A-PB	-2.39	124.66	132.67
4	B	292	ADP	C2'-C1'-N9	-2.24	110.88	114.29
4	A	291	ADP	O3B-PB-O2B	-2.07	99.51	107.38
4	A	291	ADP	O4'-C1'-N9	2.04	112.36	108.10
4	A	291	ADP	O2A-PA-O3A	2.15	114.85	105.09
4	B	292	ADP	O2A-PA-O3A	2.26	115.35	105.09
4	A	291	ADP	O2B-PB-O3A	2.68	117.26	105.09
4	B	292	ADP	O2B-PB-O3A	2.81	117.82	105.09

There are no chirality outliers.

There are no torsion outliers.

All (4) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	290	SF4	FE3-FE4-S1-S2
3	A	290	SF4	FE2-FE4-S1-S3
3	A	290	SF4	FE1-FE2-S3-S4
3	A	290	SF4	FE1-FE3-S2-S4

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	291	ADP	3	0
4	B	292	ADP	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/289 (100%)	-0.01	3 (1%) 84 77	37, 82, 99, 100	0
1	B	289/289 (100%)	0.04	8 (2%) 56 44	41, 85, 100, 100	0
All	All	578/578 (100%)	0.01	11 (1%) 70 59	37, 83, 100, 100	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	226	THR	3.4
1	B	233	LYS	2.7
1	B	1	ALA	2.6
1	B	213	ARG	2.6
1	B	82	GLY	2.4
1	B	116	GLU	2.4
1	A	85	CYS	2.3
1	A	32	LYS	2.1
1	A	1	ALA	2.1
1	B	281	ILE	2.0
1	B	65	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ADP	B	292	27/27	0.94	0.15	-0.41	77,85,89,91	0
2	MG	B	294	1/1	0.80	0.12	-0.52	57,57,57,57	0
4	ADP	A	291	27/27	0.95	0.12	-1.04	66,78,88,90	0
3	SF4	A	290	8/8	0.99	0.08	-1.60	34,37,39,42	0
2	MG	A	293	1/1	0.95	0.07	-4.91	49,49,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.